



Contents lists available at ScienceDirect

Journal of Science and Medicine in Sport

journal homepage: [www.elsevier.com/locate/jsams](http://www.elsevier.com/locate/jsams)



## Review

# Effects of different protocols of high intensity interval training for VO<sub>2</sub>max improvements in adults: A meta-analysis of randomised controlled trials

Daizong Wen<sup>a,b</sup>, Till Utesch<sup>c</sup>, Jun Wu<sup>d</sup>, Samuel Robertson<sup>e</sup>, John Liu<sup>f</sup>, Guopeng Hu<sup>b,g</sup>, Haichun Chen<sup>a,\*</sup>

<sup>a</sup> School of Physical Education and Sport Science, Fujian Normal University, China

<sup>b</sup> Physical Education and Health Science Research Center, HuaQiao University, China

<sup>c</sup> Institute of Sport and Exercise Sciences, University of Münster, Germany

<sup>d</sup> Fujian Research Institute of Sports Science, China

<sup>e</sup> Institute of Sport Exercise and Active Living (ISEAL), Victoria University, Australia

<sup>f</sup> Department of Physical Education and Health Education, Springfield College, USA

<sup>g</sup> Department of Physical Education, HuaQiao University, China

## ARTICLE INFO

### Article history:

Received 18 September 2018

Received in revised form

24 December 2018

Accepted 22 January 2019

Available online xxx

### Keywords:

Cardiorespiratory fitness

Exercise

High-intensity intermittent exercise

Meta-analysis

## ABSTRACT

**Objectives:** To examine the effects of different protocols of high-intensity interval training (HIIT) on VO<sub>2</sub>max improvements in healthy, overweight/obese and athletic adults, based on the classifications of work intervals, session volumes and training periods.

**Design:** Systematic review and meta-analysis.

**Methods:** PubMed, Scopus, Medline, and Web of Science databases were searched up to April 2018. Inclusion criteria were randomised controlled trials; healthy, overweight/obese or athletic adults; examined pre- and post-training VO<sub>2</sub>max/peak; HIIT in comparison to control or moderate intensity continuous training (MICT) groups.

**Results:** Fifty-three studies met the eligibility criteria. Overall, the degree of change in VO<sub>2</sub>max induced by HIIT varied by populations (SMD = 0.41–1.81,  $p < 0.05$ ). When compared to control groups, even short-intervals ( $\leq 30$  s), low-volume ( $\leq 5$  min) and short-term HIIT ( $\leq 4$  weeks) elicited clear beneficial effects (SMD = 0.79–1.65,  $p < 0.05$ ) on VO<sub>2</sub>max/peak. However, long-interval ( $\geq 2$  min), high-volume ( $\geq 15$  min) and moderate to long-term ( $\geq 4$ –12 weeks) HIIT displayed significantly larger effects on VO<sub>2</sub>max (SMD = 0.50–2.48,  $p < 0.05$ ). When compared to MICT, only long-interval ( $\geq 2$  min), high-volume ( $\geq 15$  min) and moderate to long-term ( $\geq 4$ –12 weeks) HIIT showed beneficial effects (SMD = 0.65–1.07,  $p < 0.05$ ).

**Conclusions:** Short-intervals ( $\leq 30$  s), low-volume ( $\leq 5$  min) and short-term ( $\leq 4$  weeks) HIIT represent effective and time-efficient strategies for developing VO<sub>2</sub>max, especially for the general population. To maximize the training effects on VO<sub>2</sub>max, long-interval ( $\geq 2$  min), high-volume ( $\geq 15$  min) and moderate to long-term ( $\geq 4$ –12 weeks) HIIT are recommended.

© 2019 Sports Medicine Australia. Published by Elsevier Ltd. All rights reserved.

## Practical implications

- HIIT appears to be an effective alternative approach for improving VO<sub>2</sub>max in healthy, overweight/obese and athletic adults.
- Short-interval ( $\leq 30$  s), low-volume ( $\leq 5$  min) and short-term ( $\leq 4$  weeks) HIIT are feasible and time-efficient strategies and come with high effectiveness for improving VO<sub>2</sub>max, especially for the general population.
- Long-interval ( $\geq 2$  min), high-volume ( $\geq 15$  min) and moderate to long-term ( $\geq 4$ –12 weeks) HIIT protocols should be adopted, if the goal is to maximize the training effects on VO<sub>2</sub>max or surpass the MICT.

\* Corresponding author.

E-mail address: [chen.wen.dr@163.com](mailto:chen.wen.dr@163.com) (H. Chen).

## 1. Introduction

Aerobic capacity is typically measured as maximal oxygen uptake ( $\text{VO}_2\text{max}$ ). It is used frequently as an indicator of cardiorespiratory fitness, which is considered critical for health promotion.<sup>1</sup> Higher relative aerobic capacity levels are related to better physical performance of athletes,<sup>2</sup> and to a lower risk of cardiovascular/coronary heart diseases and all-cause mortality in non-athletic general population.<sup>3–5</sup> Recently, high intensity interval training (HIIT) was ranked Number 1 (most popular) in the annual survey of worldwide fitness trends in 2018.<sup>6</sup> It has been widely used as an alternative to traditional endurance training and was shown to result in higher levels of endurance performance,<sup>7</sup> reduced time commitment and increased exercise adherence.<sup>8</sup> Several studies have demonstrated the effectiveness of HIIT on  $\text{VO}_2\text{max}$  in athletes,<sup>9–11</sup> healthy<sup>12–14</sup> and overweight/obese non-athletes,<sup>15,16</sup> and even cardiac patients.<sup>17,18</sup> Moreover, new training programs were developed like the Resistance and Aerobic Program (RAP) that combines resistance exercises with HIIT. It has been shown that these are even more beneficial for improving physical and mental health outcomes in healthy and diabetic populations.<sup>19,20</sup>

HIIT protocols enable individuals who exercise to maintain at maximal or near maximal oxygen uptake ( $\text{T@VO}_2\text{max}$ ) for long periods of time, because a potent stimulus elicits both central (oxygen transport) and peripheral (oxygen utilization) adaptations for  $\text{VO}_2\text{max}$  improvement.<sup>7,21,22</sup> Many different components of HIIT such as work intensity, bout duration, number of repetitions, and training periodization have been shown to have substantial influence on  $\text{T@VO}_2\text{max}$ .<sup>7,21,22</sup> Correspondingly, HIIT can currently be subdivided into different protocols. For instance, according to different combinations of work intensity and bout duration, HIIT uses different work interval protocols including long-interval (2–4 min of work/bout at sub-maximal intensity, LI-HIIT), short-interval (<45 s of work/bout at sub-maximal intensity, SI-HIIT), sprint-interval (>20–30 s of work/bout at near to maximal intensity, SIT) and repeated-sprint exercises ( $\leq 10$  s of work/bout at near to maximal intensity, RST).<sup>21,23</sup> When the number of repetitions is added, HIIT protocols can be implemented with high (16 min of work) or low (4 min of work) session volume (HV-HIIT or LV-HIIT).<sup>16,24</sup> Moreover, considering the effect of training periodization, the length of HIIT intervention is classified as long-term ( $\geq 12$  weeks) or short-term ( $\leq 4$  weeks) duration (LT-HIIT or ST-HIIT).<sup>25,26</sup>

To increase time efficiency and exercise adherence, especially for non-athletes, HIIT training programmes were optimised with shorter work interval, lower session volume or shorter training periods.<sup>14,15,27,28</sup> However, these optimisations need to be further evaluated with respect to whether they retain a meaningful effect on improving  $\text{VO}_2\text{max}$  when compared with traditional HIIT programmes in diverse populations, because they can affect  $\text{T@VO}_2\text{max}$ .

An increasing body of systematic reviews and meta-analyses<sup>26,28–32</sup> have been conducted to investigate the efficiency of HIIT for improving  $\text{VO}_2\text{max}$  in adults without disease, and also examined the impact of several moderators of training effects. Batacan et al.,<sup>26</sup> Weston et al.,<sup>28</sup> Sloth et al.,<sup>29</sup> and Gist et al.<sup>30</sup> compared the effects of HIIT on  $\text{VO}_2\text{max}$  with moderate-intensity continuous training (MICT) and no training control groups. However, these meta-analyses included only HIIT research with short work intervals (10–30 s) or low session volumes ( $\leq 4$ –6 min). Milanović et al.<sup>31</sup> and Bacon et al.<sup>32</sup> addressed this gap in their meta-analyses that investigated HIIT protocols with longer work intervals (unrestricted) and higher session volumes (unrestricted or  $\geq 10$  min). However neither study directly examine the differences in  $\text{VO}_2\text{max}$  improvements between the particular protocols mentioned above nor involve athletic or overweight/obese populations. Furthermore, most of the above mentioned meta-analyses

included non-randomised controlled trials<sup>26,28,29,32</sup> and even non-control trials<sup>28,29,32</sup>, which may have led to potential bias or overestimation of treatment effects.<sup>33</sup>

In order to address such deficiencies, this systematic review and meta-analysis aims to review all relevant randomized controlled trials (RCTs), and examined the effects of HIIT on  $\text{VO}_2\text{max}$  improvements with regard to different work intervals, session volumes or training periods in several populations (i.e., healthy, overweight/obese and athletic adults).

## 2. Methods

This systematic review and meta-analysis was conducted according to the 'Preferred Reporting Items for Systematic Reviews and Meta-Analyses' (PRISMA) guidelines.<sup>34</sup> All the following steps were implemented by two independent raters (DW, JW), and any discrepancies were resolved by discussion or consensus with a third rater (HC).

A literature search was performed until April 2018 using the scientific databases (PubMed, Scopus, Medline, and Web of Science). The initial search terms included 'high intensity interval training' OR 'high intensity interval exercise' OR 'high intensity intermittent exercise' OR 'high intensity intermittent training' OR 'repeated sprint training' OR 'sprint interval training' OR 'HIIT' OR 'HIIE' OR 'HIT' OR 'HIE'. The second search terms included 'maximum  $\text{O}_2$ ' OR 'maximum  $\text{VO}_2$ ' OR 'maximal  $\text{VO}_2$ ' OR 'maximal oxygen uptake' OR 'maximal oxygen consumption' OR 'peak oxygen uptake' OR 'maximal aerobic capacity' OR ' $\text{VO}_2\text{max}$ ' OR ' $\text{VO}_2\text{peak}$ '. The third search terms included 'adult\*' OR 'men' OR 'women'. The fourth search terms included 'randomised controlled trial' OR 'RCT' OR 'random\*'. Finally, the four search terms were combined using the operator 'AND'. Further, reference lists of the included articles and related reviews were then scanned for potentially relevant studies.

Studies were identified using the following inclusion criteria: (1) adult participants including healthy (body mass index [BMI]  $< 25 \text{ kg/m}^2$ ) or overweight/obese (BMI  $\geq 25 \text{ kg/m}^2$ ) non-athletic or athletic populations (well-trained); (2) studies comparing HIIT with either control (CON) or MICT group, where training intensity thresholds of HIIT and MICT were defined as high ( $\geq 80$ – $85\%$   $\text{VO}_2\text{max}$ ,  $\geq 85$ – $90\%$  maximal heart rate [HRmax] or  $\geq 90\%$  velocity/power at  $\text{VO}_2\text{max}$  [ $\text{v/pVO}_2\text{max}$ ]) and moderate ( $40$ – $65\%$   $\text{VO}_2\text{max}$  or  $55$ – $75\%$  HRmax) respectively;<sup>35–37</sup> (3) studies of multiple treatment arms were treated as separate trials; (4) the training effect on  $\text{VO}_2\text{max}/\text{peak}$  was reported or could be calculated; (5) RCTs. Studies were excluded if they met the following criteria: (1) not published in English; (2) included participants suffering from any kind of acute or chronic diseases; (3) training intensity did not meet the previously defined thresholds; (4) HIIT was combined with other training methods in non-athletic populations (this criterion was not used for athletic research, as HIIT intervention is generally combined with the regular training programmes, which also served as the control group in the athletic research). To investigate the effect of various training protocols in this review, HIIT was pre-classified (Fig. 1) by different work intervals (long-interval [LI-HIIT], moderate-interval [MI-HIIT], short-interval [SI-HIIT], sprint-interval [SIT] and repeated-sprint [RST]), session volumes (high-volume [HV-HIIT], moderate-volume [MV-HIIT] and low-volume [LV-HIIT]) and training periods (long-term [LT-HIIT], moderate-term [MT-HIIT] and short-term [ST-HIIT]) respectively.<sup>16,21,23–26,29,38,39</sup>

We extracted the following characteristics from each eligible trial: author; year of publication; populations; sex; mean age; mean body mass index (BMI); baseline mean  $\text{VO}_2\text{max}/\text{peak}$




Fig. 1. Classification of HIIT protocols.

(ml/kg/min); groups; sample size; exercise modality; training period and frequency; training session protocol including number of repetitions, work intensity and duration, rest modality and duration, work/rest ratio and cumulative work time; and changes in  $\text{VO}_2\text{max/peak}$ . For the data that were shown only described in figures or graphs, we used Graph digitizer software (Digitizelt, Germany) to read the data. When the magnitude of changes in  $\text{VO}_2\text{max/peak}$  was not directly reported, we calculated the effect sizes and standard deviations (SDs) based on the baseline and pro-intervention values according to the methods suggested by the Cochrane handbook.<sup>40</sup> To assess the study quality, we used the modified Physiotherapy Evidence Base Database (PEDro) scale and considered a high quality study with a score of  $\geq 7/10$  points.<sup>41</sup> Additionally, three exercise training-specific criteria from the TESTEX scale<sup>42</sup> were added to the assessment, including activity monitoring in control groups, relative exercise intensity remained constant, and exercise energy expenditure information.

A meta-analysis was conducted to determine the pooled effect of the change in  $\text{VO}_2\text{max/peak}$  (ml/kg/min) for HIIT vs CON/MICT. Standardized mean difference (SMD), weighted mean difference (WMD) and 95% confidence intervals (CIs) were calculated using the random-effects model. A  $p$  value  $<0.05$  was considered statistically significant. The effect sizes are interpreted as trivial (SMD  $<0.2$ ), small (SMD 0.2-0.6), moderate (SMD 0.6-1.2), large (SMD 1.2-2.0) or very large (SMD 2.0-4.0).<sup>25,43</sup> Heterogeneity among studies was explored using Cochrane's  $Q$  statistic and  $I^2$  value, with values of 20%, 50% and 75% indicating low, moderate and high heterogeneity, respectively.<sup>44</sup> Sensitivity analysis was performed by removing trials with scores  $<7$  points (PEDro scale). To investigate different protocols of HIIT, further meta-analyses were performed by evaluating the effect of HIIT on  $\text{VO}_2\text{max/peak}$  by different work intervals (LI-SIIT, MI-SIIT, SI-SIIT, SIT and RST), session volumes (HV-HIIT, MI-HIIT and LV-HIIT) and training periods (LT-HIIT, MT-HIIT and ST-HIIT). Meta-regression analyses were further conducted in an attempt to determine the relationship between sex, age, BMI, baseline  $\text{VO}_2\text{max/peak}$  and the work:rest ratio with training effects on  $\text{VO}_2\text{max/peak}$ . Publication bias was analysed using funnel plot and Egger test.<sup>45</sup> All analyses were using executed using Stata version 13.1 (StataCorp, LP, College Station, TX).

### 3. Results

The initial search identified 1190 articles from the databases. Additionally, nine records were found via other sources. After excluding the duplicates, the titles and abstracts of 925 articles were screened. Of these, 251 eligible articles were selected for full-text review. Finally, a total of 53 records were included in this study (Supplementary material Fig. S1). Characteristics of the included studies are summarized in Supplementary material Table S1. All included studies compared the effectiveness of HIIT on  $\text{VO}_2\text{max/peak}$  with either CON or MICT group in a total of 1514 adults covering an age range of 19–47 years and baseline  $\text{VO}_2\text{max/peak}$  values ranging from 22.7 to 66.5 ml/kg/min. Populations covered by the identified studies included healthy non-athletes (26/53 studies with 29 HIIT groups), overweight/obese non-athletes (18/53 studies with 22 HIIT groups) and athletes (9/53 studies with 13 HIIT groups). Sample sizes in the HIIT groups ranged from 6 to 34 participants. Exercise modalities comprised cycling, handcycling, running, walking, swimming, and rowing. The HIIT protocols ranged from high (80%  $\text{VO}_2\text{max/peak}$ , 85%  $\text{HRmax}$  or 90%  $v/p\text{VO}_2\text{max}$ ) to all-out in intensity, 8 s to 10 min in bout duration and 20 s to 40 min in session volume. Training periods ranged from 2 to 16 weeks. The changes in  $\text{VO}_2\text{max/peak}$  after HIIT intervention varied between  $-5.4\%$  and  $33.1\%$ .

The methodological quality of the reviewed studies is presented in Supplementary material Table S2. A mean PEDro score of 6.77/10 (range from 5 to 9) was achieved. Concealed allocation (8%), blinding of assessors (28%), an explanation of sample size calculations (2%), activity monitoring in control groups (23%) and relative exercise intensity remained constant (38%) were reported in a minority of the studies, while specific eligibility criteria (74%), randomisation (100%), similar baseline groups (98%), outcome measures assessed in 70% of patients (98%), intention-to-treat analysis (70%), between-group statistical comparisons (100%), point measures and measures of variability (100%) and exercise energy expenditure information (92%) were reported in most of the studies.

The results of the overall and subgroup meta-analyses are presented in Table 1, Supplementary material Table S3 and Figs. S2-4, and the magnitude of effects for all protocols of HIIT were integrated and ranked in Fig. 2.

In healthy populations, HIIT had an overall large beneficial effect on  $\text{VO}_2\text{max/peak}$  (WMD = 5.45 ml/kg/min; SMD = 1.81, 95%

**Table 1**  
General results of the pooled effect of comparison HIIT versus CON/MICT on VO<sub>2</sub>max/peak by different HIIT training protocols.

	Healthy			I <sup>2</sup>	Overweight/obese			I <sup>2</sup>	Athletic			I <sup>2</sup>
	N	WMD (95%CI)	SMD (95%CI)		N	WMD (95%CI)	SMD (95%CI)		N	WMD (95%CI)	SMD (95%CI)	
<b>HIIT vs CON</b>												
Work intervals												
LI	8	7.62 (5.36, 9.88)	2.27 (1.60, 2.94)	70.7%	3	3.15 (0.35, 5.94)	1.13 (0.36, 1.90)	59.7%	5	2.46 (0.28, 4.65)	0.79 (0.05, 1.53)	66.2%
MI	2	3.72 (0.98, 6.46)	1.51 (0.60, 2.40)	42.3%	2	3.13 (-1.18, 7.43)	1.17 (-0.79, 3.13)	73.4%	2	2.44 (0.65, 4.23)	1.01 (0.22, 1.80)	0.0%
SI	3	3.54 (1.85, 5.23)	1.65 (0.77, 2.52)	35.8%					1	1.00 (-2.30, 4.30)	0.28 (-0.62, 1.18)	
SIT	5	3.66 (2.10, 5.23)	1.24 (0.41, 2.07)	72.1%	2	3.49 (-0.31, 7.28)	1.22 (-0.45, 2.89)	74.3%	4	0.21 (92.24, 2.66)	0.08 (-0.90, 1.06)	75.8%
RST	1	8.30 (5.73, 10.87)	2.31 (1.37, 3.25)		2	4.13 (3.18, 5.08)	1.97 (1.40, 2.54)	0.0%				
Training volumes												
HV	7	8.26 (5.91, 10.60)	2.48 (1.76, 3.19)	67.4%	1	2.00 (0.86, 3.14)	1.14 (0.43, 1.85)		5	2.87 (0.65, 5.08)	0.90 (0.16, 1.64)	65.9%
MV	6	4.09 (2.33, 5.86)	1.48 (1.04, 1.92)	17.5%	6	3.89 (2.59, 5.19)	1.43 (0.70, 2.16)	73.3%	5	2.56 (1.33, 3.79)	0.79 (0.35, 1.23)	0.0%
LV	6	3.96 (2.73, 5.18)	1.45 (0.62, 2.27)	73.9%	2	3.49 (-0.31, 7.28)	1.22 (-0.45, 2.89)	74.3%	3	90.87 (92.26, 0.53)	-0.45 (-0.99, 0.09)	0.0%
Training periods												
LT	4	5.64 (3.03, 8.24)	1.78 (1.32, 2.24)	71.3%	4	3.37 (1.87, 4.87)	1.68 (1.04, 2.33)	70.9%	8	-1.80 (-4.06, 0.46)	-0.60 (-1.37, 0.18)	67.5%
MT	11	6.01 (3.94, 8.08)	1.96 (1.39, 2.53)	67.3%	2	5.69 (3.90, 7.48)	1.99 (1.22, 2.76)	0.0%	4	1.31 (-0.26, 2.87)	0.50 (0.02, 0.99)	0.0%
ST	4	3.96 (2.47, 5.45)	1.56 (0.29, 2.83)	18.9%	3	2.38 (-0.08, 4.85)	0.76 (-0.09, 1.61)	51.9%	1	2.20 (0.48, 3.91)	0.79 (0.17, 1.41)	
<b>HIIT vs MICT</b>												
Work intervals												
LI	9	3.66 (2.28, 5.04)	1.07 (0.62, 1.52)	58.2%	5	2.51 (1.46, 3.56)	1.02 (0.64, 1.40)	3.6%				
MI	3	1.82 (-1.84, 5.48)	0.84 (-0.80, 2.48)	75.2%	3	0.92 (-0.22, 2.05)	0.29 (-0.11, 0.69)	0.0%				
SI					2	-1.15 (-4.19, 1.88)	-0.34 (-1.22, 0.55)	54.1%				
SIT	5	0.40 (-1.66, 2.45)	0.00 (-0.92, 0.93)	79.0%	3	-0.04 (-1.76, 1.69)	-0.08 (-0.99, 0.84)	67.5%				
RST	1	1.60 (-1.30, 4.50)	0.39 (-0.33, 1.12)		2	0.69 (-0.87, 2.26)	0.23 (-0.36, 0.82)	4.1%				
Training volumes												
HV	8	3.47 (1.99, 4.94)	1.04 (0.54, 1.54)	61.8%	5	2.51 (1.46, 3.56)	1.02 (0.64, 1.40)	3.6%				
MV	4	2.72 (-0.06, 5.51)	0.94 (-0.12, 2.00)	70.8%	4	0.69 (-0.31, 1.69)	0.22 (90.14, 0.57)	0.0%				
LV	7	0.20 (-1.53, 1.94)	-0.03 (-0.73, 0.67)	73.4%	6	-0.07 (-1.34, 1.21)	-0.04 (-0.59, 0.50)	57.1%				
Training periods												
LT	3	2.28 (0.11, 4.45)	0.77 (-0.08, 1.61)	51.1%	4	1.83 (0.82, 2.84)	0.77 (0.32, 1.22)	67.3%				
MT	12	2.25 (0.32, 4.18)	0.65 (0.05, 1.24)	79.0%	5	0.25 (-1.14, 1.64)	0.08 (-0.38, 0.54)	39.1%				
ST	4	1.20 (-1.24, 3.64)	0.43 (-0.46, 1.33)	76.9%	6	0.69 (-1.45, 2.83)	0.21 -0.66, 1.08)	50.0%				

The underlined data indicate statistically significant effect ( $p < 0.05$ ).

N: number of trails, HIIT: high intensity interval training, LI: long-interval ( $\geq 2$  min of work/bout at sub-maximal intensity), MI: moderate-interval ( $>30$  s and  $<2$  min of work/bout at sub-maximal intensity), SI: shot-interval ( $\leq 30$  s of work/bout at sub-maximal intensity), SIT: sprint interval training (10 to 30 s of work/bout at near to maximal intensity), RST: repeated sprint training ( $\leq 10$  s bout of work/at near to maximal intensity), HV: high-volume ( $\geq 15$  min of work/session), MV: moderate-volume ( $>5$  and  $<15$  min of work/session), LV: low-volume ( $\leq 5$  min of work/session), LT: long-term ( $\geq 12$  weeks), MT: moderate-term ( $>4$  and  $<12$  weeks), ST: short-term ( $\leq 4$  weeks), WMD: weighted mean difference, SMD: standardized mean difference, CL: confidence interval, CON: control group, MICT: moderate intensity continuous training.

CI 1.39–2.22,  $p < 0.05$ ,  $I^2 = 68.0\%$ ) in comparison to no training controls (NT-CON), while all HIIT protocols elicited significant beneficial effects (SMD = 1.24–2.48,  $p < 0.05$ ) in subgroup analyses. When compared to MICT, HIIT showed an overall moderate effect (WMD = 2.06 ml/kg/min; SMD = 0.64, 95% CI 0.23–1.05,  $p < 0.05$ ,  $I^2 = 75.2\%$ ) on VO<sub>2</sub>max/peak, but only long-interval, high-volume, and moderate-term protocols elicited significant beneficial effects (SMD = 0.65–1.07,  $p < 0.05$ ) in subgroup analyses.

In overweight/obese populations, HIIT had an overall large beneficial effect on VO<sub>2</sub>max/peak (WMD = 3.54 ml/kg/min; SMD = 1.35, 95% CI 0.81–1.88,  $p < 0.05$ ,  $I^2 = 68.8\%$ ) in comparison to NT-CON, while most HIIT protocols (long-interval, moderate to high-volume, moderate to long-term HIIT, and RST) elicited significant beneficial effects (SMD = 1.13–1.99,  $p < 0.05$ ) in subgroup analyses. When compared to MICT, HIIT showed an overall small effect (WMD = 1.07 ml/kg/min; SMD = 0.41, 95% CI 0.08–0.75,  $p < 0.05$ ,  $I^2 = 60.1\%$ ) on VO<sub>2</sub>max/peak, but only long-interval, high-volume, and long-term protocols elicited significant beneficial effects (SMD = 0.77–1.02,  $p < 0.05$ ) in subgroup analyses.

In athletic populations, HIIT had an overall small effect (WMD = 1.71 ml/kg/min; SMD = 0.57, 95% CI 0.13–1.01,  $p < 0.05$ ,  $I^2 = 62.8\%$ ) in comparison to regular training controls (RT-CON), while most HIIT protocols (moderate to long-interval, moderate to high-volume and short to moderate-term HIIT) elicited significant beneficial effects (SMD = 0.50–1.01,  $p < 0.05$ ) in subgroup analyses.

Sensitivity analysis (Supplementary material Table S3) demonstrated little less pronounced effects of HIIT on VO<sub>2</sub>max/peak in comparison to MICT became a little less pronounced after removing 16 trials of poor quality (PEDro scores  $< 7$  points). Potential publication bias was found by funnel plot (Supplementary mate-

rial Fig. S5) and Egger test ( $p = 0.011$ ). Regarding meta-regression analyses (Supplementary material Table S4), the work:rest ratio ( $\beta = 1.123$ ,  $p = 0.001$ ) was identified as a moderator for the effect of HIIT on VO<sub>2</sub>max/peak in overweight/obese populations when HIIT was compared to MICT.

#### 4. Discussion

This study utilised data from RCTs to confirm the findings from previous meta-analyses that examined the effectiveness of HIIT on VO<sub>2</sub>max performance. It also further investigated the effects of different protocols of HIIT in various populations. Overall, irrespective of protocol, the degree of change in VO<sub>2</sub>max induced by HIIT varied by populations. Further subgroup analyses revealed that even short work interval ( $\leq 30$  s), low-volume ( $\leq 5$  min) and short-term ( $\leq 4$  weeks) HIIT could elicit clear beneficial effects on VO<sub>2</sub>max when compared to CON. However, long-interval ( $\geq 2$  min), high-volume ( $\geq 15$  min) and moderate to long-term ( $\geq 4$ –12 weeks) HIIT displayed significantly larger effects on VO<sub>2</sub>max than both CON and MICT. Interestingly, when HIIT vs CON and HIIT vs MICT were both taken into consideration, training effects of long-interval and high-volume HIIT were highest in healthy populations, whereas long-term HIIT showed advantages in overweight/obese populations. For athletic adults, HIIT effects were lower with increased training periods, while in general population, the opposite was the case.

The current study found that non-athletic populations benefited more from HIIT than athletic populations, which is consistent with previous findings stating that aerobic training in general having an apparent adaptive effect on VO<sub>2</sub>max favouring the subjects




Fig. 2. Magnitude of effect by different protocols of HIIT in different population.

with a lower baseline  $VO_2\max$  value.<sup>28,31</sup> It is therefore unlikely that large improvements in  $VO_2\max$  could occur following HIIT in already highly trained athletes. This meta-analysis also found that HIIT appeared to be slightly more effective for healthy people than for overweight/obese people. It seems possible that this result is due to the calculation method of relative  $VO_2\max$  employed in the included studies, which divides absolute  $VO_2\max$  by body weight rather than fat-free mass (FFM).<sup>46,47</sup> Previous findings<sup>48,49</sup> have demonstrated that  $VO_2\max$  did not differ between obese and normal-weight people after adjusting for FFM, and  $VO_2\max$  was significantly correlated with FFM after controlling for fat mass. Therefore, without normalising  $VO_2\max$  by FFM, the training-induced changes in relative  $VO_2\max$  in obese subjects would be underestimated due to their higher body weight and body fat percentage.

In terms of the impact of work intervals on  $VO_2\max$ , previous meta-analyses<sup>29,30</sup> have demonstrated that SIT with 10–30 s sprints at all-out intensity demonstrated beneficial effects (SMD = 0.63–0.69) on  $VO_2\max$  levels compared to no training control groups, but a trivial effect (0.04) was observed when comparing it to endurance training in healthy adults, which is in line with the present study results. We also found that short-interval HIIT elicited similar training effects as SIT, but involved lower intensity with more repetitions. This means that although SIT was more time-efficient, short-interval HIIT could be an alternative approach when considering the safety and feasibility issues regarding the application of HIIT in general population.<sup>50</sup> Nevertheless, our findings show that both SIT and short-interval HIIT evoke no significant effect on  $VO_2\max$  in overweight/obese and athletic populations. Traditional moderate to long-interval HIIT between >30 s and 2 min

exercise at sub-maximal intensity are therefore recommended to ensure or enhance the training effect across all populations.

Recently, RST has received increased attention in the literature.<sup>38,51</sup> We observed large to very large effects on  $VO_2\max$  improvements in healthy and overweight/obese populations. However, previous studies suggested that RST with overly short bout durations may allow for a limited  $T@VO_2\max$  as compared to other HIIT protocols that involved longer intervals. It was considered to be more anaerobic dependent.<sup>21,29</sup> There were only four RCTs that used RST were identified in the present review. Hence high quality studies are needed to confirm our observations in the future.

Exercise volume as determined by work intervals and repetitions together was considered as a key factor that influences  $VO_2\max$  improvements and time-efficiency of a training program.<sup>1</sup> In accordance with the previous studies,<sup>26,29</sup> we found that low-volume HIIT elicited a large effect in healthy populations as compared to CON. However, only moderate to high-volume HIIT (>5–15 min) demonstrated moderate to very large effects across the populations when compared to CON or MICT. This finding was supported by Bækkerud et al.'s study<sup>52</sup> where high-volume HIIT (16 min) was superior to low-volume group in most likely improving the  $VO_2\max$  because of an increased stroke volume.

Moreover, We found that the session volume used in RST studies (8 min) was obviously larger than that used in most SIT studies (<4 min) due to more sprint repetitions employed in RST. This may be another reason why RST presented greater beneficial effects on  $VO_2\max$  changes in the included studies. A recent meta-analysis<sup>53</sup> investigating the effect of number of sprint repetitions in SIT showed that fewer repetitions would not attenuate the improvements in  $VO_2\max$ . However, their conclusion was limited, as the

session volumes employed were less than 5 min in all the included studies, suggesting that such a small range of change may not be enough to lead to significant increases in  $\text{VO}_2\text{max}$ . Therefore, we think that at a given individualized work interval, improvements in  $\text{VO}_2\text{max}$  could also be ensured or greatly enhanced across populations by substantially increasing the session volume.

Although a very short training duration (2 weeks) was considered to be sufficiently long to promote aerobic adaptations, a longer duration was more likely to be associated with greater improvements in  $\text{VO}_2\text{max}$ .<sup>16,54</sup> Our results demonstrated that even short-term HIIT ( $\leq 4$  weeks) can improve  $\text{VO}_2\text{max}$  when compared to CON in healthy populations, but moderate to long-term HIIT ( $> 4$ –12 weeks) showed additional further beneficial effects as compared to both CON and MICT in both healthy and overweight/obese populations. These findings are similar to those reported in a previous meta-analysis<sup>26</sup> where long-term HIIT ( $\geq 12$  weeks) exerted a large positive effect (SMD = 1.20) on  $\text{VO}_2\text{max}$  in overweight/obese populations. Thus, to ensure or more greatly enhance the training effects, it is important to improve exercise adherence and maintenance in general population, especially in overweight/obese populations.

Such positive trends were not observed in athletic populations, with HIIT displaying a reduced effect on  $\text{VO}_2\text{max}$  improvement over a prolonged intervention duration. This may indicate that the early stage of the training period is more likely to be responsible for the adaptations of  $\text{VO}_2\text{max}$  through HIIT in well-trained athletes.<sup>29</sup> However, this finding should be interpreted with caution as only one RCT used a short-term HIIT protocol on athletic populations were identified, and future work is required to confirm these results.

The present study does not come without limitations. The overall analysis demonstrated significant heterogeneity ( $I^2$  ranged from 60.1% to 75.2%) among the included studies, which may affect the findings of our meta-analysis. While pre-specified subgroup and meta-regression analyses were conducted to investigate the influence of some individual characteristic and training variables on training effect, varying degrees of heterogeneity ( $I^2$  ranged from 0.0% to 79.0%) were detected among results in subgroups, and only work:rest ratio was identified as a moderator for the effect of HIIT on  $\text{VO}_2\text{max}$  in the meta-regression analysis. This may have meant that the heterogeneity is affected by multi-factors that vary across studies rather than single factors. We therefore used the random effects model in the statistical analysis to make the results more conservative.

Although this review included published RCTs, many of these studies have suffered from small sample sizes with some issues in methodological quality, and a publication bias was detected, which may affect the reliability of our results. Moreover, due to the small number of trials included in some subgroup analyses, the findings should be interpreted with caution. Additionally, we extracted the relative values (ml/kg/min) rather than absolute values (L/min) of  $\text{VO}_2\text{max}$  from the included studies, which may in turn magnify the training effect due to a possible decrease of body weight after the intervention.

## 5. Conclusions

In conclusion, our meta-analysis suggests that, irrespective of protocol, HIIT is effective for improving  $\text{VO}_2\text{max}$  in healthy, overweight/obese and athletic adults. By investigating the different protocols of HIIT, short work interval HIIT ( $\leq 30$  s of work/bout at sub-maximal to all-out intensity), low-volume HIIT ( $\leq 5$  min of work/session) and short-term HIIT ( $\leq 4$  weeks of intervention) are feasible and time-efficient strategies and come with high effectiveness for  $\text{VO}_2\text{max}$  improvements, especially for the general

population. To ensure or more greatly improve the training effects on  $\text{VO}_2\text{max}$ , long-interval ( $\geq 2$  min of work/bout at sub-maximal intensity), high-volume ( $\geq 15$  min of work/session) and moderate to long-term ( $\geq 4$ –12 weeks of intervention) HIIT are recommended.

## Acknowledgments

This study was supported by Key Laboratory of Physical Motor Function Assessment, General Administration of Sport of China (ZGD1607235). Daizong Wen, Till Utesch, Jun Wu, Samuel Robertson, John Liu, Guopeng Hu, Haichun Chen have no conflicts of interest that are directly relevant to the content of this review.

## Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <https://doi.org/10.1016/j.jsams.2019.01.013>.

## References

1. Garber CE, Blissmer B, Deschenes MR et al. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: guidance for prescribing exercise. *Med Sci Sports Exerc* 2011; 43(7):1334–1359.
2. Helgerud J, Engen LC, Wisloff U et al. Aerobic endurance training improves soccer performance. *Med Sci Sports Exerc* 2001; 33(11):1925–1931.
3. Kodama S, Saito K, Tanaka S et al. Cardiorespiratory fitness as a quantitative predictor of all-cause mortality and cardiovascular events in healthy men and women: a meta-analysis. *JAMA* 2009; 301(19):2024–2035.
4. Blair SN, Kampert JB, Kohl 3rd HW et al. Influences of cardiorespiratory fitness and other precursors on cardiovascular disease and all-cause mortality in men and women. *JAMA* 1996; 276(3):205–210.
5. Lee DC, Artero EG, Sui X et al. Mortality trends in the general population: the importance of cardiorespiratory fitness. *J Psychopharmacol* 2010; 24(4 Suppl):27–35.
6. Thompson WR. Worldwide survey of fitness trends for 2018: the CREP edition. *ACSM's Health Fitness J* 2017; 21(6):10–19.
7. Laursen PB, Jenkins DG. The scientific basis for high-intensity interval training: optimising training programmes and maximising performance in highly trained endurance athletes. *Sports Med* 2002; 32(1):53–73.
8. Thum JS, Parsons G, Whittle T et al. High-intensity interval training elicits higher enjoyment than moderate intensity continuous exercise. *PLoS One* 2017; 12(1):e0166299.
9. Siahkhouhian M, Khodadadi D, Shahmoradi K. Effects of high-intensity interval training on aerobic and anaerobic indices: comparison of physically active and inactive men. *Sci Sports* 2013; 28(5):e119–e125.
10. Esfarjani F, Laursen PB. Manipulating high-intensity interval training: effects on  $\text{VO}_2\text{max}$ , the lactate threshold and 3000m running performance in moderately trained males. *J Sci Med Sport* 2007; 10(1):27–35.
11. Slettalokken G, Ronnestad BR. High-intensity interval training every second week maintains  $\text{VO}_2\text{max}$  in soccer players during off-season. *J Strength Cond Res* 2014; 28(7):1946–1951.
12. Boutcher SH, Park Y, Dunn SL et al. The relationship between cardiac autonomic function and maximal oxygen uptake response to high-intensity intermittent-exercise training. *J Sports Sci* 2013; 31(9):1024–1029.
13. Konigsrainer I, Zieker D, Loffler M et al. Influence of exhaustive exercise on the immune system in solid organ transplant recipients. *Exerc Immunol Rev* 2010; 16:184–193.
14. Hazell TJ, Macpherson RE, Gravelle BM et al. 10 or 30-s sprint interval training bouts enhance both aerobic and anaerobic performance. *Eur J Appl Physiol* 2010; 110(1):153–160.
15. Whyte LJ, Gill JM, Cathcart AJ. Effect of 2 weeks of sprint interval training on health-related outcomes in sedentary overweight/obese men. *Metabolism* 2010; 59(10):1421–1428.
16. Tjonna AE, Leinan IM, Bartnes AT et al. Low- and high-volume of intensive endurance training significantly improves maximal oxygen uptake after 10-weeks of training in healthy men. *PLoS One* 2013; 8(5):e65382.
17. Koufaki P, Mercer TH, George KP et al. Low-volume high-intensity interval training vs continuous aerobic cycling in patients with chronic heart failure: a pragmatic randomised clinical trial of feasibility and effectiveness. *J Rehabil Med* 2014; 46(4):348–356.
18. Rognum O, Hetland E, Helgerud J et al. High intensity aerobic interval exercise is superior to moderate intensity exercise for increasing aerobic capacity in patients with coronary artery disease. *Eur J Cardiovasc Prev Rehabil* 2004; 11(3):216–222.
19. Costigan SA, Eather N, Plotnikoff RC et al. High-intensity interval training for cognitive and mental health in adolescents. *Med Sci Sports Exerc* 2016; 48(10):1985–1993.
20. Wormgoor SG, Dalleck LC, Zinn C et al. High-intensity interval training is equivalent to moderate-intensity continuous training for short- and medium-term

- outcomes of glucose control cardiometabolic risk, and microvascular complication markers in men with type 2 diabetes. *Front Endocrinol (Lausanne)* 2018; 9:475.
21. Buchheit M, Laursen PB. High-intensity interval training, solutions to the programming puzzle: part I: cardiopulmonary emphasis. *Sports Med* 2013; 43(5):313–338.
  22. Midgley A, Mc Naughton L. Time at or near VO<sub>2</sub>max during continuous and intermittent running. *J Sports Med Phys Fitness* 2006; 46:1–14.
  23. Buchheit M, Laursen PB. High-intensity interval training, solutions to the programming puzzle Part II: anaerobic energy, neuromuscular load and practical applications. *Sports Med* 2013; 43(10):927–954.
  24. Bhati P, Bansal V, Moiz JA. Comparison of different volumes of high intensity interval training on cardiac autonomic function in sedentary young women. *Int J Adolesc Med Health* 2017; 0:1–13.
  25. de Oliveira MF, Caputo F, Corvino RB et al. Short-term low-intensity blood flow restricted interval training improves both aerobic fitness and muscle strength. *Scand J Med Sci Sports* 2016; 26(9):1017–1025.
  26. Batacan Jr RB, Duncan MJ, Dalbo VJ et al. Effects of high-intensity interval training on cardiometabolic health: a systematic review and meta-analysis of intervention studies. *Br J Sports Med* 2017; 51(6):494–503.
  27. Tabata I, Nishimura K, Kouzaki M et al. Effects of moderate-intensity endurance and high-intensity intermittent training on anaerobic capacity and VO<sub>2</sub>max. *Med Sci Sports Exerc* 1996; 28(10):1327–1330.
  28. Weston M, Taylor KL, Batterham AM et al. Effects of low-volume high-intensity interval training (HIT) on fitness in adults: a meta-analysis of controlled and non-controlled trials. *Sports Med* 2014; 44(7):1005–1017.
  29. Sloth M, Sloth D, Overgaard K et al. Effects of sprint interval training on VO<sub>2</sub>max and aerobic exercise performance: a systematic review and meta-analysis. *Scand J Med Sci Sports* 2013; 23(6):e341–e352.
  30. Gist NH, Fedewa MV, Dishman RK et al. Sprint interval training effects on aerobic capacity: a systematic review and meta-analysis. *Sports Med* 2014; 44(2):269–279.
  31. Milanovic Z, Sporis G, Weston M. Effectiveness of high-intensity interval training (HIT) and continuous endurance training for VO<sub>2</sub>max improvements: a systematic review and meta-analysis of controlled trials. *Sports Med* 2015; 45(10):1469–1481.
  32. Bacon AP, Carter RE, Ogle EA et al. VO<sub>2</sub>max trainability and high intensity interval training in humans: a meta-analysis. *PLoS One* 2013; 8(9):e73182.
  33. Stroup DF, Berlin JA, Morton SC et al. Meta-analysis of observational studies in epidemiology: a proposal for reporting: Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group. *JAMA* 2000; 283(15):2008–2012.
  34. Moher D, Liberati A, Tetzlaff J et al. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ* 2009; 339, b2535–b2535.
  35. Billat LV. Interval training for performance: a scientific and empirical practice: special recommendations for middle- and long-distance running. Part I: aerobic interval training. *Sports Med* 2001; 31:13–31.
  36. Weuge M, van den Berg R, Ward RE et al. The effects of high-intensity interval training vs: moderate-intensity continuous training on body composition in overweight and obese adults: a systematic review and meta-analysis. *Obes Rev* 2017; 18(6):635–646.
  37. Norton K, Norton L, Sadgrove D. Position statement on physical activity and exercise intensity terminology. *J Sci Med Sport* 2010; 13(5):496–502.
  38. Taylor J, Macpherson T, Spears I et al. The effects of repeated-sprint training on field-based fitness measures: a meta-analysis of controlled and non-controlled trials. *Sports Med* 2015; 45(6):881–891.
  39. Garcia-Hermoso A, Cerrillo-Urbina AJ, Herrera-Valenzuela T et al. Is high-intensity interval training more effective on improving cardiometabolic risk and aerobic capacity than other forms of exercise in overweight and obese youth? A meta-analysis. *Obes Rev* 2016; 17(6):531–540.
  40. Higgins JPT, Green S. *Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0 updated March 2011*, The Cochrane Collaboration, 2011. Table 7.7.a: Formulae for combining groups.
  41. Maher CG, Sherrington C, Herbert RD et al. Reliability of the PEDro scale for rating quality of randomized controlled trials. *Phys Ther* 2003; 83(8):713–721.
  42. Smart NA, Waldron M, Ismail H et al. Validation of a new tool for the assessment of study quality and reporting in exercise training studies: TESTEX. *Int J Evid Based Healthcare* 2015; 13(1):9–18.
  43. Hopkins WG, Marshall SW, Batterham AM et al. Progressive statistics for studies in sports medicine and exercise science. *Med Sci Sports Exerc* 2009; 41(1):3–13.
  44. Higgins JP, Thompson SG, Deeks JJ et al. Measuring inconsistency in meta-analyses. *BMJ* 2003; 327(7414):557–560.
  45. Egger M, Davey Smith G, Schneider M et al. Bias in meta-analysis detected by a simple, graphical test measures of funnel plot asymmetry. *BMJ* 1997; 315(7109):629–634.
  46. Toth MJ, Goran MI, Ades PA et al. Examination of data normalization procedures for expressing peak VO<sub>2</sub> data. *J Appl Physiol (1985)* 1993; 75(5):2288–2292.
  47. Vanderburgh PM, Katch FI. Ratio scaling of VO<sub>2</sub>max penalizes women with larger percent body fat, not lean body mass. *Med Sci Sports Exerc* 1996; 28(9):1204–1208.
  48. Ekelund U, Franks PW, Wareham NJ et al. Oxygen uptakes adjusted for body composition in normal-weight and obese adolescents. *Obes Res* 2004; 12(3):513–520.
  49. Goran M, Fields DA, Hunter GR et al. Total body fat does not influence maximal aerobic capacity. *Int J Obesity* 2000; 24(7):841–848.
  50. Bayati M, Farzad B, Gharakhanlou R et al. A practical model of low-volume high-intensity interval training induces performance and metabolic adaptations that resemble 'all-out' sprint interval training. *J Sports Sci Med* 2011; 10(3):571–576.
  51. Bishop D, Girard O, Mendez-Villanueva A. Repeated-sprint ability – part II: recommendations for training. *Sports Med* 2011; 41(9):741–756.
  52. Bækkerud FH, Solberg F, Leinan IM et al. Comparison of three popular exercise modalities on VO<sub>2</sub>max in overweight and obese. *Med Sci Sports Exerc* 2016; 48(3):491–498.
  53. Vollaard NBJ, Metcalfe RS, Williams S. Effect of number of sprints in an SIT session on change in VO<sub>2</sub>max: a meta-analysis. *Med Sci Sports Exerc* 2017; 49(6):1147–1156.
  54. Murias JM, Kowalchuk JM, Paterson DH. Time course and mechanisms of adaptations in cardiorespiratory fitness with endurance training in older and young men. *J Appl Physiol (1985)* 2010; 108(3):621–627.