

Review

Polyphenol Supplementation and Antioxidant Status in Athletes: A Narrative Review

Aleksandra Bojarczuk * and Magdalena Dzitkowska-Zabielska 

Faculty of Physical Education, Gdansk University of Physical Education and Sport, 80-336 Gdansk, Poland

* Correspondence: aleksandra.bojarczuk@awf.gda.pl

Abstract: Antioxidants in sports exercise training remain a debated research topic. Plant-derived polyphenol supplements are frequently used by athletes to reduce the negative effects of exercise-induced oxidative stress, accelerate the recovery of muscular function, and enhance performance. These processes can be efficiently modulated by antioxidant supplementation. The existing literature has failed to provide unequivocal evidence that dietary polyphenols should be promoted specifically among athletes. This narrative review summarizes the current knowledge regarding polyphenols' bioavailability, their role in exercise-induced oxidative stress, antioxidant status, and supplementation strategies in athletes. Overall, we draw attention to the paucity of available evidence suggesting that most antioxidant substances are beneficial to athletes. Additional research is necessary to reveal more fully their impact on exercise-induced oxidative stress and athletes' antioxidant status, as well as optimal dosing methods.

Keywords: dietary polyphenols; athletes; antioxidant status



Citation: Bojarczuk, A.;

Dzitkowska-Zabielska, M.

Polyphenol Supplementation and Antioxidant Status in Athletes: A Narrative Review. *Nutrients* **2023**, *15*, 158. <https://doi.org/10.3390/nu15010158>

Academic Editor: Barbara Fraczek

Received: 15 November 2022

Revised: 19 December 2022

Accepted: 22 December 2022

Published: 29 December 2022



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

According to the WHO, nutrition is crucial for health and development. Better nutrition is linked to improved health, stronger immunity, lower risk of non-communicable diseases, and longevity (<https://www.who.int/health-topics/nutrition>, accessed on 11 July 2022). Dietary supplements are different from conventional food. They are concentrated sources of vitamins, minerals, or other ingredients such as polyphenol extracts. They are available in various forms, including tablets, capsules, gummies, powders, drinks, and energy bars. Dietary supplements are sold in pharmacies or health and beauty retailers, which might mislead patients and/or customers into supposing they are forms of drugs or medicine. In addition, certain scientific challenges are encountered in supplement research due to the absence of consensus on terminology for the categorization and regulation of these products. Nevertheless, dietary polyphenol supplements are widely used by athletes due to their multiple biological effects. This review discusses polyphenols as a major area of interest within the field of sports due to their antioxidative properties [1] that are important for athletes because muscle exercise induces oxidative stress and promotes reactive oxygen species (ROS) production [2–4]. Oxidation-reduction homeostasis is a critical process for human health as it regulates a myriad of biological responses [5]. Thus, polyphenols are considered key signal molecules upon introduction into the human body [6]. Recent evidence shows that oral administration of polyphenols strengthens the body's antioxidant defense system and protects against oxidative stress in athletes [7,8]. However, the debate about polyphenols has gained fresh prominence with many researchers reporting disappointing outcomes [9–13]. Discrepancies in results remain, and the extent of the effects of polyphenols remains unclear [14]. The following review focuses on the use of polyphenols by athletes and aims to summarize current knowledge of the effects of dietary polyphenol supplementation on oxidative stress and oxidative damage. This review also includes details of polyphenol doses and exercise protocols.

1.1. What Are Polyphenols?

The term polyphenol is not well-defined. Surprisingly, the term was not found in the International Union of Pure and Applied Chemistry (UPAC) online databases nor in its glossaries [15,16]. Polyphenols are secondary metabolites of plants, meaning that they are synthesized through secondary metabolism and have no direct function in essential processes, i.e., photosynthesis or respiration [17]. However, they are involved in signal transduction from the root to the shoot [18]. For example, exogenously administered caffeic acid inhibits growth and enhances the lignification of soybean roots [19]. This indicates an allelopathic and allelochemical function of the acid. Allelochemicals released into the bulk soil might be absorbed by receptor plants [20], and can affect several plant processes, e.g., water utilization [21] or seed germination [22]. Interestingly, *Rhizobium* bacteria induce the synthesis of phenolic compounds in rice plants and rhizobial strains promote the growth and productivity of rice plants, highlighting a plant–microbe interaction [23]. Moreover, polyphenols are involved in the nutrient mobilization, e.g., by reducing soluble nitrogen in the soil [24]. The aforementioned examples outline the critical role of polyphenols in interactions with the general environment.

It is worth noting that these secondary metabolites are natural products, and they occur naturally in, e.g., fruits, vegetables, cereals, and beverages [25–27]. They are also produced during abiotic stresses [18,28] through the phenylpropanoid pathway [29,30]. One potential example is salt stress, upon which a plant's total phenolic content increases [31]. Salinity promotes ROS formation in plants, and this stress is mitigated by the plant's antioxidative system, which relies on polyphenols [32–35].

Phenolic compounds have powerful antioxidant properties and are helpful in scavenging harmful ROS in plants under salt stress [36,37]. From a chemical point of view, they are characterized by phenyl rings and hydroxyl substituents [38]. However, the literature is not consistent in terms of polyphenols' nomenclature and classification. Some publications describe two classes (flavonoids and non-flavonoids) [38,39], others three (phenolic acids, flavonoids, and non-flavonoids) [26,40], or four (phenolic acids, lignans, stilbenes, and flavonoids) [41–43]. Lignans (e.g., galbacin) are plant products of low molecular weight that consist of two propyl-benzene units coupled by β, β' bonds [44]. Phenolic acids (caffeic, gallic and coumaric acids) generally have one carboxylic acid group [45]. Flavonoids have a basic structural unit of 2-phenylchromone [15]. Stilbenes contain a 1,2-diphenylethylene backbone [46]. The schematic classification of polyphenols is presented in Figure 1.




Figure 1. Classification and chemical structure of major classes of dietary polyphenols. Adapted from [47,48].

1.2. Importance of Research on Polyphenol-Based Dietary Supplements

The global dietary supplements market increases in size every year. The market is driven by the rising popularity of sports-based and athletic performance enhancement supplements. While all regulatory scientists want to protect consumers from unsafe products, regulatory frameworks vary from country to country. A product might be considered a dietary supplement in one country, but prescription medicine in another. This means that pre-market research on polyphenol supplements is not mandatory worldwide. Scientific and regulatory challenges encountered in the existing research on the safety, quality, and efficacy of dietary supplements have been expertly reviewed elsewhere [49]. According to the National Institutes of Health (NIH), however, another problem lies in the fact that much of the research has focused only on single ingredients. Since dietary supplements for exercise and athletic performance contain multiple ingredients, a continued lack of data on the effectiveness and safety of the combinations in these multi-ingredient products is of high concern. A further complication is that most studies investigating the potential value and safety of supplements suggested to increase athletic performance involve mostly well-trained athletes. Therefore, there has been little discussion about their value to recreational or occasional exercisers. Moreover, many studies have involved young adults (more often males than females), and not adolescents who might want to use these dietary supplements (<https://ods.od.nih.gov/factsheets/ExerciseAndAthleticPerformance-HealthProfessional/#h2>, accessed on 11 July 2022). A further consideration relates to amounts of supplements in products. Some items provide neither the amount of each ingredient nor their standardization for, e.g., content of –polyphenols and anthocyanins. Only standardized extracts guarantee reliable clinical trials [50,51]. However, this practice

is rarely implemented worldwide. In addition, polyphenols are of low bioavailability [47], which leads to poor therapeutic effects and will be discussed separately below.

A large and growing body of literature has investigated polyphenol-based supplements. Some of the emerging issues relate specifically to the experimental models used. Cell lines are frequently used, e.g., L929 mouse fibroblasts [52], lung epithelial carcinoma cell line (A-549), human ovarian cancer cell line (A-2780), human breast cancer cell line (MCF-7), human colon carcinoma cell line (Caco-2) [53], or RAW 264.7 macrophages [54]. However, one could argue that in vivo models seem more convincing than in vitro because they involve complex living systems, encompassing the process, pathway, or function under study while part of a larger “living test tube” [55]. By contrast, data from animal studies have been criticized due to the physiological differences between animal and human metabolisms [56]. Still, a considerable volume of in vivo polyphenol research has been reported. Examples of studies on human objects are elegantly reviewed in [47]. In animal models, mammals are used most frequently because they are biologically very similar to humans in evolutionary terms [57]. However, first and foremost, data derived from athletes is required because there is currently insufficient experimental evidence to convincingly suggest intake of preparations based on polyphenol antioxidants.

2. Properties of Polyphenol-Based Dietary Supplements

Athletes experience physiological stress, which affects their whole performance. Effects include transient inflammation, oxidative stress, and immune dysregulation, which have been linked to nutritional influences [58]. The biological properties of polyphenols include antioxidant, anti-inflammatory, anti-cancer, anti-obesity, and anti-neurodegenerative effects, among more [59]. This review mainly describes the role of polyphenols in oxidative stress and inflammation, as these processes are linked.

2.1. Polyphenols, Oxidative Stress, and Inflammation

As mentioned above, polyphenols are antioxidants. This activity is ascribable to hydroxyl groups operating as electron donors and stabilizing free radicals through the delocalization of unpaired electrons [60], and is termed a chain-breaking function [61]. Polyphenols also demonstrate metal chelation activity. Because transition metals can produce ROS that damage the genome, polyphenols play an antioxidative and DNA-protective role [62]. In addition, they inhibit certain enzymes involved in ROS production, i.e., xanthine oxidase and NADPH oxidase, whilst upregulating other endogenous antioxidant enzymes, e.g., superoxide dismutase (SOD), catalase, and glutathione peroxidase (GPX) [63]. The antioxidative property of polyphenols is important for athletes, because it is well established that exercise generates ROS formation and prolonged exercise can promote oxidative damage to active myofibers [64–66]. Oxidative stress presents an imbalance between ROS-generating and -scavenging systems. ROS contain at least one oxygen atom and one or more unpaired electrons. ROS include free radicals (e.g., superoxide anion (O_2^-), hydroxyl radical ($\cdot OH$)) and non-radicals (e.g., singlet oxygen (1O_2), hydrogen peroxide (H_2O_2)). ROS overproduction can lead to tissue injury and initiate inflammation [63]. Superoxide (O_2^-) is relatively membrane-impermeable and unreactive [67] and has a short half-life in the cell [68]. Superoxide dismutase (SOD) spontaneously converts O_2^- to non-radical hydrogen peroxide (H_2O_2) [69] which is further reduced to water by GPX [70]. H_2O_2 molecule is considered stable and can therefore diffuse considerable distances in or out of the cell [2]. O_2^- and H_2O_2 are products of NADPH oxidase [71]. In contrast to H_2O_2 , the hydroxyl radical ($\cdot OH$) is a strong oxidizing agent and is considered the most damaging [72,73]. Another free radical is nitric oxide ($NO\cdot$), which is a short-lived and weak oxidant produced by the action of nitric oxide synthase (NOS) [74]. It has been shown that oxidative stress and inflammation are related, because inflammatory response occurs after exercise [75]. During inflammation, phagocytes undergo a respiratory burst that imposes high oxidative stress on the engulfed pathogens [76]. The oxidative burst is mediated by NADPH oxidase, which produces O_2^- and H_2O_2 [77], and these products

achieve their cytotoxic effects due to reactions with other antimicrobial systems to generate ROS [67,78]. In skeletal muscle cells in culture, two primary NADPH oxidase isoforms (NOX2 and NOX4) [79] were found to be the main sources of ROS [80,81]. Thus, the role of NADPH oxidase-derived ROS is important in pathogen-mediated and sterile inflammation (i.e., without pathogen involvement). Importantly, ROS can exacerbate the inflammatory process [82]. Nonetheless, free radicals are crucial for proper cell physiology and function actively as intracellular messengers [68,82]. This means that ROS have both positive and negative physiological effects. ROS are by-products of cellular metabolism, and constitutive antioxidant defenses neutralize the housekeeping production of reactive species, e.g., superoxide dismutase (SOD) converts O_2^- to H_2O_2 [83]. Another antioxidant strategy is the scavenging of ROS by numerous molecules of low molecular weight that exist in the extracellular space and also within cells. A further method is to minimize the availability of pro-oxidants via metal-binding proteins. These chelating molecules avert these transition metals from participation in ROS formation [2,84]. The toxic effects of ROS and RNS are also counteracted by non-enzymatic reactions, e.g., ascorbic acid or vitamin E [85]. However, when the production of radicals overwhelms the antioxidant defenses, oxidative stress proceeds. This can lead to changes in cell membranes and other biological structures including lipoproteins, lipids, proteins, DNA [86], etc. Polyphenols are of particular interest because they can also protect against oxidative damage through various mechanisms. For instance, resveratrol, which is a stilbenoid and a type of natural phenol found in red grapes, reduced the expression of NADPH oxidase 4 (NOX4) in mice [87]. Quercetin, a natural flavonoid found in onions, green tea, and apples, decreases lipid peroxidation and inhibits cellular oxidation in erythrocytes in type 2 diabetic patients [88]. Another flavonoid, myricetin also protected healthy human erythrocytes from oxidative stress *in vitro* [89]. The same effect was observed for tea catechins in human diabetic erythrocytes [90]. Conversely, others showed that catechins provided only modest protection from oxidative damage in healthy men [91] and showed no effect in soccer players [12]. Notwithstanding their high training demands, athletes' diets may not contain sufficient antioxidants to support their physical activity [92,93]. This raises the question of whether they should take dietary polyphenols. Given that the health benefits of exercise are well documented, it seems unlikely that exercise-generated free radicals should negatively impact performance in the long term [92,94].

2.2. Polyphenols Bioavailability

The bioavailability of a substance is the fractional extent to which the active moiety is absorbed and rate at which the dosage reaches the therapeutic site of action [95,96]. The issue of bioavailability presents a major challenge for clarifying the therapeutic effects of polyphenols and may account for large inter-individual variations in clinical trials [97]. Many factors influence the bioavailability of dietary polyphenols in humans including external factors (sun exposure, degree of ripeness of plants), factors related to food processing (homogenization, lyophilization), interactions with other compounds (bonds with proteins (e.g., albumin), polyphenol-related factors (i.e., chemical structure, concentration in food)) [98,99].

The *in vitro* approach to studying polyphenols' absorption, metabolism, and bioavailability has mostly used the Caco-2 cell line [100–103], because the line reliably reflects biological barriers and the ability to perform phase II biotransformation by adding hydrophilic groups. Nonetheless, other cells have also been used [104]. The *in vivo* approach mostly uses rodents [105–108] and humans [109–112]. Absorption of polyphenols in mammals occurs in the small intestine after cleaving off sugar moieties [113]. This process occurs by either passive diffusion or with the use of specific transporters localized in the enterocyte membrane [99]. Only approximately 5–10% is absorbed, and the remainder accumulates in the large intestine with the bile conjugates, being thereby subjected to digestion [114]. Importantly, polyphenols of low molecular weight such as gallic or caffeic acids are partially absorbed into the body directly or after conversion. In con-

trast, polyphenols of higher molecular weight, e.g., proanthocyanidins, are very poorly absorbed [115]. Phenolic compounds absorbed in the small intestine are further transported to the colon, where they are transformed by the gut microbiota. These microorganisms perform a myriad of activities including hydrolysis, dihydroxylation, demethylation, and decarboxylation [99] thereby converting phenolic compounds into bioavailable metabolites. Meanwhile, polyphenol-rich extracts can also modulate gut microflora [116,117]. Importantly, the position of the hydroxyl groups might influence the polyphenols' breakdown. For instance, flavonoids without hydroxyl groups at the C5, C7, and C4' positions are degraded more slowly [113,118]. This implies that the slow-degrading compounds might be more bioavailable because their absorption can occur more quickly than those degraded more rapidly at the colon level [119]. When polyphenol sub-products are absorbed by the small intestine or colon, phase I and II transformation begins within enterocytes and in the liver to be distributed through the blood [120–122]. However, the metabolites may encounter enterohepatic circulation, whereby they are returned to the small intestine and transported back to the liver [97,99,123]. Polyphenols can stay in the blood or be delivered to other tissues. Indeed, polyphenol metabolites have been found in various tissues, e.g., glucuronidated and methyl-glucuronidated derivatives of catechin and epicatechin in the muscle of rats fed with grape pomace extract [124] or catechin-glucuronide, methyl catechin-glucuronide and methyl catechin-sulfate in kidneys, intestine, lungs, spleen, and thymus of rats receiving hazelnut extract [125]. Nonetheless, their targeted delivery is a major problem relating to polyphenol-rich supplements and their low bioavailability. So far, many systems have been developed to address the issue. However, these approaches mainly concentrate on achieving delivery that minimizes systemic diffusion and degradation of phenolic compounds, because some phenolics are poorly absorbed in the digestive tract while others are extensively metabolized to derivatives with a lower activity or are degraded [99,126]. This means that when polyphenols are orally administered, their therapeutic concentration might not be achieved [127]. Temperature, light, oxygen, acidic pH, and enzymatic activity in the digestive system [127] can all affect the beneficial effects of polyphenols [128]. As mentioned above, polyphenols follow phase I and II metabolic pathways. The major pathway is phase II, during which glucuronidation and sulfation lead to especially hydrophilic conjugates, and methylation to similar if not slightly more lipophilic metabolites. The methylated products are conjugated into glucuronides and sulfates, as long as there is a functional group available for conjugation into more hydrophilic metabolites [120]. Moreover, unabsorbed and hydrophilic polyphenol conjugates can undergo transformation triggered by gut microflora, which can even result in ring fission. Bacteria can dramatically reduce polyphenols' bioactivity [129]. Therefore, various research efforts have been focused on the problem of polyphenol delivery. Of note, bioavailability is a key determinant of polyphenols' potential health-promoting applications [99], and is also important in establishing dietary reference intake (RDI) [130]. This aspect is reviewed in the discussion of polyphenol supplementation strategies.

2.3. The Intake of Phenolic Compounds in an Average Daily Diet

The biodiversity of polyphenols found in food is wide. Therefore, it is extremely difficult to determine their content within food products and measure their daily intake [131]. Several research articles have extrapolated the daily dietary intake of particular flavonoids among the European population, based on the US database of flavonoid concentrations in particular foods. Many authors have referred to the data published by Kühnau et al., almost 46 years ago, where a daily intake of 1 g of total phenols was established in the US population [132]. There have been attempts to define the daily consumption of polyphenols within the diets of different populations. So far, Hertog et al., have done this for flavonols and flavones [133], and Reinli and Block for isoflavones in the Dutch population [134]. Researchers established the intake of flavonols (mainly quercetin) and flavones as 21 and 2 mg/d, respectively [133]. The average dietary consumption of isoflavones in the Japanese population was determined to be 30–40 mg/d [135,136]. Interestingly, soy products are less

popular in Western countries, which is reflected in the lower polyphenol content of the Western diet [137], explaining why the daily dietary intake of quercetin and genistein does not exceed 2–4% of total polyphenols [131]. Some authors report that daily polyphenol consumption above 650 mg decreases risk of death in comparison with those whose daily polyphenols intake is below 500 mg [138]. Other authors report health benefits ranging from intakes of 500 mg to 1500 mg per day [131]. Further sources recommend a daily dose of 0.1–1.0 g of polyphenols. Fruits such as grapes, apples, pears, cherries, and blueberries contain up to 200–300 mg of polyphenols per 100 g of fresh weight. Interestingly, a glass of red wine or a cup of tea contains about 100 mg of polyphenols, and the presence of these products in the diet may reduce the likelihood of chronic diseases. In Europe, the main sources of polyphenols are onions, black tea, red wine, and apples [138,139].

2.4. Polyphenols Delivery

Various delivery systems for polyphenols have been developed to improve their efficiency and combat their bioactivity problems. One major issue is to find a way to enhance the penetration of active substances and bring hydrophilic compounds into the tissues. This concept has been explored in numerous studies demonstrating the use of formulations with simple emulsions [140–142], cyclic glucan oligosaccharides known as cyclodextrins [143–145], gels [146–148], nanoemulsion [149–151], or liposomes [152–154]. Other solutions are also available, such as micelles, nanocomposites, metal oxide nanoparticles, etc. [127]. The main reason for encapsulating polyphenols is to tackle the problem of their stability. For instance, maltodextrins preserve the integrity of anthocyanins [155]. Encapsulation increases biocompatibility and prevents degradation caused by the external environment. It also minimizes interactions with other components of the human body [156]. In general, coating by microencapsulation in particles up to 1000 μm protects active substances and preserves their antioxidant properties [157–159]. However, nanotechnology is apparently more effective, because nanoencapsulated polyphenols have been found to increase the protection of active substances and bioavailability as well as improving controlled targeted release [160]. Particle size is generally seen as a factor strongly related to bioavailability. The development of a nanoscale delivery system is aimed at achieving improved site-selective targeting [161]. This is possible due to the small molecular size and active incorporation into cells by different endocytic pathways [162,163]. However, targeted delivery is complicated. It can be achieved actively or passively [164]; active targeting involves the therapeutic agent (in this case a polyphenol) being loaded into a carrier and this conjugate attached to tissue or cell-specific ligands [165]. By contrast, passive targeting requires loading the therapeutic agent into a nanomolecule that passively reaches the target tissue or organ. This leads to the accumulation of a drug delivery system with a specific size, molecular mass, and charge [166]. Given that a passive targeting system might utilize specific conditions in the diseased tissues or cells (e.g., low pH) [166], it is reasonable to assume that this approach is suitable to treat oxidative stress generated by exercise. After all, the higher the exercise intensity, the lower the muscle pH [167]. Nonetheless, the active approach also seems effective since it is possible to target specific cells or the inside of cells (for targeting intracellular organelles) [166].

Importantly, approaches should be compatible with another target, that of achieving the concentrations required for systemic therapies. In these circumstances, the hydrophilic part of a phenol carrier must be in balance with the lipophilic part [168], because the affinity of polyphenols for lipid bilayers partially determines their biological activity *in vitro* [169] and is of great significance in the biomedical and dietary fields. Polyphenolic extracts interact with the cell membrane by creating a protective coat around the lipid membrane, through their location on the membrane surface. This effect was shown using liposomes as models of lipid membranes, wherein trans-stilbenes and flavonoids interacted at the hydrophilic interface [170]. This finding is in agreement with another study reporting that three different types of blueberry extracts changed the arrangement of the hydrophilic region of the liposome membranes [171]. Liposomes themselves possess an aqueous central

section as well as hydrophobic and hydrophilic components comprising a lipid bilayer. The aqueous cores typically encapsulate hydrophilic compounds. By contrast, hydrophobic substances favor lipid bilayers. Thus, liposomes might be used for the delivery of diverse substances, such as hydrophilic and hydrophobic compounds [172].

3. Oxidative Stress and Athletic Antioxidant Status

3.1. ROS in Working Muscles

Contracting muscles produce ROS, as shown in muscle tissue homogenates and intact muscles in rats [65]. While cellular respiration generates ROS, there is no hard evidence that mitochondria are the main source of ROS in contracting muscle fibers [173]. Early findings indicate that superoxide production in isolated rat or bovine mitochondria occurs at a level of 2–5% of total oxygen consumed by mitochondria [174,175]. Interestingly, recent results differ from that 2–5% estimate of superoxide generation; it was reported that less than 0.15% of oxygen consumed by the mitochondria is used in superoxide [176]. This was demonstrated using intact isolated rat mitochondria. These findings may disprove the proposal that mitochondria release significant amounts of reactive oxygen species under physiological conditions. In addition, in rat skeletal muscle fibers produce more ROS in the basal state of respiration compared with the active state [177]. It is therefore questionable whether mitochondria are the primary source of ROS production in the working muscle. Another debate concerns whether exercise-induced ROS generation is detrimental to health. This is somewhat doubtful, because if it were the case then people who engage regularly in sports would experience a higher prevalence of chronic diseases associated with oxidative stress. In support of this, the study by Walsh et al., demonstrated that endurance training increased VO_2max by 21% and did not increase the sensitivity of mitochondrial oxidative function to ROS in skinned fibers [178]. By contrast, sprinting relies mostly on anaerobic energy pathways [179] and high-intensity sprint training inhibits mitochondrial respiration [180]. Thus, exercise can be perceived as a double-edged sword, whereby moderate ROS production initiates positive physiological adaptation in the active skeletal muscles, and meanwhile high levels of ROS lead to damage in, e.g., proteins, lipids, and DNA [2]. Previous studies on high-level swimmers have reported that high-intensity discontinuous training and continuous moderate-intensity training produced similar levels of oxidative stress expressed by plasma biomarkers such as protein carbonyl, thiobarbituric acid-reactive substances, and 8-hydroxy-2-deoxy guanosine. The obtained results suggested the beneficial role of exercise independent of the intensity of training [181]. Likewise, although high-intensity discontinuous training induces ROS production, a gradual decrease of ROS and return to resting values is typically seen after 10 min. Adaptive responses are responsible for this phenomenon. The human body can expand its antioxidant capacity to restrain oxidative chain reactions and thereby control oxidative stress during exercise [182]. However, reactive species appear to act in a hormetic manner [181,183]. For example, subjects with low total antioxidant capacity (TAC) values prior to training showed an increase in these after exercise. Conversely, participants with high pre-training values showed a decrease [181]. It therefore seems likely that individual differences might occur in response to training.

Furthermore, the source of ROS during anaerobic exercise is unclear. Ex vivo muscle fiber data from mice shows that NADPH oxidase is one of the potential sites of superoxide, both at rest and during contractions in skeletal muscle [184]. Nevertheless, the measurement of ROS release by muscle mitochondria isolated after exercise from humans [185] and animals [186] suggests that aerobic exercise increases ROS production. It has been even suggested that a source other than the mitochondria initially releases the ROS, damaging the mitochondria and leading to their elevated ROS release [187]. Nonetheless, exercise leads to microinjury in muscle tissue [188,189], leading to increased tumor necrosis factor alpha ($\text{TNF}\alpha$) levels due to increased TNF expression by damaged muscle fibers in addition to macrophage infiltration [190–192]. Macrophages respond to changes in their cytokine microenvironment; briefly, they undergo polarization to either the M1 or M2 activation

phenotype following exposure to a specific stimulus [193]. Activated macrophages produce reactive oxygen species [194]. TNF α has been shown to interfere with oxidative phosphorylation in isolated rat and bovine mitochondria [195] and to stimulate ROS production [196]. Strenuous exercise also increases levels of serum platelet-derived growth factor (PDGF) [197], resulting in rapid ROS generation [198]. Plasma levels of Interleukin-6 (IL-6) depend on the intensity and duration of exercise, and IL-6 is sharply reduced when the muscle becomes more energetically efficient [199]. However, a pleiotropic Interleukin-6 (IL-6) dysregulates the muscle redox balance, as shown *ex vivo* in mice [200,201], but eventually reduces ROS production, which was demonstrated *in vitro* in the mouse myoblast cell line [202], whereas chronically elevated IL-6 was shown to contribute to oxidative stress in isolated mouse muscles [200,203]. Human *ex vivo* data indicate that the increase in pro-inflammatory cytokines is balanced by the release of anti-inflammatory cytokines such as Interleukin 1 receptor antagonist (IL-1Ra), IL-4 [204], and IL-10 [205]. A shift from pro- to anti-inflammatory signaling after muscle injury enables the termination of the inflammatory response and supports later phases of myogenesis, which was shown *ex vivo* in isolated mouse muscles [206]. This process is highly complex, as not only muscles and macrophages but many other cell types and signaling molecules are involved. Ultimately, rebalancing between pro- and anti-inflammatory cytokines prevents excessive ROS accumulation and guarantees full muscle rebuilding and return to redox homeostasis [207].

Interestingly, the appearance of ROS in damaged human-muscle fibres fosters muscle remodelling and their adaptation to physical activity [208,209] for two reasons: (1) ROS cooperates in activation of transcription factors, e.g., nuclear factor kappa-light-chain-enhancer of activated B cells (NF- κ B) [210], which regulates expression of several genes including cytokine genes. NF- κ B activation stimulated by contracting muscles provides a foundation for adaptation to exercise and regulates the expression levels of antioxidant enzymes in human skeletal muscle [211]; (2) ROS activate proteasomes [212] that degrade damaged proteins sensitive to ROS, i.e., actin. This eventually increases proteolysis and synthesis of proteins and intensifies the anabolic effect. In summary, ROS have divergent roles in humans. It should be highlighted that only a fully functioning antioxidant defense system maintains ROS at a safe level [213]. Normally they are generated at low levels and restricted to a specific subcellular location, thereby initiating a multitude of signaling cascades supporting normal physiological processes [214]. A schematic illustration of the possible representative pathways of polyphenol action in the muscle is depicted in Figure 2, with the example of lignans. Briefly, lignans decrease protein degradation, enhance protein synthesis, reduce oxidative stress, and improve mitochondrial biogenesis and myogenesis [215].




Figure 2. A schematic representation of the mechanism of polyphenols' action on muscles. Adapted from [215].

3.2. Physical Activity, Oxidative Stress and Antioxidant Status

Oxidative stress markers are molecules modified by interactions with ROS in the microenvironment. Quantifications can be classified into ROS detection, measurements of antioxidant levels (e.g., glutathione peroxidase (GPX), superoxide dismutase (SOD), catalase (CAT)), oxidation products (e.g., protein carbonyl (PC), thiobarbituric acid-reactive substance (TBARS) markers of protein oxidation, F₂-isoprostanes, and malondialdehyde (MDA), which are markers of lipid peroxidation, 8-oxo-2'-deoxyguanosine (8-OHdG), which is a marker of deoxyribonucleic acid (DNA) oxidation), or measurements of the redox balance (e.g., the reduced glutathione/oxidized glutathione (GSH/GSSG) ratio) [94]. Oxidative stress can be assayed by total antioxidant status (TAS) and the free radical scavenging capacities of plant extracts can be measured with methods such as 2,2-Diphenyl-1-picrylhydrazyl (DPPH), 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS), oxygen radical absorbance capacity (ORAC), total antioxidant capacity (TAC), etc.

Numerous publications highlight that oxidative stress and ROS generation sharply increase after the workout. For example, a short-term training period (repeated sprint) performed by elite soccer players led to elevations in TBARS, PC, SOD, and CAT [216]. Another study demonstrated that ultra-endurance running increases PC, TBARS, TAC, and 8-OH-dG immediately after testing in trained endurance athletes [217], whereas others reported no changes in TBARS, PC, and TAC observed during or after running [218]. Lower plasma malondialdehyde (MDA) was found in trained subjects compared to untrained, at rest 15 min after strenuous exercise [219]. However, contradictory findings have also been reported. Handball players undergoing 6 months of training were characterized by lower MDA results compared with non-athletes [220]. Similarly, treadmill runners maintained MDA at a stable levels over a 9-month program [221]. This suggests adaptation to regular exercise. Furthermore, changes in oxidative stress and redox status according to training load have been described. For instance, PC and TBARS were elevated in soccer players at the end of the pre-season training period, reduced in the middle of the season, and increased again at the end of the season. In contrast, antioxidant status biomarkers showed an opposite pattern of variations [222]. In another example, significant elevations of TBARS, PC, CAT activity, TAC, and also significant decreases in the GSH/GSSG ratio were found [223]. The ratio GSH/GSSG represents whole-body redox status [224] and its reduction is indicative of oxidative stress, as reported previously [225]. These results were observed in boys and girls participating in intense swimming training [223], and are in line with others who reported that antioxidant status and oxidative stress develop gradually over a sports season [226] and the redox status of professional athletes is influenced by training load throughout the season [227]. Although athletes' performances demonstrate various responses expressed as redox homeostasis markers, the data imply that short-term and long-term exercise are obvious stressors for the athlete's redox status, resulting in acute and chronic responses, respectively [228]. Moreover, oxidative stress and antioxidant status might relate to training intensity, as supported by the work of Lamprecht et al. This group examined regular aerobic exercisers and investigated the effects of single bouts of exercise at three different intensities on the redox state of human serum albumin, which acts as transport and redox system, and proposed that the redox pool was accessed more completely with increasing exercise intensity [229]. Furthermore, antioxidant status seems to relate to training status. In a study comparing untrained versus trained participants, the former showed (a) a larger increase in PC in response to exercise at all times throughout the training, (b) a more pronounced response from TBARS immediately and 1 h after training, (c) lower TAC values at rest and in response to exercise throughout the test period, and (d) lower GSH values at rest during the test period, and immediately and 1 h after exercise [230]. In addition, the literature indicates that the type of exercise might exert a different effect on the level of ROS-induced end products [3,231].

Lastly, antioxidant levels in the blood increase during physical exercise [232]. It also appears that these levels depend on the types of sport involved. Interestingly, as SOD activity increased, GPX and CAT activity also increased, and this finding was associated

with the physical condition of interval-trained athletes [233]. Changes in GPX, SOD, and GR confer a counteraction to radical production [231]. Ex vivo experiments suggest that regular training might increase the activity of antioxidant enzymes and reduce oxidant production [220,234].

Taken together, these results suggest antioxidant status is of great importance in terms of health maintenance. Despite a large volume of studies describing the negative aspects of exercise (ROS and oxidative stress), supportive data also exists and many studies continue to demonstrate the health benefits of exercise. These benefits include adaptation mechanisms, e.g., a rise in antioxidant capacity with increasing training effort as shown ex vivo in adolescent athletes [235]. Thus, it seems unlikely that oxidant production via exercise negatively impacts performance and health in the long term [92]. Although differences of opinion exist, it is widely accepted that exercise-induced radical production can damage skeletal muscle fibers and lead to fatigue. This knowledge has prompted many athletes to consume antioxidant supplements [93].

3.3. Physical Activity, Stress Hormones, Oxidative Stress and Antioxidant Status

Physical and psychosocial stressors activate the sympathomedullary pathway (SAM) and the hypothalamic–pituitary–adrenal axis (HPA axis) [236], and these routes trigger release of stress hormones. For instance, the SAM pathway leads to catecholamine (e.g., adrenaline and noradrenaline) secretion from the adrenal medulla, which are part of a short fight-or-flight response. The HPA pathway induces a longer transient hormonal cascade leading to glucocorticoids (cortisol) release from the adrenal cortex [237]. Researchers have studied the close relationship between stress hormones and athletic performance [238,239], and it has been found that circulating catecholamine levels increase during exercise in healthy athletes [240]. Catecholamines have beneficial effects on the heart, skeletal muscle, and central nervous system in relation to exercise performance [241]. Nonetheless, during pronounced oxidative stress in sustained or ongoing exercise, excess secretion of catecholamines contributes to immune suppression and a rise in ROS production due to catecholamine peroxidation [242–244]. Catecholamines released during exercise can also bind to β -adrenergic receptors, increasing sympathetic activity and alleviating skeletal muscle oxidative metabolism [245].

Competitive situations also change the levels of cortisol, the primary hormone of the HPA system, axis activation of which improves alertness for extended durations [246]. The paraventricular nucleus (PVN) of the hypothalamus releases corticotropin-releasing hormone (CRH). Adrenocorticotrophic hormone (ACTH) is then released from the anterior pituitary and acts on the adrenal cortex to release cortisol. According to Filaire et al., the typical cortisol response pattern exhibits an anticipatory spike before competition [247] and a second rise shortly after competition, as a result of extreme mental and physical strain [248]. The cortisol–performance relationship is inconsistent. For instance, there was no correlation between cortisol levels measured before multiple games of basketball and retrospective athlete and coach performance evaluations [249]. A negative correlation between performance and cortisol levels on the day of competition was observed during a 36-hole golf competition, suggesting that higher cortisol levels are linked to worse golfing performance [250]. In a study that utilized more time-points, cortisol levels in the loser of a tennis match were higher before (17%), during (65%), and after (54%) the game. Additionally, there was a negative correlation between cortisol and other performance indicators (e.g., unforced errors and return performance) [251], implying variations in cortisol. Another compound worth mentioning is 11 β -hydroxysteroid dehydrogenase (11 β -HSD), which catalyzes the conversion of cortisone to biologically active cortisol. According to Dovio et al. [252], strenuous exercise markedly elevates systemic 11 β -HSD type 1 activity in humans. In addition, their findings suggest that 11 β -HSD activity can profoundly upregulate cellular concentrations of cortisol. Therefore, it is plausible to suppose that increased local endogenous glucocorticoid availability might help to reduce exercise-induced muscle damage by helping to stop the inflammatory response in the

muscles [252]. Meanwhile, when cortisol levels are elevated, glucocorticoid-induced muscle atrophy might occur with the breakdown of contractile proteins. It was also reported that exercise-induced cortisol is linked with reactive oxygen species production [253]. Interestingly, antioxidant supplementation has been linked with reduced secretion of cortisol in athletes [254–256].

3.4. Dietary Polyphenols and Protocol Considerations

Oxidative stress is an active field of athletic research, and a wide range of different oxidative stress biomarkers has been used for its assessment. Since these markers typically result from diverse reactions, they also vary in how quickly they manifest and vanish following exposure [257]. Furthermore, oxidative stress indices also have drawbacks. For instance, in the TBARS assay *in vitro* and/or *ex vivo*, up to 98% of the measured MDA might be produced by the high temperature during the protocol itself [258]. TAC measures the total antioxidant capacity of plasma-bound small molecules, excluding peroxidation and the crucial role of intracellular enzymes including SOD or CAT. Thus, measurements cannot be taken as an accurate representation of *in vivo* antioxidant capacity [259]. An additional direct measurement technique for assessing ROS generation *ex vivo* is electron paramagnetic resonance [182,260], while the majority of studies use indirect markers of oxidative stress as surrogate markers [261]. These factors along with the pharmacokinetics of different dietary supplements must be considered when planning the blood-collection schedule of a study [257]. Thus, although a placebo-controlled crossover study design seems ideal and straightforward, many complex choices must be made, e.g., the length of pre-supplementation, timing of intake before testing, the dosage of the supplement, and the ideal washout period [257]. The gold standard for studying the effects of dietary supplements on athletic health should also include verification that the dietary supplement was ingested and elicited a biological response (e.g., by muscle, blood, urine, or saliva sampling) [262].

In elite athletes, it might be important to take into account the part of the season in which acute study was carried out. Overloading of exercise during the early part of the season might put additional stress on the body's immunological, muscular, and antioxidant systems [257,263–265]. The literature also points to an inter-individual variability of redox responses [266]. According to Lehmann et al., individual heterogeneity in recovery potential, exercise capacity, stress tolerance, and training tolerance accounts for the different ways in which athletes are vulnerable to overtraining under the same training conditions [267]. Even in subjects with the same training status, variations between individuals regarding the changes in oxidative stress markers were found *ex vivo* [268]. Inter-individual variability in response to consumption of polyphenols has also been reported [265]. Firstly, the bioavailability of polyphenols is impacted by inter-individual variability of absorption, distribution, metabolism, and excretion. Secondly, the target tissues' cellular molecular metabolism and the inter-individual variability of the cellular processes influenced by the bioavailable polyphenols (and/or their metabolites), are likely to have an impact on the bioactivity of these compounds [269]. A range of factors including genetic background, gut microbiota, age, gender, health, and training status may be involved in these inter-individual variations, but the current knowledge is sparse and fragmented. Another major problem in the literature investigating the effects of dietary polyphenols lies in the fact that many researchers did not use pure polyphenols or their mixtures, but blends with other antioxidants (e.g., juices containing polyphenols and vitamins [270–274]). Therefore, the antioxidant role of the active molecules could be ascribed not only to the polyphenols but also to, e.g., vitamins or the combined effect of polyphenols and vitamins, the latter also being found in berry juices. In other words, if there is no comprehensive information regarding the supplement content, i.e., if the supplement did not consist of polyphenols only, the experimental results cannot be attributed to polyphenols alone. This does not mean that the product to be tested cannot contain auxiliary substances or fillers. Nonetheless, the ideal dietary polyphenol supplement should be standardized to ensure the constant concentration of polyphenols,

their quality, and the therapeutic efficacy of each product without batch-to-batch variations. In contrast, unstandardized plant extracts contain variable amounts of active substances and use of the same dose does not always produce the same therapeutic effect. For instance, Lafay et al., studied the performance of elite male athletes given supplements of 400 mg of grape extract. This extract contained no purified polyphenol, and chemical analysis showed that gallic acid, catechin, and epicatechin were present in comparatively high amounts. The authors stated that the extract was from whole grapes, which means that the seeds might have been used [266]; these contain vitamin E, which is an antioxidant [275,276]. Therefore, the findings that grape extract improves antioxidant status and physical performance in elite male athletes do not seem to be polyphenol-specific. Another example can be found in the work Morillas-Ruiz et al. [270], who provided professional cyclists with a supplement in the form of a sports drink containing berry concentrates (total ingested polyphenol 2.3 g), vitamin C (20 mg/L), maltodextrin, pectin, and whey protein (unspecified quantities), as well as vitamin B1 (15% of recommended daily intake). Vitamin C was present in the placebo, which contained but no carbohydrates or other active ingredients. The fruit concentrate of the supplement drink (black grape [81 g/L], raspberry [93 g/L], and red currant [39 g/L]) provided almost all of its carbohydrates. The concentration of anthocyanin was highest (759 mg/L), followed by hydroxycinnamic acid derivatives (246 mg/L) and ellagic acid (168 mg/L) [270]. The authors concluded that the supplement protected against oxidative stress [270]. However, because the dietary supplement contained berry concentrates that provided polyphenols, carbohydrates, and vitamin C it is difficult to attribute these differences to polyphenols alone. An equivocal conclusion was also found in the study by Pilaczynska-Szczesniak et al., who supplemented male rowers with chokeberry juice (23 mg of anthocyanins in 50 mL of juice, three times per day for four weeks). Ex vivo analysis demonstrated that following a 2000 m ergometer rowing test, TBARS were dramatically decreased 24 h post-exercise in the supplemented group as opposed to those receiving the placebo [277]. The authors did not assess vitamin C content, found in aronia fruit which is a good source of vitamin C. The latter is well known for its antioxidative properties [278]. Thus, since the supplement might potentially have provided vitamin C as well as polyphenols, it is difficult to assign the differences between the supplement and placebo groups to the polyphenols alone.

As described, there are a number of methodological considerations regarding polyphenol supplementation in athletes, which can greatly affect experimental outcomes. Meanwhile, the optimal dosage and dietary reference intake remain undetermined. These issues are discussed in the supplementation strategies section, below (Section 3.5).

3.5. Polyphenols Antioxidant Intervention and Antioxidant Status in Athletes

Some athletes endorse dietary supplements based on data that supports improved exercise performance [279]. Studies have confirmed that polyphenols, i.e., quercetin [280], resveratrol [281], and polyphenolic compounds from grape extract [266] or beetroot juice [282–284] improve exercise performance. Quercetin was found to increase mitochondrial biogenesis and endurance capacity in mice ex vivo [285]. This was shown by increased mRNA expression of *PGC-1 γ* and *SIRT1*, as well as increases in cytochrome c concentration [285]. *PGC-1 γ* and *SIRT1* promote mitochondrial biogenesis [286]. Increased cytochrome c concentrations coincidence with other concentrations of mitochondrial enzymes involved in the electron transport chain (ETC), the tricarboxylic acid cycle (TCA), and the β -oxidation pathway, which increases mitochondrial capacity [285,287]. Mitochondrial biogenesis is important in sports performance because it expands the total mitochondrial volume and improves capacity to respond to contraction as cells increase their numbers of mitochondria [288]. Resveratrol, which has a similar structure to quercetin, also improves mitochondrial function and aerobic capacity in mice [281] and rats [289], as shown experimentally ex vivo.

The reason for the interest in polyphenols' antioxidants is the discovery that highly reactive chemical species can increase during exercise. Unfortunately, there are very limited

and contradictory studies on the effects of polyphenol supplementation on exercise-induced oxidative stress and antioxidant status in athletes [290,291]. In contrast, there is abundant literature on exercise in animals [292–296] or non-athletes [271,272,297–301]. In humans, blood is the most frequently analyzed biological sample, while a limited number of studies have examined skeletal muscle. For instance, 8-week supplementation with 500 mg/d hesperidin, a flavanone abundant in citrus fruits [302], was shown to reduce oxidized glutathione in amateur cyclists, indicating reduced oxidative stress. The effect was not seen in the placebo group [303]. This is in line with *in vitro* studies where hesperidin was demonstrated to reduce the by-products of lipid peroxidation, measured as TBARS in human erythrocytes [304] and MDA in human hepatic cells [305]. TBARS also decreased in cyclists after one-time consumption of a polyphenol blend including black grape, raspberry, and redcurrant (2000 mg polyphenols, including 1212 mg anthocyanins) [270]. Furthermore, 1 month of supplementation with 400 mg grape extract in elite athletes resulted in decreased creatine phosphokinase (CPK), a marker of skeletal muscle damage, supporting the idea of improved recovery [266]. It was also shown that oral administration of grape extract increased antioxidant capacity, while levels of SOD, CAT, and lipid peroxidation markers did not change between the placebo and treatment groups. However, significant results were obtained for isoprostanes and athletic performance in handball players. Although not all markers demonstrated the anticipated results, the data were encouraging. The authors suggested that the grape extract might be beneficial for maintaining the balance of oxidative stress and antioxidant status balance, enhancing physical performance in one category of athlete [266].

In studies of catechins, which represent flavanols, a single dose of 640 mg of green tea catechins in soccer players did not impair exercise-induced oxidative stress or muscle damage [12]. Similarly, Dean et al. concluded that 6-day intake of 270 mg epigallocatechin-3-gallate by male cyclists had no effect on TBARS [306]. However, data for dark chocolate, which is flavonoid-rich [307], appears very convincing. Consumption of 80 g/day for 2 weeks resulted in reduced plasma F₂-isoprostanes at exhaustion and after 1-h recovery compared with the placebo group [308]. That study produced results that corroborated the findings of Davison et al., who found that 100 g of dark chocolate 48 h before a cycling challenge significantly increased TAS [299]. Chronic intake of dark chocolate (40 g/day equivalent) was also able to reduce oxidative stress and muscle damage biomarkers during elite football players' training sessions [309]. The opposite was seen for Montmorency cherry (600 mg of phenolic compounds and 40 mg of anthocyanins 10 days before testing) in resistance-trained males. Following resistance exercise, markers of oxidative stress, lipid peroxidation, and antioxidant activity did not change significantly over time [310].

The redox-related effects of supplementation with a wide range of phenolic compounds on oxidative stress induced by severe physical exercise was the subject of a 2022 review by Kruk et al. These effects were thoroughly reviewed but the research largely focused on combined effects of polyphenols in sports and not specifically on antioxidative parameters. However, they listed 12 publications in the recent literature that demonstrated the reduction of oxidative stress markers or improvements in antioxidant capacity in professional sportsmen [290], although the list included sedentary and other non-athletes that are beyond the scope of this current review. Therefore, based on [290] and our own research, relevant positions in the literature are denoted that directly relate to polyphenol intervention and redox status in athletes [11,12,270–272,308,311], summarized in Table 1. The table does not refer only to polyphenol intake alone. For instance, the research reported by Jówko et al., 2015 [11] used green tea extract, which might contain antioxidant compounds including flavonoids, epigallocatechin-3-gallate, and vitamins. Further examples can be found [266,310,312–315] wherein Montmorency cherry extract was examined. None of these publications except from Levers et al., 2015 [310], mentioned whether the skin of the Montmorency cherry was used for preparation of the extract. The skin contains melatonin, which acts as an antioxidant [316], but none of these publications analyzed the extract for melatonin. This means that the effects described in Table 1 cannot be ascribed

to polyphenols alone. It is interesting to note that some of the included research reported significant drops in just a few oxidative stress markers while finding no change in others. In addition, there was variation in the doses used. Research has shown that selected antioxidant doses might not always be advantageous or effective. For instance, 4-day supplementation with 480 mg of resveratrol daily had no effect on oxidative stress when compared with a placebo in physically active men subjected to a cycling challenge [13]. Authors have emphasized the presence of a variety of antioxidants in human cells, some of which are enzymatic in character, such as CAT and SOD. Intracellular antioxidants often reach millimolar concentrations, whereas polyphenols' circulating concentrations normally do not exceed the low micromolar range [317].

The maximal daily dose of dietary polyphenols remains unknown because they have no established dietary reference intake value (DRI) [59]. This is because polyphenols are classified as nutritional supplements and no such value exists in the pharmacopoeia. Classically, DRIs include the recommended dietary allowance (RDA), adequate intake (AI), the tolerable upper intake level (UL), and the estimated average requirement (EAR). All of these values are linked; no RDA will be set if an EAR cannot be established. When evidence remains insufficient to develop an RDA, AI can be established. The UL is not intended to indicate a recommended level of intake (RDA or AI) but warns against a harmful excess of nutrients [318]. The DRIs apply to the healthy general population, which implies a gap in the knowledge regarding ill individuals. In addition, it might be that life stage and the subject's sex can affect DRIs. Nonetheless, these values for polyphenols have not been set. To complicate matters, there exist numerous polyphenols with diverse structural properties and it does not seem feasible to obtain sufficient data to establish a DRI for each [319]. Dietary recommendations for polyphenol intake should be based on bioavailability and bioactivity, taking into account inter-individual variability [269], and the DRIs should be based on from randomized placebo-controlled studies in humans [51,319].

Table 1. Effects of polyphenol-enriched supplementation on oxidative stress induced by heavy exercise training in athletes.

Author and Date, Sport	Participant Characteristics	Polyphenols Supplement	Exercise Protocol	Results	Conclusions
Jówko et al., 2015 [11], sprint	randomized double-blinded study; male sprinters aged 21.6 ± 1.5 years ($n = 16$);	green tea extract (980 mg of PP/day taken as four capsules daily) ($n = 8$) or placebo treatment ($n = 8$) for 4 weeks	double cycle sprint test on a bicycle ergometer with submaximal load up to 130–150 heartbeats/min	significantly increased blood MDA, TAC and SOD in the placebo group and CK activity in both groups tested after exercise; P intake increased resting TAC levels, decreased post-exercise SOD and MDA	treatment with P prevents exercise-induced oxidative stress, but prevention of exercise-induced muscle oxidative damage or improvement in sprint performance were not identified
Jówko et al., 2012 [12], soccer	randomized double-blinded placebo-controlled design; soccer players aged 22.9 ± 5.5 years ($n = 16$)	green tea PP (640 mg /day taken as two capsules) ($n = 8$) or placebo ($n = 8$) administered 2 h after breakfast	muscular endurance test: three sets of two strength exercises (bench press, back press, squat) to exhaustion with a load of 60% 1RM	significantly increased levels of TBARS, uric acid, TAS, CK and total catechins after exercise in both groups	green tea PP supplementation did not affect oxidative stress
Morillas-Ruiz et al., 2006, [270], cycling	controlled double-blinded clinical trial; male cyclists aged 23.6 ± 0.9 years ($n = 60$);	a beverage (black grape, raspberry, red currant) containing (2.3 g PP/trial) ($n = 30$) as opposed to placebo-treated ($n = 30$)	submaximal 90 min aerobic exercise on a bicycle ergometer at 70% VO_2 max	no significant changes in plasma TAS levels in either group after exercise; lower increases in CK and TBARS in the supplemented group compared with the control group; decreased content of carbonyl groups in the PP-treated group	supplementing with PP can protect against oxidative stress caused by physical exertion
Allgrove et al., 2011 [308]; cycling	randomized controlled trial in regularly exercising men aged 22 ± 4 years ($n = 20$)	40 g of dark chocolate containing 98.7 mg PP ($n = 10$), twice daily and once 2 h before training for 2 weeks, versus control group ($n = 10$)	cycling for 90 min with varying VO_2 max from 60 to 90% for 30 s every 10 min, followed by this activity to exhaustion at 90% VO_2 max	significantly lower blood levels of F2-isoprostans during fatigue and after 1 h recovery; oxidized LDL before and after exercise; increased FFA levels in the supplemented group; no significant effect on IL-6, IL-10, IL-1Ra, glucose, glucagon, insulin and cortisol levels, and time to exhaustion	dark chocolate supplementation reduced some oxidative stress markers and increased free fatty acid mobilization post-workout

Table 1. Cont.

Author and Date, Sport	Participant Characteristics	Polyphenols Supplement	Exercise Protocol	Results	Conclusions
Sadowska-Krepa et al., 2008 [311]; swimming	male physical education students ($n = 14$)	three capsules of 390 mg/day of red grape skin extract (188 mg/g PP plus 35 mg/g anthocyanidins) ($n = 9$) three times a day for 6 weeks versus placebo ($n = 5$)	interval swim test of moderate to high intensity (six repetitions of 50 m)	significantly reduced CK activity, increased GSH, uric acid, TAS in plasma and increased swimming performance, minor changes in antioxidant enzymes (SOD, CAT, GPX, GR)	red grape skin supplementation in sports training improved hemodynamic status and performance, and had little effect on antioxidant defense system activity
Bell et al., 2014 [312], cycling	randomised controlled trial; male trained cyclists aged 30 ± 8 years; ($n = 16$)	30 mL of Montmorency cherry concentrate (274 mg of anthocyanins) twice daily for 7 days ($n = 8$) as opposed to a placebo ($n = 8$)	a 109-min high-intensity, stochastic cycling trial on a cycle ergometer on days 5, 6, and 7	significant reduction of lipid hydroperoxides in the cherry concentrate	lipid peroxidation was reduced with Montmorency cherry concentrate supplementation
Bell et al., 2015 [313], cycling	randomised controlled trial; male trained cyclists aged 30 ± 8 years; ($n = 16$)	30 mL of Montmorency cherry concentrate (276 mg of anthocyanins) twice daily for 8 days ($n = 8$) as opposed to a placebo ($n = 8$)	a 109-min high-intensity, stochastic cycling trial on a cycle ergometer on day 5	no change in plasma markers of damage (CK and lipid hydroperoxides)	oxidative stress was attenuated with Montmorency cherry concentrate supplementation
Levers et al., 2015, running [310]	randomised controlled trial; resistance trained males aged 20.9 ± 2.6 years ($n = 23$)	480 mg of powdered Montmorency cherry ($n = 11$) versus placebo ($n = 12$) \times 1 per day for 10 days	10 sets of 10 back squats with a barbell at 70% 1RM	no change in plasma marker of oxidative damage (TBARS, MDA, TAS, SOD) compared with placebo	supplementation with Montmorency cherry powder did not change markers of oxidative stress, lipid peroxidation, or antioxidant activity
McCormick et al., 2016 [314], water polo	double-blinded crossover trial; trained male water polo players ($n = 9$) aged 18.6 ± 1.4 years	30 mL Montmorency cherry concentrate (274 mg anthocyanins) in the morning and 60 mL of the concentrate (547 mg anthocyanins) in the evening for 6 days; no data regarding the number of participants under treatment or taking placebo	simulated water polo team game	no change in plasma F2-isoprostanes between supplement and control groups	supplementing with cherry juice did not affect oxidative stress

Table 1. Cont.

Author and Date, Sport	Participant Characteristics	Polyphenols Supplement	Exercise Protocol	Results	Conclusions
Lafay et al., 2009 [266], jumping	randomized, double-blind crossover design; elite male athletes aged 21.6 ± 2 ($n = 9$)	400 mg/day of grape extract (2 capsule/day; total polyphenol content >90%; gallic acid equivalents not shown) for 1 month; no data regarding the number of participants taking a supplement or placebo	jumps	antioxidant capacity (ORAC) increased and no change in SOD or CAT, GPx in treated group versus control GPx decreased in the placebo	grape extract maintained a healthy balance between oxidative stress and antioxidant status throughout competition period
Nieman et al., 2013 [315], running	randomized, double-blinded crossover design; male or female runners aged 19–45 years ($n = 31$)	20 g blueberry polyphenol–soy protein complex (2136 mg/d gallic acid equivalents); ($n = 16$) twice a day for 17 days versus placebo ($n = 15$)	2.5 h/day running bouts	PC did not differ between groups; no change in F2-isoprostanes	no apparent benefit in the polyphenol–soy protein complex group

PP—polyphenol, PC—protein carbonyls, TAS—total antioxidant status, TAC—total antioxidant capacity, SOD—superoxide dismutase, CAT—catalase, CK—creatinine kinase, MDA—malonaldehyde, TBARS—thiobarbituric acid reactive substances, $VO_2\max$ —maximal oxygen intake, IL-6—interleukin-6, IL-8—interleukin-8, IL-10—interleukin-10, IL-12p70—interleukin-12 antibody, IL-1Ra—interleukin-1-receptor antagonist, IL-1—interleukin-1 β , TNF α —tumor necrosis factor alpha, LDH—lactate dehydrogenase, GPX—glutathione peroxidase, GR—glutathione reductase, GSH—glutathione, LDL—low density lipoproteins, FFA—free fatty acids, 1RM—one repetition maximum, HR—maximum heart rate; adapted from [290].

As described, a large body of literature has reported the antioxidative properties of dietary polyphenols *in vitro*, in animal models, and for non-trained participants (e.g., recreationally active, subjected to a training protocol only for the research purposes, or sedentary). These publications mostly describe the effects of mixtures of antioxidants, e.g., beverages containing polyphenols and vitamins (comprehensively reviewed in [257,259,290]). Thus, their results cannot be ascribed to polyphenols alone. It is beyond the scope of this current study to discuss data from non-athletes and concerning antioxidant supplementation strategies that are not based on dietary polyphenols alone. Unfortunately, there is limited evidence from athletes *per se* (i.e., professionals regularly engaged in sports) supplemented with pure polyphenols. This current review points out that the likely reasons for the conflicting results in human studies are the administration of antioxidant cocktails, limited information on the antioxidant dosage required for effectiveness, and different durations of antioxidant supplementation. Additionally, a cooperative network of cellular antioxidants is formed by interactions between exogenous antioxidants in food and endogenous antioxidants [320], which further complicates the interpretation of the research results.

A wide range of doses were used in the group of interest in this current review. Table 1 lists doses for acute and chronic supplementation with supplements containing nutritional polyphenols. For example, in [270] (Table 1), a drink with flavonoids and vitamin C was used, which does not meet the criteria for Table 2. Nonetheless, in this case the antioxidant effect was ascribed to polyphenols because the amount of vitamin C present in the beverage was the same as that found in the placebo [270]. Table 2 presents examples in athletes of acute and chronic supplementation with polyphenols alone, considering the redox-related effects only (as opposed to Table 1 that presents effects ascribable to polyphenol-enriched supplements). Contrary to expectations, discrepancies in the experimental outcomes can be seen, even when polyphenol-enriched supplements are considered (Table 1). However, this might be due to the assessment of oxidative stress using blood markers, which is by far the most commonly analyzed sampling method; these markers are not sufficient to determine the extent of oxidative stress in skeletal muscle [290]. The basic problem in the detection of free radicals lies in the fact that ROS have a short half-life and react quickly with components that control the redox state. The direct measurement of ROS levels is difficult and requires high accuracy and precision. Nonetheless, new approaches for its direct measurement are being developed [321], and procedures for less intrusive biopsies may become possible. Regarding TAS, some authors have even speculated that dietary flavonoids have a significant antioxidant effect *in vivo* that does not emerge from the flavonoids themselves, but from increased uric acid levels resulting from fructose metabolism after consumption of flavonoid-rich fruits [317].

Another issue is that the antioxidant activity of polyphenols appears to result from their structure [322,323]. Thus, it would be of great interest for athletes to know which classes of polyphenols are most beneficial for scavenging free radicals.

More recently, the use of polyphenols has been criticized as reviews carried out failed to demonstrate remarkable antioxidant activity in athletes [290,291]. In general, therefore, it seems that the current promotion of polyphenols as antioxidants and their supposed beneficial role in athletes' antioxidant status is not based on unequivocal results.

Table 2. Effect of dietary polyphenol supplementation on oxidative stress induced by heavy exercise training in athletes.

Author and Date, Sport	Participant Characteristics	Polyphenols Supplement	Exercise Protocol	Results	Conclusions
Decroix et al., 2017 [324], cycling	crossover trial; trained male cyclists aged 30 ± 3 years ($n = 12$)	900 mg cocoa flavanols 1.5 h and 3 h pre-exercise (low PP diet consumed 24 h pre-exercise); number of participants in the PP or placebo group not provided	two cycling time-trials (each lasting 30 min at 75% of peak power output) with a passive recovery period of 100 min	increased TAC (corrected for uric acid); no change in plasma MDA	acute cocoa flavanol intake had minimal effects on exercise-induced oxidative stress
Yarahmadi et al., 2014 [325], treadmill running	double-blinded clinical trial; male and female athletes ($n = 54$) aged 24.96 ± 7.37 and 22.82 ± 6.84 years	100 mg supplemental anthocyanin pills; a 28-day supplementation of a single dose of 10 mL/kg/day before the run; no data regarding the number of participants under treatment or taking placebo	incremental graded Bruce protocol	no change in CK; no change in LDH	supplementation with anthocyanin in athletes might improve some indices of performance rather than oxidative stress
McAnulty et al., 2013 [326], running	double-blinded crossover trial; trained males aged 18–40 years ($n = 14$)	120 mg resveratrol and 225 mg quercetin for 6 days and 240 mg resveratrol and 450 mg quercetin on day 7 immediately prior to exercise; number of participants in the PP or placebo group not provided	1 h run at a 3% grade, approximately 80% VO_2 max	post-exercise PC, FRAP, ORAC, and TEAC significantly increased but not affected by PP treatment; a significant reduction in the post-exercise F2-isoprostanes in the PP group	resveratrol and quercetin supplementation attenuates post-exercise oxidative stress by reducing F2-isoprostanes
Giuriato et al. 2022 [327], cycling	single-blinded, crossover trial; physically active males aged 22.3 ± 3.6 years ($n = 10$)	capsaicin supplements ($n = 3$); 0.957 mg capsaicin/tablet versus control fiber supplements; 2×390 mg of capsaicin capsules; number of participants in the PP or placebo group not provided	a maximal incremental test using an ergometer	redox-related effects not analyzed	antioxidant potential suggested due to attenuation of the development of peripheral fatigue

PP—polyphenol; CAT—catalase, CK—creatin kinase, LDH—lactate dehydrogenase, FRAP—ferric-reducing ability of plasma (FRAP), TEAC—trolox equivalent antioxidant capacity (TEAC), MDA—malondialdehyde, ORAC—oxygen radical absorptive capacity PC—protein carbonyls, PP—polyphenol, TAC—total antioxidant capacity, TBARS—thiobarbituric acid reactive substances, SOD—superoxide dismutase.

The relationship between oxidative stress and sport is highly complex. To encourage the upregulation of endogenous antioxidant defences, ROS must be released. Indeed, ROS are generated during cellular metabolism including muscle contractions. It would therefore be very useful to know to what extent athletes experience oxidative stress, as ROS generation has been observed to rise during exercise to a level that can outpace antioxidant defenses. To our knowledge, there are no rapid tests for ROS detection or oxidative stress available on the market. There are several diagnostic methods available to detect ROS or to measure antioxidant levels, but are time-consuming and require diagnosticians. Self-conviction about undergoing oxidative stress is based on an unbalanced diet and physical effort. In such situations, many athletes reach for dietary polyphenols. The failure of some of the antioxidants reviewed here to exhibit direct and significant redox-modulating actions in vivo might reflect their lack of efficacy due to low bioavailability. Most importantly, evidence is currently scarce on outcomes related to antioxidant status and the redox-modulating effects of polyphenols alone in vivo as opposed to polyphenol-enriched supplementation.

4. Conclusions and Future Perspectives

It is evident that professional athletes are familiar with the concept that exercise causes oxidative stress, which needs to be prevented by exogenous doses of antioxidant polyphenols. The use of antioxidant-polyphenol-rich dietary supplements can upregulate the endogenous antioxidant defence system, which in turn might have crucial consequences for preventing excessive oxidative damage and promoting recovery. However, according to the body of literature presented here, the use of polyphenols in the diet is debatable. The variability in research findings means the determination of the antioxidative effects of dietary polyphenols remains elusive. For instance, protocols are still variable and further systematically designed studies are required to strengthen the evidence. This research should include well-constructed trials in humans.

It appears that switching dietary polyphenols' status from nutritional supplement to medicine would change their entire management. Safety certificates, clinical trials, and further work would be required. Currently, polyphenols are considered foodstuffs and therefore do not require such restrictive regulations as medicinal substances or medicines. Thus, all unanswered issues and measures concerning the antioxidant action of dietary polyphenols in athletes should come within the remit of researchers.

Author Contributions: Conceptualization, A.B. and M.D.-Z.; methodology, A.B. and M.D.-Z.; investigation, A.B. and M.D.-Z.; writing—original draft preparation, A.B.; writing—review and editing, A.B. and M.D.-Z.; supervision, A.B. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Olszowy, M. What Is Responsible for Antioxidant Properties of Polyphenolic Compounds from Plants? *Plant Physiol. Biochem.* **2019**, *144*, 135–143. [[CrossRef](#)] [[PubMed](#)]
2. Powers, S.K.; Deminice, R.; Ozdemir, M.; Yoshihara, T.; Bomkamp, M.P.; Hyatt, H. Exercise-Induced Oxidative Stress: Friend or Foe? *J. Sport Health Sci.* **2020**, *9*, 415–425. [[CrossRef](#)] [[PubMed](#)]
3. Hadžović-Džuvo, A.; Valjevac, A.; Lepara, O.; Pjanić, S.; Hadžimuratović, A.; Mekić, A. Oxidative Stress Status in Elite Athletes Engaged in Different Sport Disciplines. *Bosn. J. Basic Med. Sci.* **2014**, *14*, 56–62. [[CrossRef](#)]
4. Radak, Z.; Powers, S.K. Introduction to Special Topic on Exercise and Oxidative Stress. *J. Sport Health Sci.* **2020**, *9*, 385. [[CrossRef](#)]
5. Le Gal, K.; Schmidt, E.E.; Sayin, V.I. Cellular Redox Homeostasis. *Antioxidants* **2021**, *10*, 1377. [[CrossRef](#)]
6. Sorrenti, V.; Fortinguerra, S.; Caudullo, G.; Burianni, A. Deciphering the Role of Polyphenols in Sports Performance: From Nutritional Genomics to the Gut Microbiota toward Phytonutritional Epigenomics. *Nutrients* **2020**, *12*, 1265. [[CrossRef](#)]

7. Hadi, A.; Pourmasoumi, M.; Kafeshani, M.; Karimian, J.; Maracy, M.R.; Hasan Entezari, M. The Effect of Green Tea and Sour Tea (*Hibiscus sabdariffa* L.) Supplementation on Oxidative Stress and Muscle Damage in Athletes. *J. Diet Suppl.* **2017**, *14*, 346–357. [[CrossRef](#)]
8. Panza, V.S.P.; Wazlawik, E.; Schütz, G.R.; Comin, L.; Hecht, K.C.; da Silva, E.L. Consumption of Green Tea Favorably Affects Oxidative Stress Markers in Weight-Trained Men. *Nutrition* **2008**, *24*, 433–442. [[CrossRef](#)]
9. Visioli, F.; de la Lastra, C.A.; Andres-Lacueva, C.; Aviram, M.; Calhau, C.; Cassano, A.; D'Archivio, M.; Faria, A.; Favé, G.; Fogliano, V.; et al. Polyphenols and Human Health: A Prospectus. *Crit. Rev. Food Sci. Nutr.* **2011**, *51*, 524–546. [[CrossRef](#)]
10. Gomez-Cabrera, M.C.; Ristow, M.; Viña, J. Antioxidant Supplements in Exercise: Worse than Useless? *Am. J. Physiol. Endocrinol. Metab.* **2012**, *302*, 476–477. [[CrossRef](#)]
11. Jówko, E.; Długołęcka, B.; Makaruk, B.; Cieśliński, I. The Effect of Green Tea Extract Supplementation on Exercise-Induced Oxidative Stress Parameters in Male Sprinters. *Nutr. Res.* **2015**, *54*, 783–791. [[CrossRef](#)]
12. Jówko, E.; Sacharuk, J.; Balasinska, B.; Wilczak, J.; Charmas, M.; Ostaszewski, P.; Charmas, R. Effect of a Single Dose of Green Tea Polyphenols on the Blood Markers of Exercise-Induced Oxidative Stress in Soccer Players. *Int. J. Sport Nutr. Exerc. Metab.* **2012**, *22*, 486–496. [[CrossRef](#)]
13. Tsao, J.P.; Liu, C.C.; Wang, H.F.; Bernard, J.R.; Huang, C.C.; Cheng, I.S. Oral Resveratrol Supplementation Attenuates Exercise-Induced Interleukin-6 but Not Oxidative Stress after a High Intensity Cycling Challenge in Adults. *J. Med. Sci.* **2021**, *18*, 2137–2145. [[CrossRef](#)]
14. Stankiewicz, B.; Cieślińska, M.; Kujawski, S.; Piskorska, E.; Kowalik, T.; Korycka, J.; Skarpańska-Stejnborn, A. Effects of Antioxidant Supplementation on Oxidative Stress Balance in Young Footballers- a Randomized Double-Blind Trial. *J. Int. Soc. Sport Nut.* **2021**, *18*, 44. [[CrossRef](#)]
15. Moss, G.P.; Smith, P.A.S.; Tavernier, D. Glossary of Class Names of Organic Compounds and Reactive Intermediates Based on Structure (IUPAC Recommendations 1995). *Pure Appl. Chem.* **1995**, *67*, 1307–1375. [[CrossRef](#)]
16. Liebecq, C. IUPAC-IUBMB Joint Commission on Biochemical Nomenclature (JCBN) and Nomenclature Committee of IUBMB (NC-IUBMB): Newsletter. *J. Mol. Biol.* **1998**, *275*, 527–537.
17. Kessler, A.; Kalske, A. Plant Secondary Metabolite Diversity and Species Interactions. *Annu. Rev. Ecol. Evol. Syst.* **2018**, *49*, 115–138. [[CrossRef](#)]
18. Sharma, A.; Shahzad, B.; Rehman, A.; Bhardwaj, R.; Landi, M.; Zheng, B. Response of Phenylpropanoid Pathway and the Role of Polyphenols in Plants under Abiotic Stress. *Molecules* **2019**, *24*, 2452. [[CrossRef](#)]
19. Bubna, G.A.; Lima, R.B.; Zanardo, D.Y.L.; dos Santos, W.D.; Ferrarese, M.d.L.L.; Ferrarese-Filho, O. Exogenous Caffeic Acid Inhibits the Growth and Enhances the Lignification of the Roots of Soybean (*Glycine Max*). *J. Plant Physiol.* **2011**, *168*, 1627–1633. [[CrossRef](#)]
20. L Weir, T.; Park, S.-W.; Vivanco, M.J. Biochemical and Physiological Mechanisms Mediated by Allelochemicals. *Curr. Opin. Plant Biol.* **2004**, *7*, 472–479. [[CrossRef](#)]
21. Cheng, F.; Cheng, Z. Research Progress on the Use of Plant Allelopathy in Agriculture and the Physiological and Ecological Mechanisms of Allelopathy. *Front. Plant Sci.* **2015**, *6*, 01020. [[CrossRef](#)] [[PubMed](#)]
22. Ghimire, B.K.; Ghimire, B.; Yu, C.Y.; Chung, I.M. Allelopathic and Autotoxic Effects of Medicago Sativa—Derived Allelochemicals. *Plants* **2019**, *8*, 233. [[CrossRef](#)] [[PubMed](#)]
23. Mishra, R.P.N.; Singh, R.K.; Jaiswal, H.K.; Kumar, V.; Maurya, S. Rhizobium-Mediated Induction of Phenolics and Plant Growth Promotion in Rice (*Oryza sativa* L.). *Curr Microbiol* **2006**, *52*, 383–389. [[CrossRef](#)]
24. Halvorson, J.J.; Schmidt, M.A.; Hagerman, A.E.; Gonzalez, J.M.; Liebig, M.A. Reduction of Soluble Nitrogen and Mobilization of Plant Nutrients in Soils from U.S Northern Great Plains Agroecosystems by Phenolic Compounds. *Soil Biol. Biochem.* **2016**, *94*, 211–221. [[CrossRef](#)]
25. Pandey, K.B.; Rizvi, S.I. Plant Polyphenols as Dietary Antioxidants in Human Health and Disease. *Oxid. Med. Cell. Longev.* **2009**, *2*, 270–278. [[CrossRef](#)] [[PubMed](#)]
26. Beconcini, D.; Felice, F.; Fabiano, A.; Sarmiento, B.; Zambito, Y.; Di Stefano, R. Antioxidant and Anti-Inflammatory Properties of Cherry Extract: Nanosystems-Based Strategies to Improve Endothelial Function and Intestinal Absorption. *Foods* **2020**, *9*, 207. [[CrossRef](#)] [[PubMed](#)]
27. Sakakibara, H.; Honda, Y.; Nakagawa, S.; Ashida, H.; Kanazawa, K. Simultaneous Determination of All Polyphenols in Vegetables, Fruits, and Teas. *Agric. Food Chem.* **2003**, *51*, 571–581. [[CrossRef](#)]
28. Kowalczewski, P.Ł.; Radzikowska, D.; Ivanišová, E.; Szwengiel, A.; Kačániová, M.; Sawinska, Z. Influence of Abiotic Stress Factors on the Antioxidant Properties and Polyphenols Profile Composition of Green Barley (*Hordeum vulgare* L.). *Int. J. Mol. Sci.* **2020**, *21*, 397. [[CrossRef](#)]
29. Gómez-Martínez, H.; Gil-Muñoz, F.; Bermejo, A.; Zuriaga, E.; Badenes, M.L. Insights of Phenolic Pathway in Fruits: Transcriptional and Metabolic Profiling in Apricot (*Prunus armeniaca*). *Int. J. Mol. Sci.* **2021**, *22*, 3411. [[CrossRef](#)]
30. Rossi, L.; Borghi, M.; Francini, A.; Lin, X.; Xie, D.Y.; Sebastiani, L. Salt Stress Induces Differential Regulation of the Phenylpropanoid Pathway in Olea Europaea Cultivars Frantoio (Salt-Tolerant) and Leccino (Salt-Sensitive). *J. Plant Physiol.* **2016**, *204*, 8–15. [[CrossRef](#)]
31. Sarker, U.; Islam, M.T.; Oba, S. Salinity Stress Accelerates Nutrients, Dietary Fiber, Minerals, Phytochemicals and Antioxidant Activity in Amaranthus Tricolor Leaves. *PLoS ONE* **2019**, *13*, e0206388. [[CrossRef](#)] [[PubMed](#)]

32. Cho, B.O.; Che, D.N.; Kim, J.S.; Kim, J.H.; Shin, J.Y.; Kang, H.J.; Jang, S.I. In Vitro Anti-Inflammatory and Anti-Oxidative Stress Activities of Kushenol C Isolated from the Roots of *Sophora Flavescens*. *Molecules* **2020**, *25*, 1768. [[CrossRef](#)] [[PubMed](#)]
33. Oke, I.M.; Ramorobi, L.M.; Mashele, S.S.; Bonnet, S.L.; Makhafole, T.J.; Eze, K.C.; Noreljaleel, A.E.M.; Chukwuma, C.I. Vanillic Acid–Zn(II) Complex: A Novel Complex with Antihyperglycaemic and Anti-Oxidative Activity. *J. Pharm. Pharmacol.* **2021**, *73*, 1703–1714. [[CrossRef](#)]
34. Zheng, W.; Wang, S.Y. Antioxidant Activity and Phenolic Compounds in Selected Herbs. *J. Agric. Food Chem.* **2001**, *49*, 5165–5170. [[CrossRef](#)]
35. Dumanović, J.; Nepovimova, E.; Natić, M.; Kuča, K.; Jačević, V. The Significance of Reactive Oxygen Species and Antioxidant Defense System in Plants: A Concise Overview. *Front. Plant Sci.* **2021**, *11*, 552969. [[CrossRef](#)]
36. Kiani, R.; Arzani, A.; Mirmohammady Maibody, S.A.M. Polyphenols, Flavonoids, and Antioxidant Activity Involved in Salt Tolerance in Wheat, *Aegilops Cylindrica* and Their Amphidiploids. *Front. Plant. Sci.* **2021**, *12*, 646221. [[CrossRef](#)]
37. Şirin, S.; Aslım, B. Determination of Antioxidant Capacity, Phenolic Acid Composition and Antiproliferative Effect Associated with Phenylalanine Ammonia Lyase (PAL) Activity in Some Plants Naturally Growing under Salt Stress. *Med. Chem. Res.* **2019**, *28*, 229–238. [[CrossRef](#)]
38. Singla, R.K.; Dubey, A.K.; Garg, A.; Sharma, R.K.; Fiorino, M.; Ameen, S.M.; Haddad, M.A.; Al-Hiary, M. Natural Polyphenols: Chemical Classification, Definition of Classes, Subcategories, and Structures. *J. AOAC Int.* **2019**, *102*, 1397–1400. [[CrossRef](#)]
39. Abbas, M.; Saeed, F.; Anjum, F.M.; Afzaal, M.; Tufail, T.; Bashir, M.S.; Ishtiaq, A.; Hussain, S.; Suleria, H.A.R. Natural Polyphenols: An Overview. *Int. J. Food Prop.* **2017**, *20*, 1689–1699. [[CrossRef](#)]
40. Majidinia, M.; Bishayee, A.; Yousefi, B. Polyphenols: Major Regulators of Key Components of DNA Damage Response in Cancer. *DNA Repair* **2019**, *82*, 102679. [[CrossRef](#)]
41. Koszowska, A.; Dittfeld, A.; Puzo, A.; Nowak, J. Polifenole w Profilaktyce Chorób Cywilizacyjnych. *Postępy Fitoter.* **2013**, *4*, 263–266.
42. Szaniawska, M.; Taraba, A.; Szymczyk, K. Structure, Properties and Application of Anthocyanins. *Eng. Sci. Technol.* **2015**, *2*, 63–78.
43. Zhou, Y.; Zheng, J.; Li, Y.; Xu, D.P.; Li, S.; Chen, Y.M.; Li, H.B. Natural Polyphenols for Prevention and Treatment of Cancer. *Nutrients* **2016**, *8*, 515. [[CrossRef](#)]
44. Rodríguez-García, C.; Sánchez-Quesada, C.; Toledo, E.; Delgado-Rodríguez, M.; Gaforio, J.J. Naturally Lignan-Rich Foods: A Dietary Tool for Health Promotion? *Molecules* **2019**, *24*, 917. [[CrossRef](#)]
45. Kumar, N.; Goel, N. Phenolic Acids: Natural Versatile Molecules with Promising Therapeutic Applications. *Biotechnol. Rep.* **2019**, *2019*, e00370. [[CrossRef](#)]
46. El Khawand, T.; Courtois, A.; Valls, J.; Richard, T.; Krisa, S. A Review of Dietary Stilbenes: Sources and Bioavailability. *Phytochem. Rev.* **2018**, *17*, 1007–1029. [[CrossRef](#)]
47. Di Lorenzo, C.; Colombo, F.; Biella, S.; Stockley, C.; Restani, P. Polyphenols and Human Health: The Role of Bioavailability. *Nutrients* **2021**, *13*, 273. [[CrossRef](#)]
48. Martin, K.R.; Appel, C.L. Polyphenols as Dietary Supplements: A Double-Edged Sword. *Nutr. Diet Suppl.* **2009**, *2*, 6403722. [[CrossRef](#)]
49. Dwyer, J.T.; Coates, P.M.; Smith, M.J. Dietary Supplements: Regulatory Challenges and Research Resources. *Nutrients* **2018**, *10*, 41. [[CrossRef](#)]
50. Bonati, A. How and Why Should We Standardize Phytopharmaceutical Drugs for Clinical Validation? *J. Ethnopharmacol.* **1991**, *32*, 195–197. [[CrossRef](#)]
51. Lyu, W.; Rodriguez, D.; Ferruzzi, M.G.; Pasinetti, G.M.; Murrugh, J.W.; Simon, J.E.; Wu, Q. Chemical, Manufacturing, and Standardization Controls of Grape Polyphenol Dietary Supplements in Support of a Clinical Study: Mass Uniformity, Polyphenol Dosage, and Profiles. *Front. Nutr.* **2021**, *8*, 780226. [[CrossRef](#)] [[PubMed](#)]
52. Truzzi, F.; Valerii, M.C.; Tibaldi, C.; Zhang, Y.; Abduazizova, V.; Spisni, E.; Dinelli, G. Are Supplements Safe? Effects of Gallic and Ferulic Acids on in Vitro Cell Models. *Nutrients* **2020**, *12*, 1591. [[CrossRef](#)] [[PubMed](#)]
53. Banach, M.; Khaidakov, B.; Korewo, D.; Węsierska, M.; Cyplik, W.; Kujawa, J.; Ahrné, L.M.; Kujawski, W. The Chemical and Cytotoxic Properties of *Sambucus Nigra* Extracts—A Natural Food Colorant. *Sustainability* **2021**, *13*, 12702. [[CrossRef](#)]
54. Banach, M.; Wiloch, M.; Zawada, K.; Cyplik, W.; Kujawski, W. Evaluation of Antioxidant and Anti-Inflammatory Activity of Anthocyanin-Rich Water-Soluble *Aronia* Dry Extracts. *Molecules* **2020**, *25*, 4055. [[CrossRef](#)] [[PubMed](#)]
55. Bojarczuk, A. *Insights into the Opportunistic Fungal Pathogen *Cryptococcus* and Neutrophilic Inflammation Using Zebrafish Models*; The University of Sheffield: Sheffield, UK, 2020.
56. Maloney, S.K.; Fuller, A.; Mitchell, D.; Gordon, C.; Michael Overton, J. Translating Animal Model Research: Does It Matter That Our Rodents Are Cold? *Physiology* **2014**, *29*, 413–420. [[CrossRef](#)]
57. Gandhi, G.R.; Vasconcelos, A.B.S.; Wu, D.T.; Li, H.-B.; Antony, P.J.; Li, H.; Geng, F.; Gurgel, R.Q.; Narain, N.; Gan, R.Y. Citrus Flavonoids as Promising Phytochemicals Targeting Diabetes and Related Complications: A Systematic Review of in Vitro and in Vivo Studies. *Nutrients* **2020**, *12*, 2907. [[CrossRef](#)]
58. Nieman, D.C.; Mitmesser, S.H. Potential Impact of Nutrition on Immune System Recovery from Heavy Exertion: A Metabolomics Perspective. *Nutrients* **2017**, *9*, 513. [[CrossRef](#)]
59. Cory, H.; Passarelli, S.; Szeto, J.; Tamez, M.; Mattei, J. The Role of Polyphenols in Human Health and Food Systems: A Mini-Review. *Front. Nutr.* **2018**, *5*, 87. [[CrossRef](#)]

60. Quideau, S.; Deffieux, D.; Douat-Casassus, C.; Pouységu, L. Plant Polyphenols: Chemical Properties, Biological Activities, and Synthesis. *Angew. Chem. Int. Ed. Eng.* **2011**, *50*, 586–621. [[CrossRef](#)]
61. Viglianisi, C.; Menichetti, S. Chain Breaking Antioxidant Activity of Heavy (S, Se, Te) Chalcogens Substituted Polyphenols. *Antioxidants* **2019**, *8*, 487. [[CrossRef](#)]
62. Lakey-Beitia, J.; Burillo, A.M.; La Penna, G.; Hegde, M.L.; Rao, K.S. Polyphenols as Potential Metal Chelation Compounds against Alzheimer's Disease. *J. Alzheimers Dis.* **2021**, *82*, S335–S357. [[CrossRef](#)]
63. Yahfoufi, N.; Alsadi, N.; Jambi, M.; Matar, C. The Immunomodulatory and Anti-Inflammatory Role of Polyphenols. *Nutrients* **2018**, *10*, 1618. [[CrossRef](#)] [[PubMed](#)]
64. Alessio, H.M.; Goldfarb, A.H.; Cutler, R.G. MDA Content Increases in Fast- and Slow-Twitch Skeletal Muscle with Intensity of Exercise in a Rat. *Am. J. Physiol.* **1988**, *255*, C874–C877. [[CrossRef](#)] [[PubMed](#)]
65. Davies, K.J.; Quintanilha, A.T.; Brooks, A.G.; Packer, L. Free Radicals and Tissue Damage Produced by Exercise. *Biochem. Biophys. Res. Commun.* **1982**, *107*, 1198–1205. [[CrossRef](#)] [[PubMed](#)]
66. Duthie, G.G.; Robertson, J.D.; Maughan, R.J.; Morrice, P.C. Blood Antioxidant Status and Erythrocyte Lipid Peroxidation Following Distance Running. *Arch. Biochem. Biophys.* **1990**, *282*, 78–83. [[CrossRef](#)]
67. Powers, S.K.; Ji, L.L.; Kavazis, A.N.; Jackson, M.J. Reactive Oxygen Species: Impact on Skeletal Muscle. *Compr. Physiol.* **2011**, *1*, 941–969. [[CrossRef](#)]
68. Phaniendra, A.; Jestadi, D.B.; Periyasamy, L. Free Radicals: Properties, Sources, Targets, and Their Implication in Various Diseases. *Indian J. Clin. Biochem.* **2015**, *30*, 11–26. [[CrossRef](#)]
69. Fukai, T.; Ushio-Fukai, M. Superoxide Dismutases: Role in Redox Signaling, Vascular Function, and Diseases. *Antioxid. Redox Signal* **2011**, *15*, 1583–1606. [[CrossRef](#)]
70. Wang, Y.; Hekimi, S. Understanding Ubiquinone. *Trends Cell Biol.* **2016**, *26*, 367–378. [[CrossRef](#)]
71. Amanso, A.; Lyle, A.N.; Griendling, K.K. NADPH Oxidases and Measurement of Reactive Oxygen Species. *Methods Mol. Biol.* **2017**, *1527*, 219–232. [[CrossRef](#)]
72. Ward, J.F.; Evans, J.W.; Limoli, C.L.; Calabro-Jones, P.M. Radiation and Hydrogen Peroxide Induced Free Radical Damage to DNA. *Br. J. Cancer Suppl.* **1987**, *55*, 105–112.
73. Ransy, C.; Vaz, C.; Lombès, A.; Bouillaud, F. Use of H₂O₂ to Cause Oxidative Stress, the Catalase Issue. *Int. J. Mol. Sci.* **2020**, *21*, 9149. [[CrossRef](#)]
74. Radi, R. Oxygen Radicals, Nitric Oxide, and Peroxynitrite: Redox Pathways in Molecular Medicine. *Proc. Natl. Acad. Sci. USA* **2018**, *115*, 5839–5848. [[CrossRef](#)]
75. Cerqueira, É.; Marinho, D.A.; Neiva, H.P.; Lourenço, O. Inflammatory Effects of High and Moderate Intensity Exercise—A Systematic Review. *Front. Physiol.* **2020**, *10*, 01550. [[CrossRef](#)]
76. Cavinato, L.; Genise, E.; Luly, F.R.; Domenico, E.G.D.; Del Porto, P.; Ascenzioni, F. Escaping the Phagocytic Oxidative Burst: The Role of SODB in the Survival of Pseudomonas Aeruginosa Within Macrophages. *Front. Microbiol.* **2020**, *11*, 326. [[CrossRef](#)]
77. Nauseef, W.M. Detection of Superoxide Anion and Hydrogen Peroxide Production by Cellular NADPH Oxidases. *Biochim. Biophys. Acta* **2014**, *1840*, 757–767. [[CrossRef](#)]
78. Vatansever, F.; de Melo, W.C.M.A.; Avci, P.; Vecchio, D.; Sadasivam, M.; Gupta, A.; Chandran, R.; Karimi, M.; Parizotto, N.A.; Yin, R.; et al. Antimicrobial Strategies Centered around Reactive Oxygen Species—Bactericidal Antibiotics, Photodynamic Therapy, and Beyond. *FEMS Microbiol. Rev.* **2013**, *37*, 955–989. [[CrossRef](#)]
79. Ferreira, L.F.; Laitano, O. Regulation of NADPH Oxidases in Skeletal Muscle. *Free Radic. Biol. Med.* **2016**, *98*, 18–28. [[CrossRef](#)]
80. Michaelson, L.P.; Shi, G.; Ward, C.W.; Rodney, G.G. Mitochondrial Redox Potential during Contraction in Single Intact Muscle Fibers. *Muscle Nerve* **2010**, *42*, 522–529. [[CrossRef](#)]
81. D'amico, A.; Cavarretta, E.; Fossati, C.; Borriore, P.; Pigozzi, F.; Frati, G.; Sciarretta, S.; Costa, V.; De Grandis, F.; Nigro, A.; et al. Platelet Activation Favours NOX2-Mediated Muscle Damage in Elite Athletes: The Role of Cocoa-Derived Polyphenols. *Nutrients* **2022**, *14*, 1558. [[CrossRef](#)]
82. Mittal, M.; Siddiqui, M.R.; Tran, K.; Reddy, S.P.; Malik, A.B. Reactive Oxygen Species in Inflammation and Tissue Injury. *Antioxid. Redox Signal* **2014**, *20*, 1126–1167. [[CrossRef](#)] [[PubMed](#)]
83. Snezhkina, A.V.; Kudryavtseva, A.V.; Kardymon, O.L.; Savvateeva, M.V.; Melnikova, N.V.; Krasnov, G.S.; Dmitriev, A.A. ROS Generation and Antioxidant Defense Systems in Normal and Malignant Cells. *Oxid. Med. Cell Longev.* **2019**, *2019*, 6175804. [[CrossRef](#)] [[PubMed](#)]
84. Powers, S.K.; Jackson, M.J. Exercise-Induced Oxidative Stress: Cellular Mechanisms and Impact on Muscle Force Production. *Physiol. Rev.* **2008**, *88*, 243–276. [[CrossRef](#)] [[PubMed](#)]
85. Landete, J.M. Dietary Intake of Natural Antioxidants: Vitamins and Polyphenols. *Crit. Rev. Food Sci. Nutr.* **2013**, *53*, 706–721. [[CrossRef](#)] [[PubMed](#)]
86. Rudrapal, M.; Khairnar, S.J.; Khan, J.; Dukhyil, A.B.; Ansari, M.A.; Alomary, M.N.; Alshabrimi, F.M.; Palai, S.; Deb, P.K.; Devi, R. Dietary Polyphenols and Their Role in Oxidative Stress-Induced Human Diseases: Insights into Protective Effects, Antioxidant Potentials and Mechanism(s) of Action. *Front. Pharmacol.* **2022**, *13*, 806470. [[CrossRef](#)]
87. Xian, Y.; Gao, Y.; Lv, W.; Ma, X.; Hu, J.; Chi, J.; Wang, W.; Wang, Y. Resveratrol Prevents Diabetic Nephropathy by Reducing Chronic Inflammation and Improving the Blood Glucose Memory Effect in Non-Obese Diabetic Mice. *Naunyn Schmiedeberg's Arch. Pharmacol.* **2020**, *393*, 2009–2017. [[CrossRef](#)]

88. Rizvi, S.I.; Mishra, N. Anti-Oxidant Effect of Quercetin on Type 2 Diabetic Erythrocytes. *J. Food Biochem.* **2009**, *33*, 404–415. [[CrossRef](#)]
89. Maurya, P.K.; Rizvi, S.I. Protective Role of Tea Catechins on Erythrocytes Subjected to Oxidative Stress during Human Aging. *Nat. Prod. Commun.* **2009**, *4*, 221–226. [[CrossRef](#)]
90. Rizvi, S.I.; Zaid, M.A.; Anis, R.; Mishra, N. Protective Role of Tea Catechins against Oxidation-Induced Damage of Type 2 Diabetic Erythrocytes. *Clin. Exp. Pharmacol. Physiol.* **2005**, *32*, 70–75. [[CrossRef](#)]
91. Jówko, E.; Sacharuk, J.; Balasińska, B.; Ostaszewski, P.; Charnas, M.; Charnas, R. Green Tea Extract Supplementation Gives Protection against Exercise-Induced Oxidative Damage in Healthy Men. *Nutr. Res.* **2011**, *31*, 813–821. [[CrossRef](#)]
92. Higgins, M.R.; Izadi, A.; Kaviani, M. Antioxidants and Exercise Performance: With a Focus on Vitamin e and c Supplementation. *Int. J. Environ. Res. Public Health* **2020**, *17*, 8452. [[CrossRef](#)]
93. Powers, S.K.; Sollanek, K.J. Endurance Exercise and Antioxidant Supplementation: Sense or Nonsense?—Part 1. *Sport Sci. Exch.* **2014**, *27*, 1–4.
94. Kawamura, T.; Muraoka, I. Exercise-Induced Oxidative Stress and the Effects of Antioxidant Intake from a Physiological Viewpoint. *Antioxidants* **2018**, *7*, 119. [[CrossRef](#)]
95. Kim, M.T.; Sedykh, A.; Chakravarti, S.K.; Saiakhov, R.D.; Zhu, H. Critical Evaluation of Human Oral Bioavailability for Pharmaceutical Drugs by Using Various Cheminformatics Approaches. *Pharm. Res.* **2014**, *31*, 1002–1014. [[CrossRef](#)]
96. Chow, S.-C. Bioavailability and Bioequivalence in Drug. *Wiley Interdiscip. Rev. Comput. Stat.* **2014**, *6*, 304–312. [[CrossRef](#)]
97. Teng, H.; Chen, L. Polyphenols and Bioavailability: An Update. *Crit. Rev. Food Sci. Nutr.* **2019**, *59*, 2040–2051. [[CrossRef](#)]
98. D’Archivio, M.; Filesì, C.; Vari, R.; Scaccocchio, B.; Masella, R. Bioavailability of the Polyphenols: Status and Controversies. *Int. J. Mol. Sci.* **2010**, *11*, 1321–1342. [[CrossRef](#)] [[PubMed](#)]
99. Cosme, P.; Rodríguez, A.B.; Espino, J.; Garrido, M. Plant Phenolics: Bioavailability as a Key Determinant of Their Potential Health-Promoting Applications. *Antioxidants* **2020**, *9*, 1263. [[CrossRef](#)]
100. CenciĀ, A.; Langerholc, T. Functional Cell Models of the Gut and Their Applications in Food Microbiology—A Review. *Int. J. Food Microbiol.* **2010**, *141* (Suppl. 1), S4–S14. [[CrossRef](#)] [[PubMed](#)]
101. Li, Q.; Wang, C.; Liu, F.; Hu, T.; Shen, W.; Li, E.; Liao, S.; Zou, Y. Mulberry Leaf Polyphenols Attenuated Postprandial Glucose Absorption: Via Inhibition of Disaccharidases Activity and Glucose Transport in Caco-2 Cells. *Food Funct.* **2020**, *11*, 1835–1844. [[CrossRef](#)] [[PubMed](#)]
102. Lingua, M.S.; Theumer, M.G.; Kruzynski, P.; Wunderlin, D.A.; Baroni, M.V. Bioaccessibility of Polyphenols and Antioxidant Properties of the White Grape by Simulated Digestion and Caco-2 Cell Assays: Comparative Study with Its Winemaking Product. *Food Res. Intal.* **2019**, *122*, 496–505. [[CrossRef](#)]
103. Zhang, H.; Hassan, Y.I.; Renaud, J.; Liu, R.; Yang, C.; Sun, Y.; Tsao, R. Bioaccessibility, Bioavailability, and Anti-Inflammatory Effects of Anthocyanins from Purple Root Vegetables Using Mono- and Co-Culture Cell Models. *Mol. Nutr. Food Res.* **2017**, *61*, 1600928. [[CrossRef](#)]
104. Grootaert, C.; Kamiloglu, S.; Capanoglu, E.; Van Camp, J. Cell Systems to Investigate the Impact of Polyphenols on Cardiovascular Health. *Nutrients* **2015**, *7*, 9229–9255. [[CrossRef](#)]
105. Olivero-David, R.; Ruiz-Roso, M.B.; Caporaso, N.; Perez-Olleros, L.; De las Heras, N.; Lahera, V.; Ruiz-Roso, B. In Vivo Bioavailability of Polyphenols from Grape By-Product Extracts, and Effect on Lipemia of Normocholesterolemic Wistar Rats. *J. Sci. Food Agric.* **2018**, *98*, 5581–5590. [[CrossRef](#)]
106. Peng, Y.; Meng, Q.; Zhou, J.; Chen, B.; Xi, J.; Long, P.; Zhang, L.; Hou, R. Nanoemulsion Delivery System of Tea Polyphenols Enhanced the Bioavailability of Catechins in Rats. *Food Chem.* **2018**, *242*, 527–532. [[CrossRef](#)]
107. Lambert, J.D.; Hong, J.; Kim, D.H.; Mishin, V.M.; Yang, C.S. Piperine Enhances the Bioavailability of the Tea Polyphenol (–)-Epigallocatechin-3-Gallate in Mice. *J. Nutr.* **2004**, *134*, 1948–1952. [[CrossRef](#)]
108. Curti, V.; Zaccaria, V.; Sokeng, A.J.T.; Dacrema, M.; Masiello, I.; Mascaro, A.; D’antona, G.; Daglia, M. Bioavailability and in Vivo Antioxidant Activity of a Standardized Polyphenol Mixture Extracted from Brown Propolis. *Int. J. Mol. Sci.* **2019**, *20*, 1250. [[CrossRef](#)]
109. Scholz, S.; Williamson, G. Interactions Affecting the Bioavailability of Dietary Polyphenols in Vivo. *Int. J. Vitam. Nutr. Res.* **2007**, *77*, 224–235. [[CrossRef](#)]
110. Zhong, S.; Sandhu, A.; Edirisinghe, I.; Burton-Freeman, B. Characterization of Wild Blueberry Polyphenols Bioavailability and Kinetic Profile in Plasma over 24-h Period in Human Subjects. *Mol. Nutr. Food Res.* **2017**, *61*, 1700405. [[CrossRef](#)]
111. Vitaglione, P.; Barone Lumaga, R.; Ferracane, R.; Sellitto, S.; Morelló, J.R.; Reguant Miranda, J.; Shimoni, E.; Fogliano, V. Human Bioavailability of Flavanols and Phenolic Acids from Cocoa-Nut Creams Enriched with Free or Microencapsulated Cocoa Polyphenols. *Br. J. Nutr.* **2013**, *109*, 1832–1843. [[CrossRef](#)]
112. Martínez-Huélamo, M.; Vallverdú-Queralt, A.; Di Lecce, G.; Valderas-Martínez, P.; Tulipani, S.; Jáuregui, O.; Escribano-Ferrer, E.; Estruch, R.; Illan, M.; Lamuela-Raventós, R.M. Bioavailability of Tomato Polyphenols Is Enhanced by Processing and Fat Addition: Evidence from a Randomized Feeding Trial. *Mol. Nutr. Food Res.* **2016**, *60*, 1578–1589. [[CrossRef](#)] [[PubMed](#)]
113. Marín, L.; Miguélez, E.M.; Villar, C.J.; Lombó, F. Bioavailability of Dietary Polyphenols and Gut Microbiota Metabolism: Antimicrobial Properties. *Biomed. Res. Int.* **2015**, *2015*, 905215. [[CrossRef](#)] [[PubMed](#)]

114. Singh, A.K.; Cabral, C.; Kumar, R.; Ganguly, R.; Rana, H.K.; Gupta, A.; Lauro, M.R.; Carbone, C.; Reis, F.; Pandey, A.K. Beneficial Effects of Dietary Polyphenols on Gut Microbiota and Strategies to Improve Delivery Efficiency. *Nutrients* **2019**, *11*, 2216. [[CrossRef](#)] [[PubMed](#)]
115. Kawabata, K.; Yoshioka, Y.; Terao, J. Role of Intestinal Microbiota in the Bioavailability and Physiological Functions of Dietary Polyphenols. *Molecules* **2019**, *24*, 370. [[CrossRef](#)] [[PubMed](#)]
116. Zhu, Y.; Zhang, J.-Y.; Wei, Y.-L.; Hao, J.-Y.; Lei, Y.-Q.; Zhao, W.-B.; Xiao, Y.-H.; Sun, A.-D. The Polyphenol-Rich Extract from Chokeberry (*Aronia Melanocarpa* L.) Modulates Gut Microbiota and Improves Lipid Metabolism in Diet-Induced Obese Rats. *Nutr. Metab.* **2020**, *17*, 54. [[CrossRef](#)]
117. Istas, G.; Wood, E.; Le Sayec, M.; Rawlings, C.; Yoon, J.; Dandavate, V.; Cera, D.; Rampelli, S.; Costabile, A.; Fromentin, E.; et al. Effects of Aronia Berry (Poly)Phenols on Vascular Function and Gut Microbiota: A Double-Blind Randomized Controlled Trial in Adult Men. *Am. J. Clin. Nutr.* **2019**, *110*, 316–329. [[CrossRef](#)]
118. Winter, J.; Popoff, M.R.; Grimont, P.; Bokkenheuser, V.D. *Clostridium Orbiscindens* Sp. Nov., a Human Intestinal Bacterium Capable of Cleaving the Flavonoid C-Ring. *Int. J. Syst. Bacteriol.* **1991**, *41*, 355–357. [[CrossRef](#)]
119. Simons, A.L.; Renouf, M.; Hendrich, S.; Murphy, P.A. Human Gut Microbial Degradation of Flavonoids: Structure–Function Relationships. *J. Agric. Food Chem.* **2005**, *53*, 4258–4263. [[CrossRef](#)]
120. Liu, Z.; Hu, M. Natural Polyphenol Disposition via Coupled Metabolic Pathways. *Expert Opin. Drug Metab. Toxicol.* **2007**, *3*, 389–406. [[CrossRef](#)]
121. Van Duynhoven, J.; Vaughan, E.E.; Jacobs, D.M.; Kemperman, R.A.; Van Velzen, E.J.J.; Gross, G.; Roger, L.C.; Possemiers, S.; Smilde, A.K.; Doré, J.; et al. Metabolic Fate of Polyphenols in the Human Superorganism. *Proc. Natl. Acad. Sci. USA* **2011**, *108* (Suppl. 1), 4531–4538. [[CrossRef](#)]
122. Luca, S.V.; Macovei, I.; Bujor, A.; Miron, A.; Skalicka-Woźniak, K.; Aprotosoia, A.C.; Trifan, A. Bioactivity of Dietary Polyphenols: The Role of Metabolites. *Crit. Rev. Food Sci. Nutr.* **2020**, *60*, 626–659. [[CrossRef](#)]
123. Tu, Y.; Wang, L.; Rong, Y.; Tam, V.; Yin, T.; Gao, S.; Singh, R.; Hu, M. Hepatoenteric Recycling Is a New Disposition Mechanism for Orally Administered Phenolic Drugs and Phytochemicals in Rats. *Elife* **2021**, *10*, e58820. [[CrossRef](#)]
124. Andres-Lacueva, C.; Macarulla, M.T.; Rotches-Ribalta, M.; Boto-Ordóñez, M.; Urpi-Sarda, M.; Rodríguez, V.M.; Portillo, M.P. Distribution of Resveratrol Metabolites in Liver, Adipose Tissue, and Skeletal Muscle in Rats Fed Different Doses of This Polyphenol. *J. Agric. Food Chem.* **2012**, *60*, 4833–4840. [[CrossRef](#)]
125. Serra, A.; Macià, A.; Romero, M.-P.; Anglès, N.; Morelló, J.R.; Motilva, M.-J. Distribution of Procyanidins and Their Metabolites in Rat Plasma and Tissues after an Acute Intake of Hazelnut Extract. *Food Funct.* **2011**, *2*, 562–568. [[CrossRef](#)]
126. Makarewicz, M.; Drożdż, I.; Tarko, T.; Duda-Chodak, A. The Interactions between Polyphenols and Microorganisms, Especially Gut Microbiota. *Antioxidants* **2021**, *10*, 188. [[CrossRef](#)] [[PubMed](#)]
127. Enaru, B.; Socaci, S.; Farcas, A.; Socaciu, C.; Danciu, C.; Stanila, A.; Diaconeasa, Z. Novel Delivery Systems of Polyphenols and Their Potential Health Benefits. *Pharmaceuticals* **2021**, *14*, 946. [[CrossRef](#)] [[PubMed](#)]
128. Pimentel-Moral, S.; Teixeira, M.C.; Fernandes, A.R.; Arráez-Román, D.; Martínez-Férez, A.; Segura-Carretero, A.; Souto, E.B. Lipid Nanocarriers for the Loading of Polyphenols—A Comprehensive Review. *Adv. Colloid Interface* **2018**, *260*, 85–94. [[CrossRef](#)] [[PubMed](#)]
129. Miksits, M.; Maier-Salamon, A.; Aust, S.; Thalhammer, T.; Reznicek, G.; Kunert, O.; Haslinger, E.; Szekeres, T.; Jaeger, W. Sulfation of Resveratrol in Human Liver: Evidence of a Major Role for the Sulfotransferases SULT1A1 and SULT1E1. *Xenobiotica* **2005**, *35*, 1101–1119. [[CrossRef](#)]
130. Hambidge, K.M. Micronutrient Bioavailability: Dietary Reference Intakes and a Future Perspective. *Am. J. Clin. Nutr.* **2010**, *91*, 1430–1432. [[CrossRef](#)]
131. Scalbert, A.; Williamson, G. Dietary Intake and Bioavailability of Polyphenols. *J. Nutr.* **2000**, *130* (Suppl. 8), 2073S–2085S. [[CrossRef](#)]
132. Kühnau, J. The Flavonoids. A Class of Semi-Essential Food Components: Their Role in Human Nutrition. *World Rev. Nutr. Diet* **1976**, *24*, 117–191.
133. Hertog, M.G.L.; Hollman, P.C.H.; Hertog, M.G.L.; Katan, M.B. Content of Potentially Anticarcinogenic Flavonoids of 28 Vegetables and 9 Fruits Commonly Consumed in the Netherlands. *J. Agric. Food Chem.* **1992**, *40*, 2379–2383. [[CrossRef](#)]
134. Reinli, K.; Block, G. Phytoestrogen Content of Foods—A Compendium of Literature Values. *Nutr. Cancer* **1996**, *26*, 123–148. [[CrossRef](#)]
135. Kimira, M.; Arai, Y.; Shimoi, K.; Watanabe, S. Japanese Intake of Flavonoids and Isoflavonoids from Foods. *J. Epidemiol.* **1998**, *8*, 168–175. [[CrossRef](#)]
136. Wakai, K.; Egami, I.; Kato, K.; Kawamura, T.; Tamakoshi, A.; Lin, Y.; Nakayama, T.; Wada, M.; Ohno, Y. Dietary Intake and Sources of Isoflavones among Japanese. *Nutr. Cancer* **1999**, *33*, 139–145. [[CrossRef](#)]
137. Kirk, P.; Patterson, R.E.; Lampe, J. Development of a Soy Food Frequency Questionnaire to Estimate Isoflavone Consumption in US Adults. *J. Am. Diet Assoc.* **1999**, *99*, 558–563. [[CrossRef](#)]
138. Gutiérrez-Grijalva, E.P.; Ambriz-Pere, D.L.; Leyva-Lopez, N.; Castillo-Lopez, R.I.; Heiedia, J.B. Review: Dietary Phenolic Compounds, Health Benefits and Bioaccessibility. *Arch. Latinoam. Nutr.* **2016**, *66*, 87–100.

139. Del Bo, C.; Bernardi, S.; Marino, M.; Porrini, M.; Tucci, M.; Guglielmetti, S.; Cherubini, A.; Carrieri, B.; Kirkup, B.; Kroon, P.; et al. Systematic Review on Polyphenol Intake and Health Outcomes: Is There Sufficient Evidence to Define a Health-Promoting Polyphenol-Rich Dietary Pattern? *Nutrients* **2019**, *11*, 1355. [[CrossRef](#)]
140. Wang, J.; Ballon, A.; Schroën, K.; de Lamo-Castellví, S.; Ferrando, M.; Güell, C. Polyphenol Loaded W1/O/W2 Emulsions Stabilized with Lesser Mealworm (*Alphitobius Diaperinus*) Protein Concentrate Produced by Membrane Emulsification: Stability under Simulated Storage, Process, and Digestion Conditions. *Foods* **2021**, *10*, 2997. [[CrossRef](#)]
141. Zembyla, M.; Murray, B.S.; Sarkar, A. Water-In-Oil Pickering Emulsions Stabilized by Water-Insoluble Polyphenol Crystals. *Langmuir* **2018**, *34*, 10001–10011. [[CrossRef](#)]
142. Martins, C.; Higaki, N.T.F.; Montrucchio, D.P.; de Oliveira, C.F.; Gomes, M.L.S.; Miguel, M.D.; Miguel, O.G.; Zanin, S.M.W.; Dias, J.d.F.G. Development of W1/O/W2 Emulsion with Gallic Acid in the Internal Aqueous Phase. *Food Chem.* **2020**, *314*, 126174. [[CrossRef](#)] [[PubMed](#)]
143. Pralhad, T.; Rajendrakumar, K. Study of Freeze-Dried Quercetin–Cyclodextrin Binary Systems by DSC, FT-IR, X-Ray Diffraction and SEM Analysis. *J. Pharm. Biomed. Anal.* **2004**, *34*, 333–339. [[CrossRef](#)] [[PubMed](#)]
144. Hunt, L.E.; Bourne, S.A.; Caira, M.R. Inclusion of Hydroxycinnamic Acids in Methylated Cyclodextrins: Host-Guest Interactions and Effects on Guest Thermal Stability. *Biomolecules* **2021**, *11*, 45. [[CrossRef](#)] [[PubMed](#)]
145. Matencio, A.; Hernández-García, S.; García-Carmona, F.; López-Nicolás, J.M. An Integral Study of Cyclodextrins as Solubility Enhancers of α -Methylstilbene, a Resveratrol Analogue. *Food Funct.* **2017**, *8*, 270–277. [[CrossRef](#)] [[PubMed](#)]
146. Nguyen Thi, D.P.; Tran, D.L.; Le Thi, P.; Park, K.D.; Hoang Thi, T.T. Supramolecular Gels Incorporating Cordyline Terminalis Leaf Extract as a Polyphenol Release Scaffold for Biomedical Applications. *Int. J. Mol. Sci.* **2021**, *22*, 8759. [[CrossRef](#)]
147. Boyacı, D.; Kavur, P.B.; Gulec, S.; Yemenicioğlu, A. Physicochemical and Active Properties of Gelatine-Based Composite Gels Loaded with Lysozyme and Green Tea Polyphenols. *Food Technol. Biotechnol.* **2021**, *59*, 337–348. [[CrossRef](#)]
148. Stanciauskaite, M.; Marksa, M.; Ivanauskas, L.; Perminaitė, K.; Ramanauskienė, K. Ophthalmic in Situ Gels with Balsam Poplar Buds Extract: Formulation, Rheological Characterization, and Quality Evaluation. *Pharmaceutics* **2021**, *13*, 953. [[CrossRef](#)]
149. Niknam, S.M.; Kashaninejad, M.; Escudero, I.; Sanz, M.T.; Beltrán, S.; Benito, J.M. Preparation of Water-in-Oil Nanoemulsions Loaded with Phenolic-Rich Olive Cake Extract Using Response Surface Methodology Approach. *Foods* **2022**, *11*, 279. [[CrossRef](#)]
150. Ragelle, H.; Crauste-Manciet, S.; Seguin, J.; Brossard, D.; Scherman, D.; Arnaud, P.; Chabot, G.G. Nanoemulsion Formulation of Fisetin Improves Bioavailability and Antitumour Activity in Mice. *Int. J. Pharm.* **2012**, *427*, 452–459. [[CrossRef](#)]
151. Rabelo, C.A.S.; Taarji, N.; Khalid, N.; Kobayashi, I.; Nakajima, M.; Neves, M.A. Formulation and Characterization of Water-in-Oil Nanoemulsions Loaded with Açai Berry Anthocyanins: Insights of Degradation Kinetics and Stability Evaluation of Anthocyanins and Nanoemulsions. *Food Res. Int.* **2018**, *106*, 542–548. [[CrossRef](#)] [[PubMed](#)]
152. Seguin, J.; Brullé, L.; Boyer, R.; Lu, Y.M.; Romano Ramos, M.; Touil, Y.S.; Scherman, D.; Bessodes, M.; Mignet, N.; Chabot, G.G. Liposomal Encapsulation of the Natural Flavonoid Fisetin Improves Bioavailability and Antitumor Efficacy. *Int. J. Pharm.* **2013**, *444*, 146–154. [[CrossRef](#)] [[PubMed](#)]
153. Sultana, F.; Neog, M.K.; Rasool, M.K. Targeted Delivery of Morin, a Dietary Bioflavanol Encapsulated Mannosylated Liposomes to the Macrophages of Adjuvant-Induced Arthritis Rats Inhibits Inflammatory Immune Response and Osteoclastogenesis. *Eur. J. Pharm. Biopharm.* **2017**, *115*, 229–242. [[CrossRef](#)] [[PubMed](#)]
154. Wei, X.Q.; Zhu, J.F.; Wang, X.B.; Ba, K. Improving the Stability of Liposomal Curcumin by Adjusting the Inner Aqueous Chamber PH of Liposomes. *ACS Omega* **2020**, *5*, 1120–1126. [[CrossRef](#)] [[PubMed](#)]
155. Vega, E.N.; Molina, A.K.; Pereira, C.; Dias, M.I.; Heleno, S.A.; Rodrigues, P.; Fernandes, I.P.; Barreiro, M.F.; Stojković, D.; Soković, M.; et al. Anthocyanins from *Rubus Fruticosus* L. and *Morus Nigra* L. Applied as Food Colorants: A Natural Alternative. *Plants* **2021**, *10*, 1181. [[CrossRef](#)] [[PubMed](#)]
156. Yang, B.; Dong, Y.; Wang, F.; Zhang, Y. Nanoformulations to Enhance the Bioavailability and Physiological Functions of Polyphenols. *Molecules* **2020**, *25*, 4613. [[CrossRef](#)] [[PubMed](#)]
157. Shaygannia, S.; Eshaghi, M.R.; Fazel, M.; Hashemiravan, M. The Effect of Microencapsulation of Phenolic Compounds from Lemon Waste by Persian and Basil Seed Gums on the Chemical and Microbiological Properties of Mayonnaise. *Prev. Nutr. Food Sci.* **2021**, *26*, 82–91. [[CrossRef](#)] [[PubMed](#)]
158. Desai, N.M.; Haware, D.J.; Basavaraj, K.; Murthy, P.S. Microencapsulation of Antioxidant Phenolic Compounds from Green Coffee. *Prep. Biochem. Biotechnol.* **2019**, *49*, 400–406. [[CrossRef](#)]
159. de Meneses Costa Ferreira, L.M.; Pereira, R.R.; de Carvalho-Guimarães, F.B.; do Nascimento Remígio, M.S.; Barbosa, W.L.R.; Ribeiro-Costa, R.M.; Silva-Júnior, J.O.C. Microencapsulation by Spray Drying and Antioxidant Activity of Phenolic Compounds from Tucuma Coproduct. *Polymers* **2022**, *14*, 2905. [[CrossRef](#)]
160. Rosales, T.K.O.; Hassimotto, N.M.A.; Lajolo, F.M.; Fabi, J.P. Nanotechnology as a Tool to Mitigate the Effects of Intestinal Microbiota on Metabolization of Anthocyanins. *Antioxidants* **2022**, *11*, 506. [[CrossRef](#)]
161. Saifullah, M.; Shishir, M.R.I.; Ferdowsi, R.; Tanver Rahman, M.R.; Van Vuong, Q. Micro and Nano Encapsulation, Retention and Controlled Release of Flavor and Aroma Compounds: A Critical Review. *Trends Food Sci. Technol.* **2019**, *86*, 230–251. [[CrossRef](#)]
162. Behzadi, S.; Serpooshan, V.; Tao, W.; Hamaly, M.A.; Alkawareek, M.Y.; Dreaden, E.C.; Brown, D.; Alkilany, A.M.; Farokhzad, O.C.; Mahmoudi, M. Cellular Uptake of Nanoparticles: Journey Inside the Cell. *Chem. Soc. Rev.* **2017**, *46*, 4218–4244. [[CrossRef](#)]
163. Mosquera, J.; García, I.; Liz-Marzán, L.M. Cellular Uptake of Nanoparticles versus Small Molecules: A Matter of Size. *Acc. Chem. Res.* **2018**, *51*, 2305–2313. [[CrossRef](#)]

164. Rajesh, S.; Lillard, J.W., Jr. Nanoparticle-Based Targeted Drug Delivery. *Exp. Mol. Pathol.* **2009**, *86*, 215–223. [[CrossRef](#)]
165. Lamprecht, A.; Ubrich, N.; Yamamoto, H.; Schäfer, U.; Takeuchi, H.; Maincent, P.; Kawashima, Y.; Lehr, C.M. Biodegradable Nanoparticles for Targeted Drug Delivery in Treatment of Inflammatory Bowel Disease. *J. Pharmacol. Exp. Ther.* **2001**, *299*, 775–781.
166. Majumder, J.; Taratula, O.; Minko, T. Nanocarrier-Based Systems for Targeted and Site Specific Therapeutic Delivery. *Adv. Drug Deliv. Rev.* **2019**, *144*, 57–77. [[CrossRef](#)]
167. Street, D.; Bangsbo, J.; Juel, C. Interstitial PH in Human Skeletal Muscle during and after Dynamic Graded Exercise. *J. Physiol.* **2001**, *537*, 993–998. [[CrossRef](#)]
168. Parisi, O.I.; Puoci, F.; Restuccia, D.; Farina, G.; Iemma, F.; Picci, N. Polyphenols in Human Health and Disease. In *Polyphenols in Human Health and Disease*; Watson, R.R., Preedy, V.R., Zibadi, S., Eds.; Academic Press-Elsevier Inc.: Amsterdam, The Netherlands, 2014. [[CrossRef](#)]
169. Nakayama, T.; Hashimoto, T.; Kajiya, K.; Kumazawa, S. Affinity of Polyphenols for Lipid Bilayers. *Biofactors* **2000**, *13*, 147–151. [[CrossRef](#)]
170. Phan, H.T.T.; Yoda, T.; Chahal, B.; Morita, M.; Takagi, M.; Vestergaard, M.C. Structure-Dependent Interactions of Polyphenols with a Biomimetic Membrane System. *Biochim. Biophys. Acta* **2014**, *1838*, 2670–2677. [[CrossRef](#)]
171. Pruchnik, H.; Bonarska-Kujawa, D.; Żyłka, R.; Oszmiański, J.; Kleszczyńska, H. Application of the DSC and Spectroscopy Methods in the Analysis of the Protective Effect of Extracts from the Blueberry Fruit of the Genus *Vaccinium* in Relation to the Lipid Membrane. *J. Therm. Anal. Calorim.* **2018**, *134*, 679–689. [[CrossRef](#)]
172. Malekar, S.A.; Sarode, A.L.; Bach, A.C.; Worthen, D.R. The Localization of Phenolic Compounds in Liposomal Bilayers and Their Effects on Surface Characteristics and Colloidal Stability. *AAPS PharmSciTech* **2016**, *17*, 1468–1476. [[CrossRef](#)]
173. Powers, S.K.; Talbert, E.E.; Adihetty, P.J. Reactive Oxygen and Nitrogen Species as Intracellular Signals in Skeletal Muscle. *J. Physiol.* **2011**, *589*, 2129–2138. [[CrossRef](#)] [[PubMed](#)]
174. Boveris, A.; Chance, B. The Mitochondrial Generation of Hydrogen Peroxide. General Properties and Effect of Hyperbaric Oxygen. *Biochem. J.* **1973**, *134*, 707–716. [[CrossRef](#)] [[PubMed](#)]
175. Loschen, G.; Azzi, A.; Richter, C.; Flohe, L. Superoxide Radicals as Precursors of Mitochondrial Hydrogen Peroxide. *FEBS Lett.* **1974**, *42*, 68–72. [[CrossRef](#)] [[PubMed](#)]
176. St-Pierre, J.; Buckingham, J.A.; Roebuck, S.J.; Brand, M.D. Topology of Superoxide Production from Different Sites in the Mitochondrial Electron Transport Chain. *J. Biol. Chem.* **2002**, *277*, 44784–44790. [[CrossRef](#)] [[PubMed](#)]
177. Anderson, E.J.; Neuffer, P.D. Type II Skeletal Myofibers Possess Unique Properties That Potentiate Mitochondrial H₂O₂ Generation. *Am. J. Physiol. Cell Physiol.* **2006**, *290*, 844–851. [[CrossRef](#)]
178. Walsh, B.; Tonkonogi, M.; Sahlin, K. Effect of Endurance Training on Oxidative and Antioxidative Function in Human Permeabilized Muscle Fibres. *Pflugers Arch.* **2001**, *442*, 420–425. [[CrossRef](#)]
179. He, F.; Li, J.; Liu, Z.; Chuang, C.C.; Yang, W.; Zuo, L. Redox Mechanism of Reactive Oxygen Species in Exercise. *Front. Physiol.* **2016**, *7*, 486. [[CrossRef](#)]
180. Larsen, F.J.; Schiffer, T.A.; Ørtenblad, N.; Zinner, C.; Morales-Alamo, D.; Willis, S.J.; Calbet, J.A.; Holmberg, H.C.; Boushel, R. High-Intensity Sprint Training Inhibits Mitochondrial Respiration through Aconitase Inactivation. *FASEB J.* **2016**, *30*, 417–427. [[CrossRef](#)]
181. Vezzoli, A.; Pugliese, L.; Marzorati, M.; Serpiello, F.R.; La Torre, A.; Porcelli, S. Time-Course Changes of Oxidative Stress Response to High-Intensity Discontinuous Training versus Moderate-Intensity Continuous Training in Masters Runners. *PLoS ONE* **2014**, *9*, e87506. [[CrossRef](#)]
182. Mrakic-Spota, S.; Gussoni, M.; Porcelli, S.; Pugliese, L.; Pavei, G.; Bellistri, G.; Montorsi, M.; Tacchini, P.; Vezzoli, A. Training Effects on ROS Production Determined by Electron Paramagnetic Resonance in Master Swimmers. *Oxid. Med. Cell Longev.* **2015**, *2015*, 804794. [[CrossRef](#)]
183. Radak, Z.; Chung, H.Y.; Koltai, E.; Taylor, A.W.; Goto, S. Exercise, Oxidative Stress and Hormesis. *Ageing Res. Rev.* **2008**, *7*, 34–42. [[CrossRef](#)]
184. Sakellariou, G.K.; Vasilaki, A.; Palomero, J.; Kayani, A.; Zibrik, L.; McArdle, A.; Jackson, M.J. Studies of Mitochondrial and Nonmitochondrial Sources Implicate Nicotinamide Adenine Dinucleotide Phosphate Oxidase(s) in the Increased Skeletal Muscle Superoxide Generation That Occurs during Contractile Activity. *Antioxid. Redox Signal* **2013**, *18*, 603–621. [[CrossRef](#)]
185. Sahlin, K.; Shabalina, I.G.; Mattsson, C.M.; Bakkman, L.; Fernström, M.; Rozhdestvenskaya, Z.; Enqvist, J.K.; Nedergaard, J.; Ekblom, B.; Tonkonogi, M. Ultraendurance Exercise Increases the Production of Reactive Oxygen Species in Isolated Mitochondria from Human Skeletal Muscle. *J. Appl. Physiol.* **2010**, *108*, 780–787. [[CrossRef](#)]
186. Zoladz, J.A.; Koziel, A.; Woyda-PLoSzczyca, A.; Celichowski, J.; Jarmuszkiewicz, W. Endurance Training Increases the Efficiency of Rat Skeletal Muscle Mitochondria. *Pflugers Arch.* **2016**, *468*, 1709–1724. [[CrossRef](#)]
187. Di Meo, S.; Napolitano, G.; Venditti, P. Mediators of Physical Activity Protection against Ros-Linked Skeletal Muscle Damage. *Int. J. Mol. Sci.* **2019**, *20*, 3024. [[CrossRef](#)]
188. Nakama, L.H.; King, K.B.; Abrahamsson, S.; Rempel, D.M. Evidence of Tendon Microtears Due to Cyclical Loading in an in Vivo Tendinopathy Model. *J. Orthop. Res.* **2005**, *23*, 1199–1205. [[CrossRef](#)]
189. Friden, J.; Sjøstrom, M.; Ekblom, B. Myofibrillar Damage Following Intense Eccentric Exercise in Man. *Int. J. Sport Med.* **1983**, *4*, 170–176. [[CrossRef](#)]

190. Warren, G.L.; Hulderman, T.; Jensen, N.; McKinstry, M.; Mishra, M.; Luster, M.I.; Simeonova, P.P. Physiological Role of Tumor Necrosis Factor Alpha in Traumatic Muscle Injury. *FASEB J.* **2002**, *16*, 1630–1632. [[CrossRef](#)]
191. Saclier, M.; Cuvellier, S.; Magnan, M.; Mounier, R.; Chazaud, B. Monocyte/Macrophage Interactions with Myogenic Precursor Cells during Skeletal Muscle Regeneration. *FEBS J.* **2013**, *280*, 4118–4130. [[CrossRef](#)]
192. Collins, R.A.; Grounds, M.D. The Role of Tumor Necrosis Factor-Alpha (TNF- α) in Skeletal Muscle Regeneration: Studies in TNF- α (-/-) and TNF- α (-/-)/LT- α (-/-) Mice. *J. Histochem. Cytochem.* **2001**, *49*, 989–1001. [[CrossRef](#)]
193. Martinez, F.O.; Gordon, S. The M1 and M2 Paradigm of Macrophage Activation: Time for Reassessment. *F1000Prime Rep.* **2014**, *6*, 13. [[CrossRef](#)] [[PubMed](#)]
194. Castaneda, O.A.; Lee, S.C.; Ho, C.T.; Huang, T.C. Macrophages in Oxidative Stress and Models to Evaluate the Antioxidant Function of Dietary Natural Compounds. *J. Food Drug. Anal.* **2017**, *25*, 111–118. [[CrossRef](#)] [[PubMed](#)]
195. Samavati, L.; Lee, I.; Mathes, I.; Lottspeich, F.; Hüttemann, M. Tumor Necrosis Factor α Inhibits Oxidative Phosphorylation through Tyrosine Phosphorylation at Subunit I of Cytochrome c Oxidase. *J. Biol. Chem.* **2008**, *283*, 21134–21144. [[CrossRef](#)] [[PubMed](#)]
196. Langen, R.C.J.; Schols, A.M.W.J.; Kelders, M.C.J.M.; Van Der Velden, J.L.J.; Wouters, E.F.M.; Janssen-Heininger, Y.M.W. Tumor Necrosis Factor-Alpha Inhibits Myogenesis through Redox-Dependent and -Independent Pathways. *Am. J. Physiol. Cell Physiol.* **2002**, *283*, 714–721. [[CrossRef](#)] [[PubMed](#)]
197. Czarkowska-Paczek, B.; Bartłomiejczyk, I.; Przybylski, J. The Serum Levels of Growth Factors: PDGF, TGF-Beta and VEGF Are Increased after Strenuous Physical Exercise. *J. Physiol. Pharmacol.* **2006**, *57*, 189–197.
198. Kreuzer, J.; Viedt, C.; Brandes, R.P.; Seeger, F.; Rosenkranz, A.S.; Sauer, H.; Babich, A.; Nürnberg, B.; Kather, H.; Krieger-Brauer, H.I. Platelet-Derived Growth Factor Activates Production of Reactive Oxygen Species by NAD(P)H Oxidase in Smooth Muscle Cells through Gi1,2. *FASEB J.* **2003**, *17*, 38–40. [[CrossRef](#)]
199. Fischer, C.P. Interleukin-6 in Acute Exercise and Training: What Is the Biological Relevance? *Exerc. Immunol. Rev.* **2006**, *12*, 6–33.
200. Forcina, L.; Miano, C.; Scicchitano, B.M.; Rizzuto, E.; Berardinelli, M.G.; De Benedetti, F.; Pelosi, L.; Musarò, A. Increased Circulating Levels of Interleukin-6 Affect the Redox Balance in Skeletal Muscle. *Oxid. Med. Cell Longev.* **2019**, *2019*, 301858. [[CrossRef](#)]
201. Kistner, T.M.; Pedersen, B.K.; Lieberman, D.E. Interleukin 6 as an Energy Allocator in Muscle Tissue. *Nat. Metab.* **2022**, *4*, 170–179. [[CrossRef](#)]
202. Abid, H.; Ryan, Z.C.; Delmotte, P.; Sieck, G.C.; Lanza, I.R. Extramyocellular Interleukin-6 Influences Skeletal Muscle Mitochondrial Physiology through Canonical JAK/STAT Signaling Pathways. *FASEB J.* **2020**, *34*, 14458–14472. [[CrossRef](#)]
203. Cheng, A.J.; Jude, B.; Lanner, J.T. Intramuscular Mechanisms of Overtraining. *Redox Biol.* **2020**, *35*, 101480. [[CrossRef](#)]
204. Ostrowski, K.; Rohde, T.; Asp, S.; Schjerling, P.; Pedersen, B.K. Pro- and Anti-Inflammatory Cytokine Balance in Strenuous Exercise in Humans. *J. Physiol.* **1999**, *515*, 287–291. [[CrossRef](#)]
205. Peake, J.M.; Suzuki, K.; Hordern, M.; Wilson, G.; Nosaka, K.; Coombes, J.S. Plasma Cytokine Changes in Relation to Exercise Intensity and Muscle Damage. *Eur. J. Appl. Physiol.* **2005**, *95*, 514–521. [[CrossRef](#)]
206. Arnold, L.; Henry, A.; Poron, F.; Baba-Amer, Y.; Van Rooijen, N.; Plonquet, A.; Gherardi, R.K.; Chazaud, B. Inflammatory Monocytes Recruited after Skeletal Muscle Injury Switch into Antiinflammatory Macrophages to Support Myogenesis. *J. Exp. Med.* **2007**, *204*, 1057–1069. [[CrossRef](#)] [[PubMed](#)]
207. Thirupathi, A.; Gu, Y.; Pinho, R.A. Exercise Cuts Both Ways with ROS in Remodifying Innate and Adaptive Responses: Rewiring the Redox Mechanism of the Immune System during Exercise. *Antioxidants* **2021**, *10*, 1846. [[CrossRef](#)]
208. Kozakowska, M.; Pietraszek-Gremplewicz, K.; Jozkowicz, A.; Dulak, J. The Role of Oxidative Stress in Skeletal Muscle Injury and Regeneration: Focus on Antioxidant Enzymes. *J. Muscle Res. Cell Motil.* **2015**, *36*, 377–393. [[CrossRef](#)]
209. Powers, S.K.; Duarte, J.; Kavazis, A.N.; Talbert, E.E. Reactive Oxygen Species Are Signalling Molecules for Skeletal Muscle Adaptation. *Exp. Physiol.* **2010**, *95*, 1–9. [[CrossRef](#)]
210. Morgan, M.J.; Liu, Z.G. Crosstalk of Reactive Oxygen Species and NF-KB Signaling. *Cell Res.* **2011**, *21*, 103–115. [[CrossRef](#)]
211. Gallego-Selles, A.; Galvan-Alvarez, V.; Martinez-Canton, M.; Garcia-Gonzalez, E.; Morales-Alamo, D.; Santana, A.; Gonzalez-Henriquez, J.J.; Dorado, C.; Calbet, J.A.L.; Martin-Rincon, M. Redox Biology Fast Regulation of the NF- κ B Signalling Pathway in Human Skeletal Muscle Revealed by High-Intensity Exercise and Ischaemia at Exhaustion: Role of Oxygenation and Metabolite Accumulation. *Redox Biol.* **2022**, *55*, 102398. [[CrossRef](#)]
212. Kabe, Y.; Ando, K.; Hirao, S.; Yoshida, M.; Handa, H. Redox Regulation of NF-KB Activation: Distinct Redox Regulation Between the Cytoplasm and the Nucleus. *Antioxid. Redox Signal* **2005**, *7*, 395–403. [[CrossRef](#)]
213. Nunes Silva, A. The Association between Physical Exercise and Reactive Oxygen Species (ROS) Production. *J. Sport Med. Doping Stud.* **2015**, *5*. [[CrossRef](#)]
214. Canton, M.; Sánchez-Rodríguez, R.; Spera, I.; Venegas, F.C.; Favia, M.; Viola, A.; Castegna, A. Reactive Oxygen Species in Macrophages: Sources and Targets. *Front. Immunol.* **2021**, *12*, 734229. [[CrossRef](#)]
215. Nikawa, T.; Ulla, A.; Sakakibara, I. Polyphenols and Their Effects on Muscle Atrophy and Muscle Health. *Molecules* **2021**, *26*, 4887. [[CrossRef](#)] [[PubMed](#)]
216. Escobar, M.; Oliveira, M.; Behr, G.; Zanotto-Filho, A.; Ilha, L.; Cunha, G.D.; De Oliveira, A.R.D.; Moreira, J.C.F. Oxidative Stress in Young Football (Soccer) Players in Intermittent High Intensity Exercise Protocol. *JEPonline* **2009**, *12*, 1–10.

217. Vezzoli, A.; Dellanoce, C.; Mrakic-Spota, S.; Montorsi, M.; Moretti, S.; Tonini, A.; Pratali, L.; Accinni, R. Oxidative Stress Assessment in Response to Ultraendurance Exercise: Thiols Redox Status and ROS Production According to Duration of a Competitive Race. *Oxid. Med. Cell Longev.* **2016**, *2016*, 6439037. [[CrossRef](#)]
218. Spanidis, Y.; Stagos, D.; Orfanou, M.; Goutzourelas, N.; Bar-Or, D.; Spandidos, D.; Kouretas, D. Variations in Oxidative Stress Levels in 3 Days Follow-up in Ultramarathon Mountain Race Athletes. *J. Strength Cond. Res.* **2017**, *31*, 582–594. [[CrossRef](#)]
219. Niess, A.M.; Hartmann, A.; Grünert-Fuchs, M.; Poch, B.; Speit, G. DNA Damage after Exhaustive Treadmill Running in Trained and Untrained Men. *Int. J. Sport Med.* **1996**, *17*, 397–403. [[CrossRef](#)]
220. Sharifi, G.; Najafabadi, A.B.; Ghashghaei, F.E. Oxidative Stress and Total Antioxidant Capacity in Handball Players. *Adv. Biomed. Res.* **2014**, *3*, 181. [[CrossRef](#)]
221. Muñoz Marin, D.; Barrientos, G.; Alves, J.; Grijota, F.J.; Robles, M.C.; May Nar, M. Oxidative Stress, Lipid Peroxidation Indexes and Antioxidant Vitamins in Long and Middle Distance Athletes during a Sport Season. *J. Sport Med. Phys. Fit.* **2018**, *58*, 1713–1719. [[CrossRef](#)]
222. Becatti, M.; Mannucci, A.; Barygina, V.; Mascherini, G.; Emmi, G.; Silvestri, E.; Wright, D.; Taddei, N.; Galanti, G.; Fiorillo, C. Redox Status Alterations during the Competitive Season in Elite Soccer Players: Focus on Peripheral Leukocyte-Derived ROS. *Intern. Emerg. Med.* **2017**, *12*, 777–788. [[CrossRef](#)]
223. Nikolaidis, M.G.; Kyparos, A.; Hadziioannou, M.; Panou, N.; Samaras, L.; Jamurtas, A.Z.; Kouretas, D. Acute Exercise Markedly Increases Blood Oxidative Stress in Boys and Girls. *Appl. Physiol. Nutr. Metab.* **2007**, *32*, 197–205. [[CrossRef](#)]
224. Alkazemi, D.; Rahman, A.; Habra, B. Alterations in Glutathione Redox Homeostasis among Adolescents with Obesity and Anemia. *Sci. Rep.* **2021**, *11*, 3034. [[CrossRef](#)]
225. Sastre, J.; Asensi, M.; Gasco, E.; Pallardo, F.V.; Ferrero, J.A.; Furukawa, T.; Vina, J. Exhaustive Physical Exercise Causes Oxidation of Glutathione Status in Blood: Prevention by Antioxidant Administration. *Am. J. Physiol.* **1992**, *263*, R992–R995. [[CrossRef](#)]
226. Schippinger, G.; Wonisch, W.; Abuja, P.M.; Fankhauser, F.; Winklhofer-Roob, B.M.; Halwachs, G. Lipid Peroxidation and Antioxidant Status in Professional. *Eur. J. Clin. Invest.* **2002**, *32*, 686–692. [[CrossRef](#)]
227. Le Moal, E.; Groussard, C.; Paillard, T.; Chaory, K.; Le Bris, R.; Plantet, K.; Pincemail, J.; Zouhal, H. Redox Status of Professional Soccer Players Is Influenced by Training Load Throughout a Season. *Inter. J. Sport Med.* **2016**, *37*, 680–686. [[CrossRef](#)]
228. Varamenti, E.; Tod, D.; Pullinger, S.A. Redox Homeostasis and Inflammation Responses to Training in Adolescent Athletes: A Systematic Review and Meta-Analysis. *Sport Med. Open* **2020**, *6*, 34. [[CrossRef](#)]
229. Lamprecht, M.; Greilberger, J.F.; Schwaberg, G.; Hofmann, P.; Oettl, K. Single Bouts of Exercise Affect Albumin Redox State and Carbonyl Groups on Plasma Protein of Trained Men in a Workload-Dependent Manner. *J. Appl. Physiol.* **2008**, *104*, 1611–1617. [[CrossRef](#)]
230. Zalavras, A.; Fatouros, I.G.; Deli, C.K.; Draganidis, D.; Theodorou, A.A.; Soulas, D.; Koutsioras, Y.; Koutedakis, Y.; Jamurtas, A.Z. Age-Related Responses in Circulating Markers of Redox Status in Healthy Adolescents and Adults during the Course of a Training Macrocycle. *Oxid. Med. Cell Longev.* **2015**, *2015*, 283921. [[CrossRef](#)]
231. Ammar, A.; Trabelsi, K.; Boukhris, O.; Glenn, J.M.; Bott, N.; Masmoudi, L.; Hakim, A.; Chtourou, H.; Driss, T.; Hoekelmann, A.; et al. Effects of Aerobic-, Anaerobic- and Combined-Based Exercises on Plasma Oxidative Stress Biomarkers in Healthy Untrained Young Adults. *Int. J. Environ. Res. Public Health* **2020**, *17*, 2601. [[CrossRef](#)]
232. Brites, F.D.; Evelson, P.A.; Christiansen, M.G.; Nicol, M.F.; Basílico, M.J.; Wikinski, R.W.; Llesuy, S.F. Soccer Players under Regular Training Show Oxidative Stress but an Improved Plasma Antioxidant Status. *Clin. Sci.* **1999**, *96*, 381–385. [[CrossRef](#)]
233. Dékány, M.; Nemeskéri, V.; Györe, I.; Harbula, I.; Malomsoki, J.; Pucso, J. Antioxidant Status of Interval-Trained Athletes in Various Sports. *Inter. J. Sport Med.* **2006**, *27*, 112–116. [[CrossRef](#)] [[PubMed](#)]
234. Melikoglu, M.A.; Kaldirimci, M.; Katkat, D.; Sen, I.; Kaplan, I.; Senel, K. The Effect of Regular Long Term Training on Antioxidant Enzymatic Activities. *J. Sport Med. Phys. Fit.* **2008**, *48*, 388–390.
235. Carlsohn, A.; Rohn, S.; Bittmann, F.; Raila, J.; Mayer, F.; Schweigert, F.J. Exercise Increases the Plasma Antioxidant Capacity of Adolescent Athletes. *Ann. Nutr. Metab.* **2008**, *53*, 96–103. [[CrossRef](#)]
236. Caplin, A.; Chen, F.S.; Beauchamp, M.R.; Puterman, E. The Effects of Exercise Intensity on the Cortisol Response to a Subsequent Acute Psychosocial Stressor. *Psychoneuroendocrinology* **2021**, *131*, 105336. [[CrossRef](#)] [[PubMed](#)]
237. Godoy, L.D.; Rossignoli, M.T.; Delfino-Pereira, P.; Garcia-Cairasco, N.; de Lima Umeoka, E.H. A Comprehensive Overview on Stress Neurobiology: Basic Concepts and Clinical Implications. *Front. Behav. Neurosci.* **2018**, *12*, 127. [[CrossRef](#)]
238. Ziembra, A.; Adamczyk, J.G.; Barczak, A.; Boguszewski, D.; Kozacz, A.; Dąbrowski, J.; Steczkowska, M.; Peplowska, B.; Żekanowski, C. Changes in the Hormonal Profile of Athletes Following a Combat Sports Performance. *Biomed. Res. Int.* **2020**, *2020*, 9684792. [[CrossRef](#)]
239. de Oliveira, L.F.G.; Souza-Junior, T.P.; Fechio, J.J.; Gomes-Santos, J.A.F.; Sampaio, R.C.; Vardaris, C.V.; Lambertucci, R.H.; de Barros, M.P. Uric Acid and Cortisol Levels in Plasma Correlate with Pre-Competition Anxiety in Novice Athletes of Combat Sports. *Brain Sci.* **2022**, *12*, 712. [[CrossRef](#)]
240. Messan, F.; Tito, A.; Gouthon, P.; Nouatin, K.B.; Nigan, I.B.; Blagbo, A.S.; Lounana, J.; Medelli, J. Comparison of Catecholamine Values before and after Exercise-Induced Bronchospasm in Professional Cyclists. *Tanaffos* **2017**, *16*, 136–143.
241. Zouhal, H.; Jacob, C.; Delamarche, P.; Gratas-Delamarche, A. Catecholamines and the Effects of Exercise, Training and Gender. *Sport. Med.* **2008**, *38*, 401–423. [[CrossRef](#)]

242. Singal, P.K.; Kapur, N.; Dhillon, K.S.; Beamish, R.E.; Dhalla, N.S. Role of Free Radicals in Catecholamine-Induced Cardiomyopathy. *Can. J. Physiol. Pharmacol.* **1982**, *60*, 1390–1397. [[CrossRef](#)]
243. Albino Teixeira, A.; Azevedo, I.; Branco, D.; Rodrigues-Pereira, E.; Osswald, W. Sympathetic Denervation Caused by Long-Term Noradrenaline Infusions; Prevention by Desipramine and Superoxide Dismutase. *Br. J. Pharmacol.* **1989**, *97*, 95–102. [[CrossRef](#)]
244. Kruk, J.; Kotarska, K.; Aboul-Enein, B.H. Physical Exercise and Catecholamines Response: Benefits and Health Risk: Possible Mechanisms. *Free Radic. Res.* **2020**, *54*, 105–125. [[CrossRef](#)]
245. Mehta, J.L.; Li, D. Epinephrine Upregulates Superoxide Dismutase in Human Coronary Artery Endothelial Cells. *Free Radic. Biol. Med.* **2001**, *30*, 148–153. [[CrossRef](#)]
246. Herman, J.P.; Mcklveen, J.M.; Ghosal, S.; Kopp, B.; Wulsin, A.; Makinson, R.; Scheimann, J.; Myers, B. Regulation of the Hypothalamic-Pituitary-Adrenocortical Stress Response. *Compr. Physiol.* **2016**, *6*, 603–621. [[CrossRef](#)]
247. Filaire, E.; Sagnol, M.; Ferrand, C.; Maso, F.; Lac, G. Psychophysiological Stress in Judo Athletes during Competitions. *J. Sport Med. Phys. Fit.* **2001**, *41*, 263–268.
248. Kraemer, W.J.; Fry, A.C.; Rubin, M.R.; Triplett-Mcbride, T.; Gordon, S.E.; Perry Koziris, L.; Lynch, J.M.; Volek, J.S.; Meuffels, D.E.; Newton, R.U.; et al. Physiological and Performance Responses to Tournament Wrestling. *Med. Sci. Sport Exerc.* **2001**, *33*, 1367–1378. [[CrossRef](#)]
249. Robazza, C.; Gallina, S.; D’Amico, M.A.; Izzicupo, P.; Bascelli, A.; Di Fonso, A.; Mazzafo, C.; Capobianco, A.; Di Baldassarre, A. Relationship between Biological Markers and Psychological States in Elite Basketball Players across a Competitive Season. *Psychol. Sport Exerc.* **2012**, *13*, 509–517. [[CrossRef](#)]
250. Doan, B.K.; Newton, R.U.; Kraemer, W.J.; Kwon, Y.H.; Scheet, T.P. Salivary Cortisol, Testosterone, and T/C Ratio Responses During a 36-Hole Golf Competition. *Int. J. Sport Med.* **2007**, *28*, 470–479. [[CrossRef](#)]
251. Lautenbach, F.; Laborde, S.; Klämpfl, M.; Achtzehn, S. A Link between Cortisol and Performance: An Exploratory Case Study of a Tennis Match. *Int. J. Psychophysiol.* **2015**, *98*, 167–173. [[CrossRef](#)]
252. Dovio, A.; Roveda, E.; Sciolla, C.; Montaruli, A.; Raffaelli, A.; Saba, A.; Calogiuri, G.; De Francia, S.; Borrione, P.; Salvadori, P.; et al. Intense Physical Exercise Increases Systemic 11 β -Hydroxysteroid Dehydrogenase Type 1 Activity in Healthy Adult Subjects. *Eur. J. Appl. Physiol.* **2010**, *108*, 681–687. [[CrossRef](#)]
253. Abate, M.; Salini, V. Oxidative Stress, Testosterone, Cortisol, and Vitamin D: Differences in Professional Soccer Players of African and Caucasian Origin. *Med. Princ. Pract.* **2022**, *31*, 352–358. [[CrossRef](#)] [[PubMed](#)]
254. Davison, G.; Gleeson, M.; Phillips, S. Antioxidant Supplementation and Immunoendocrine Responses to Prolonged Exercise. *Med. Sci. Sport Exerc.* **2007**, *39*, 645–652. [[CrossRef](#)] [[PubMed](#)]
255. Peters, E.M.; Anderson, R.; Theron, A.J. Attenuation of Increase in Circulating Cortisol and Enhancement of the Acute Phase Protein Response in Vitamin C-Supplemented Ultramarathoners. *Int. J. Sport Med.* **2001**, *22*, 120–126. [[CrossRef](#)] [[PubMed](#)]
256. Teixeira, V.; Valente, H.; Casal, S.; Pereira, L.; Marques, F.; Moreira, P. Antioxidant Status, Oxidative Stress, and Damage in Elite Kayakers after 1 Year of Training and Competition in 2 Seasons. *Int. J. Sport Nutr. Exerc. Metab.* **2009**, *34*, 716–724. [[CrossRef](#)] [[PubMed](#)]
257. Myburgh, K.H. Polyphenol Supplementation: Benefits for Exercise Performance or Oxidative Stress? *Sport Med.* **2014**, *44* (Suppl. 1), S57–S570. [[CrossRef](#)]
258. Moore, K.; Roberts, L.J., 2nd. Measurement of Lipid Peroxidation. *Free Radic. Res.* **1998**, *28*, 659–671. [[CrossRef](#)]
259. Bowtell, J.; Kelly, V. Fruit-Derived Polyphenol Supplementation for Athlete Recovery and Performance. *Sport Med.* **2019**, *49* (Suppl. 1), 3–23. [[CrossRef](#)]
260. Zeng, Z.; Jendricke, P.; Centner, C.; Storck, H.; Gollhofer, A.; König, D. Acute Effects of Oatmeal on Exercise-Induced Reactive Oxygen Species Production Following High-Intensity Interval Training in Women: A Randomized Controlled Trial. *Antioxidants* **2020**, *10*, 3. [[CrossRef](#)]
261. Zeng, Z.; Centner, C.; Gollhofer, A.; König, D. Effects of Dietary Strategies on Exercise-Induced Oxidative Stress: A Narrative Review of Human Studies. *Antioxidants* **2021**, *10*, 542. [[CrossRef](#)]
262. Maughan, R.J.; Burke, L.M.; Dvorak, J.; Larson-Meyer, D.E.; Peeling, P.; Phillips, S.M.; Rawson, E.S.; Walsh, N.P.; Garthe, I.; Geyer, H.; et al. IOC Consensus Statement: Dietary Supplements and the High-Performance Athlete. *Br. J. Sport Med.* **2018**, *52*, 439–455. [[CrossRef](#)]
263. Palazzetti, S.; Richard, M.J.; Favier, A.; Margaritis, I. Overloaded Training Increases Exercise-Induced Oxidative Stress and Damage. *Can. J. Appl. Physiol.* **2003**, *28*, 588–604. [[CrossRef](#)]
264. Ponce-Gonzalez, J.G.; Corral-Pérez, J.; De Villarreal, E.S.; Gutierrez-Manzanedo, J.V.; De Castro-Maqueda, G.; Casals, C. Antioxidants Markers of Professional Soccer Players during the Season and Their Relationship with Competitive Performance. *J. Hum. Kinet.* **2021**, *80*, 113–123. [[CrossRef](#)]
265. Spanidis, Y.; Goutzourelas, N.; Stagos, D.; Mpesios, A.; Priftis, A.; Bar-Or, D.; Spandidos, D.A.; Tsatsakis, A.M.; Leon, G.; Kouretas, D. Variations in Oxidative Stress Markers in Elite Basketball Players at the Beginning and End of a Season. *Exp. Ther. Med.* **2016**, *11*, 147–153. [[CrossRef](#)]
266. Lafay, S.; Jan, C.; Nardon, K.; Lemaire, B.; Ibarra, A.; Roller, M.; Houvenaeghel, M.; Juhel, C.; Cara, L. Grape Extract Improves Antioxidant Status and Physical Performance in Elite Male Athletes. *J. Sport Sci. Med.* **2009**, *8*, 468–480.
267. Lehmann, M.; Foster, C.; Keul, J. Overtraining in Endurance Athletes: A Brief Review. *Med. Sci. Sport Exerc.* **1993**, *25*, 854–862. [[CrossRef](#)]

268. Margaritelis, N.V.; Kyparos, A.; Paschalis, V.; Theodorou, A.A.; Panayiotou, G.; Zafeiridis, A.; Dipla, K.; Nikolaidis, M.G.; Vrabas, I.S. Reductive Stress after Exercise: The Issue of Redox Individuality. *Redox Biol.* **2014**, *2*, 520–528. [[CrossRef](#)]
269. Ruskovska, T.; Maksimova, V.; Milenkovic, D. Polyphenols in Human Nutrition: From the in Vitro Antioxidant Capacity to the Beneficial Effects on Cardiometabolic Health and Related Inter-Individual Variability—An Overview and Perspective. *Br. J. Nutr.* **2020**, *123*, 241–254. [[CrossRef](#)]
270. Morillas-Ruiz, J.M.; Villegas García, J.A.; López, F.J.; Vidal-Guevara, M.L.; Zafrilla, P. Effects of Polyphenolic Antioxidants on Exercise-Induced Oxidative Stress. *Clin. Nutr.* **2006**, *25*, 444–453. [[CrossRef](#)]
271. De Lima Tavares Toscano, L.; Silva, A.S.; de França, A.C.L.; de Sousa, B.R.V.; de Almeida Filho, E.J.B.; da Silveira Costa, M.; Marques, A.T.B.; da Silva, D.F.; de Farias Sena, K.; Cerqueira, G.S.; et al. A Single Dose of Purple Grape Juice Improves Physical Performance and Antioxidant Activity in Runners: A Randomized, Crossover, Double-Blind, Placebo Study. *Eur. J. Nutr.* **2020**, *59*, 2997–3007. [[CrossRef](#)]
272. Toscano, L.T.; Tavares, R.L.; Toscano, L.T.; Silva, C.S.O.d.; Almeida, A.E.M.d.; Biasoto, A.C.T.; Gonçalves, M.d.C.R.; Silva, A.S. Potential Ergogenic Activity of Grape Juice in Runners. *Appl. Physiol. Nutr. Metab.* **2015**, *40*, 899–906. [[CrossRef](#)]
273. McAnulty, S.R.; McAnulty, L.S.; Nieman, D.C.; Quindry, J.C.; Hosick, P.A.; Hudson, M.H.; Still, L.; Henson, D.A.; Milne, G.L.; Morrow, J.D.; et al. Chronic Quercetin Ingestion and Exercise-Induced Oxidative Damage and Inflammation. *Appl. Physiol. Nutr. Metab.* **2008**, *33*, 254–262. [[CrossRef](#)] [[PubMed](#)]
274. Quindry, J.C.; McAnulty, S.R.; Hudson, M.B.; Hosick, P.; Dumke, C.; McAnulty, L.S.; Henson, D.; Morrow, J.D.; Nieman, D. Oral Quercetin Supplementation and Blood Oxidative Capacity in Response to Ultramarathon Competition. *Int. J. Sport Nutr. Exerc. Metab.* **2008**, *18*, 601–616. [[CrossRef](#)] [[PubMed](#)]
275. Miyazawa, T.; Burdeos, G.C.; Itaya, M.; Nakagawa, K.; Miyazawa, T. Vitamin E: Regulatory Redox Interactions. *IUBMB Life* **2019**, *71*, 430–441. [[CrossRef](#)] [[PubMed](#)]
276. Wolf, G. The Discovery of the Antioxidant Function of Vitamin E: The Contribution of Henry A. Mattill. *J. Nutr.* **2005**, *135*, 363–366. [[CrossRef](#)] [[PubMed](#)]
277. Pilaczynska-Szczesniak, L.; Skarpanska-Steinborn, A.; Deskur, E.; Basta, P.; Horoszkiewicz-Hassan, M. The Influence of Chokeberry Juice Supplementation on the Reduction of Oxidative Stress Resulting from an Incremental Rowing Ergometer Exercise. *Int. J. Sport Nutr. Exerc. Metab.* **2005**, *15*, 48–58. [[CrossRef](#)]
278. Padayatty, S.J.; Katz, A.; Wang, Y.; Eck, P.; Kwon, O.; Lee, J.H.; Chen, S.; Corpe, C.; Levine, M.; Dutta, A.; et al. Vitamin C as an Antioxidant: Evaluation of Its Role in Disease Prevention. *J. Am. Coll. Nutr.* **2003**, *22*, 18–35. [[CrossRef](#)]
279. Rawson, E.S.; Miles, M.P.; Larson-Meyer, D.E. Dietary Supplements for Health, Adaptation, and Recovery in Athletes. *Int. J. Sport Nutr. Exerc. Metab.* **2018**, *28*, 188–199. [[CrossRef](#)]
280. Davis, J.A.; Horn, D.L.; Marr, K.A. Central Nervous System Involvement in Cryptococcal Infection in Individuals after Solid Organ Transplantation or with AIDS. *Transpl. Infect. Dis.* **2009**, *11*, 432–437. [[CrossRef](#)]
281. Lagouge, M.; Argmann, C.; Gerhart-Hines, Z.; Meziane, H.; Lerin, C.; Daussin, F.; Messadeq, N.; Milne, J.; Lambert, P.; Elliott, P.; et al. Resveratrol Improves Mitochondrial Function and Protects against Metabolic Disease by Activating SIRT1 and PGC-1 α . *Cell* **2006**, *127*, 1109–1122. [[CrossRef](#)]
282. Bailey, S.J.; Fulford, J.; Vanhatalo, A.; Winyard, P.G.; Blackwell, J.R.; DiMenna, F.J.; Wilkerson, D.P.; Benjamin, N.; Jones, A.M. Dietary Nitrate Supplementation Enhances Muscle Contractile Efficiency during Knee-Extensor Exercise in Humans. *J. Appl. Physiol.* **2010**, *109*, 135–148. [[CrossRef](#)]
283. Vanhatalo, A.; Bailey, S.J.; Blackwell, J.R.; DiMenna, F.J.; Pavey, T.G.; Wilkerson, D.P.; Benjamin, N.; Winyard, P.G.; Jones, A.M. Acute and Chronic Effects of Dietary Nitrate Supplementation on Blood Pressure and the Physiological Responses to Moderate-Intensity and Incremental Exercise. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **2010**, *299*, 1121–1131. [[CrossRef](#)]
284. Lansley, K.E.; Winyard, P.G.; Fulford, J.; Vanhatalo, A.; Bailey, S.J.; Blackwell, J.R.; DiMenna, F.J.; Gilchrist, M.; Benjamin, N.; Jones, A.M. Dietary Nitrate Supplementation Reduces the O₂ Cost of Walking and Running: A Placebo-Controlled Study. *J. Appl. Physiol.* **2011**, *110*, 591–600. [[CrossRef](#)]
285. Davis, J.M.; Murphy, E.A.; Carmichael, M.D.; Davis, B. Quercetin Increases Brain and Muscle Mitochondrial Biogenesis and Exercise Tolerance. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **2009**, *296*, 1071–1077. [[CrossRef](#)]
286. Rodgers, J.T.; Lerin, C.; Haas, W.; Gygi, S.P.; Spiegelman, B.M.; Puigserver, P. Nutrient Control of Glucose Homeostasis through a Complex of PGC-1 α and SIRT1. *Nature* **2005**, *434*, 113–118. [[CrossRef](#)]
287. Holloszy, J.O. Biochemical Adaptations in Muscle. Effects of Exercise on Mitochondrial Oxygen Uptake and Respiratory Enzyme Activity in Skeletal Muscle. *J. Biol. Chem.* **1967**, *242*, 2278–2282. [[CrossRef](#)]
288. Rodgers, J.T.; Lerin, C.; Haas, W.; Gygi, S.P.; Spiegelman, B.M.; Puigserver, P. Molecular Basis of Exercise-Induced Skeletal Muscle Mitochondrial Biogenesis: Historical Advances, Current Knowledge, and Future Challenges. *Cold Spring Harb. Perspect. Med.* **2018**, *8*, a029686. [[CrossRef](#)]
289. Dolinsky, V.W.; Jones, K.E.; Sidhu, R.S.; Haykowsky, M.; Czubyrt, M.P.; Gordon, T.; Dyck, J.R.B. Improvements in Skeletal Muscle Strength and Cardiac Function Induced by Resveratrol during Exercise Training Contribute to Enhanced Exercise Performance in Rats. *J. Physiol.* **2012**, *590*, 2783–2799. [[CrossRef](#)]
290. Kruk, J.; Hassan, B.; Enein, A.; Duchnik, E.; Marchlewicz, M. Antioxidative Properties of Phenolic Compounds and Their Effect on Oxidative Stress Induced by Severe Physical Exercise. *J. Physiol. Sci.* **2022**, *72*, 19. [[CrossRef](#)]

291. Mason, S.A.; Trewin, A.J.; Parker, L.; Wadley, G.D. Antioxidant Supplements and Endurance Exercise: Current Evidence and Mechanistic Insights. *Redox Biol.* **2020**, *35*, 101471. [[CrossRef](#)]
292. Haramizu, S.; Ota, N.; Hase, T.; Murase, T. Catechins Attenuate Eccentric Exercise-Induced Inflammation and Loss of Force Production in Muscle in Senescence-Accelerated Mice. *J. Appl. Physiol.* **2011**, *111*, 1654–1663. [[CrossRef](#)]
293. Nakazato, K.; Ochi, E.; Waga, T. Dietary Apple Polyphenols Have Preventive Effects against Lengthening Contraction-Induced Muscle Injuries. *Mol. Nutr. Food Res.* **2010**, *54*, 364–372. [[CrossRef](#)] [[PubMed](#)]
294. Swamy, M.S.L.; Naveen, S.; Singsit, D.; Naika, M.; Khanum, F. Anti-Fatigue Effects of Polyphenols Extracted from Pomegranate Peel. *Int. J. Integr. Biol.* **2011**, *11*, 69–72.
295. Nogueira, L.; Ramirez-Sanchez, I.; Perkins, G.A.; Murphy, A.; Taub, P.R.; Ceballos, G.; Villarreal, F.J.; Hogan, M.C.; Malek, M.H. (-)-Epicatechin Enhances Fatigue Resistance and Oxidative Capacity in Mouse Muscle. *J. Physiol.* **2011**, *589*, 4615–4631. [[CrossRef](#)] [[PubMed](#)]
296. Yada, K.; Roberts, L.A.; Oginome, N.; Suzuki, K. Effect of Acacia Polyphenol Supplementation on Exercise-Induced Oxidative Stress in Mice Liver and Skeletal Muscle. *Antioxidants* **2020**, *9*, 29. [[CrossRef](#)] [[PubMed](#)]
297. Braakhuis, A.J.; Hopkins, W.G.; Lowe, T.E. Effects of Dietary Antioxidants on Training and Performance in Female Runners. *Eur. J. Sport Sci.* **2014**, *14*, 160–168. [[CrossRef](#)]
298. Cases, J.; Romain, C.; Marín-Pagán, C.; Chung, L.H.; Rubio-Pérez, J.M.; Laurent, C.; Gaillet, S.; Prost-Camus, E.; Prost, M.; Alcaraz, P.E. Supplementation with a Polyphenol-Rich Extract, Perload[®], Improves Physical Performance during High-Intensity Exercise: A Randomized, Double Blind, Crossover Trial. *Nutrients* **2017**, *9*, 421. [[CrossRef](#)]
299. Davison, G.; Callister, R.; Williamson, G.; Cooper, K.A.; Gleeson, M. The Effect of Acute Pre-Exercise Dark Chocolate Consumption on Plasma Antioxidant Status, Oxidative Stress and Immunoendocrine Responses to Prolonged Exercise. *Eur. J. Nutr.* **2011**, *51*, 69–79. [[CrossRef](#)]
300. Voduc, N.; la Porte, C.; Tessier, C.; Mallick, R.; Cameron, D.W. Effect of Resveratrol on Exercise Capacity: A Randomized Placebo-Controlled Crossover Pilot Study. *Appl. Physiol. Nutr. Metab.* **2014**, *39*, 1183–1187. [[CrossRef](#)]
301. Howatson, G.; McHugh, M.P.; Hill, J.A.; Brouner, J.; Jewell, A.P.; Van Someren, K.A.; Shave, R.E.; Howatson, S.A. Influence of Tart Cherry Juice on Indices of Recovery Following Marathon Running. *Scand. J. Med. Sci. Sport* **2010**, *20*, 843–852. [[CrossRef](#)]
302. Garg, A.; Garg, S.; Zaneveld, L.J.D.; Singla, A.K. Chemistry and Pharmacology of the Citrus Bioflavonoid Hesperidin. *Phytother. Res.* **2001**, *15*, 655–669. [[CrossRef](#)]
303. Martínez-Noguera, F.J.; Marín-pagán, C.; Carlos-vivas, J.; Alcaraz, P.E. 8-Week Supplementation of 2S-Hesperidin Modulates Antioxidant and Inflammatory Status After Exercise Until Exhaustion in Amateur Cyclists. *Antioxidants* **2021**, *10*, 432. [[CrossRef](#)]
304. Kalpana, K.B.; Srinivasan, M.; Menon, V.P. Evaluation of Antioxidant Activity of Hesperidin and Its Protective Effect on H₂O₂ Induced Oxidative Damage on PBR322 DNA and RBC Cellular Membrane. *Mol. Cell Biochem.* **2009**, *323*, 21–29. [[CrossRef](#)]
305. Chen, M.; Gu, H.; Ye, Y.; Lin, B.; Sun, L.; Deng, W.; Zhang, J.; Liu, J. Protective Effects of Hesperidin against Oxidative Stress of Tert-Butyl Hydroperoxide in Human Hepatocytes. *Food Chem. Toxicol.* **2010**, *48*, 2980–2987. [[CrossRef](#)]
306. Dean, S.; Braakhuis, A.; Paton, C. The Effects of EGCG on Fat Oxidation and Endurance Performance in Male Cyclists. *Int. J. Sport Nutr. Exerc. Metab.* **2009**, *19*, 624–644. [[CrossRef](#)]
307. Magrone, T.; Russo, M.A.; Jirillo, E. Cocoa and Dark Chocolate Polyphenols: From Biology to Clinical Applications. *Front. Immunol.* **2017**, *8*, 677. [[CrossRef](#)]
308. Allgrove, J.; Farrell, E.; Gleeson, M.; Williamson, G.; Cooper, K. Regular Dark Chocolate Consumption's Reduction of Oxidative Stress and Increase of Free-Fatty-Acid Mobilization in Response to Prolonged Cycling. *Int. J. Sport Nutr. Exerc. Metab.* **2011**, *21*, 113–123. [[CrossRef](#)]
309. Cavarretta, E.; Peruzzi, M.; Del Vescovo, R.; Di Pilla, F.; Gobbi, G.; Serdoz, A.; Ferrara, R.; Schirone, L.; Sciarretta, S.; Nocella, C.; et al. Dark Chocolate Intake Positively Modulates Redox Status and Markers of Muscular Damage in Elite Football Athletes: A Randomized Controlled Study. *Oxid. Med. Cell Longev.* **2018**, *2018*, 4061901. [[CrossRef](#)]
310. Levers, K.; Dalton, R.; Galvan, E.; Goodenough, C.; O'Connor, A.; Simbo, S.; Barringer, N.; Mertens-Talcott, S.U.; Rasmussen, C.; Greenwood, M.; et al. Effects of Powdered Montmorency Tart Cherry Supplementation on an Acute Bout of Intense Lower Body Strength Exercise in Resistance Trained Males. *J. Int. Soc. Sport Nutr.* **2015**, *12*, 41. [[CrossRef](#)]
311. Sadowska-Krepa, E.; Kłapcińska, B.; Kimsa, E.; Karpiński, R. Effects of Supplementation with Red Grape Skin Polyphenolic Extract and Interval Swimming Test on the Blood Antioxidant Status in Healthy Men. *Med. Sport* **2008**, *12*, 1–7. [[CrossRef](#)]
312. Bell, P.G.; Walshe, I.H.; Davison, G.W.; Stevenson, E.; Howatson, G. Montmorency Cherries Reduce the Oxidative Stress and Inflammatory Responses to Repeated Days High-Intensity Stochastic Cycling. *Nutrients* **2014**, *6*, 829–843. [[CrossRef](#)]
313. Bell, P.G.; Walshe, I.H.; Davison, G.W.; Stevenson, E.J.; Howatson, G. Recovery Facilitation with Montmorency Cherries Following High-Intensity, Metabolically Challenging Exercise. *Appl. Physiol. Nutr. Metab.* **2015**, *40*, 414–423. [[CrossRef](#)] [[PubMed](#)]
314. McCormick, R.; Peeling, P.; Binnie, M.; Dawson, B.; Sim, M. Effect of Tart Cherry Juice on Recovery and next Day Performance in Well-Trained Water Polo Players. *J. Int. Soc. Sport Nutr.* **2016**, *13*, 4–11. [[CrossRef](#)] [[PubMed](#)]
315. Nieman, D.C.; Gillitt, N.D.; Knab, A.M.; Shanely, R.A.; Pappan, K.L.; Jin, F.; Lila, M.A. Influence of a Polyphenol-Enriched Protein Powder on Exercise-Induced Inflammation and Oxidative Stress in Athletes: A Randomized Trial Using a Metabolomics Approach. *PLoS ONE* **2013**, *8*, e72215. [[CrossRef](#)] [[PubMed](#)]

316. Ortiz-Franco, M.; Planells, E.; Quintero, B.; Acuña-Castroviejo, D.; Rusanova, I.; Escames, G.; Molina-López, J. Effect of Melatonin Supplementation on Antioxidant Status and DNA Damage in High Intensity Trained Athletes. *Int. J. Sport Med.* **2017**, *38*, 1117–1125. [[CrossRef](#)] [[PubMed](#)]
317. Lotito, S.B.; Frei, B. Consumption of Flavonoid-Rich Foods and Increased Plasma Antioxidant Capacity in Humans: Cause, Consequence, or Epiphenomenon? *Free Radic. Biol. Med.* **2006**, *41*, 1727–1746. [[CrossRef](#)]
318. Institute of Medicine (US) Panel on Dietary Antioxidants and Related Compounds. *Dietary Reference Intakes for Vitamin C, Vitamin E, Selenium, and Carotenoids*; National Academies Press: Washington, DC, USA, 2000. [[CrossRef](#)]
319. Williamson, G.; Holst, B. Dietary Reference Intake (DRI) Value for Dietary Polyphenols: Are We Heading in the Right Direction? *Br. J. Nutr.* **2008**, *99* (Suppl. 3), 55–58. [[CrossRef](#)]
320. Powers, S.K.; DeRuisseau, K.C.; Quindry, J.; Hamilton, K.L. Dietary Antioxidants and Exercise. *J. Sport Sci.* **2004**, *22*, 81–94. [[CrossRef](#)]
321. Katerji, M.; Filippova, M.; Duerksen-Hughes, P. Approaches and Methods to Measure Oxidative Stress in Clinical Samples: Research Applications in the Cancer Field. *Oxid. Med. Cell Longev.* **2019**, *2019*, 1279250. [[CrossRef](#)]
322. Tsao, R. Chemistry and Biochemistry of Dietary Polyphenols. *Nutrients* **2010**, *2*, 1231–1246. [[CrossRef](#)]
323. Chen, J.; Yang, J.; Ma, L.; Li, J.; Shahzad, N.; Kim, C.K. Structure-Antioxidant Activity Relationship of Methoxy, Phenolic Hydroxyl, and Carboxylic Acid Groups of Phenolic Acids. *Sci. Rep.* **2020**, *10*, 2611; Erratum in *Sci. Rep.* **2020**, *10*, 5666. [[CrossRef](#)]
324. Decroix, L.; Tonoli, C.; Soares, D.D.; Descat, A.; Driessens, M.J.; Weseler, A.R.; Bast, A.; Stahl, W.; Heyman, E.; Meeusen, R. Acute Cocoa Flavanols Intake Has Minimal Effects on Exercise-Induced Oxidative Stress and Nitric Oxide Production in Healthy Cyclists: A Randomized Controlled Trial. *J. Int. Soc. Sport Nutr.* **2017**, *14*, 28. [[CrossRef](#)]
325. Yarahmadi, M.; Askari, G.; Kargarfard, M.; Ghiasvand, R.; Hoseini, M.; Mohamadi, H.; Asadi, A. The Effect of Anthocyanin Supplementation on Body Composition, Exercise Performance and Muscle Damage Indices in Athletes. *Int. J. Prev. Med.* **2014**, *5*, 1594–1600.
326. Mcanulty, L.S.; Miller, L.E.; Hosick, P.A.; Utter, A.C.; Quindry, J.C.; Mcanulty, S.R. Effect of Resveratrol and Quercetin Supplementation on Redox Status and Inflammation after Exercise. *Appl. Physiol. Nutr. Metab.* **2013**, *38*, 760–765. [[CrossRef](#)]
327. Giuriato, G.; Venturelli, M.; Matias, A.; Soares, E.M.K.V.K.; Gaetgens, J.; Frederick, K.A.; Ives, S.J. Capsaicin and Its Effect on Exercise Performance, Fatigue and Inflammation after Exercise. *Nutrients* **2022**, *14*, 232. [[CrossRef](#)]

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.