

Research

High-velocity power training has similar effects to traditional resistance training for functional performance in older adults: a systematic review

Robert T Morrison ^a, Sue Taylor ^a, John Buckley ^b, Craig Twist ^c, Chris Kite ^{a,d,e,f}

^aChester Medical School, Faculty of Health, Medicine and Society, University of Chester, University Centre Shrewsbury, Shrewsbury, UK; ^bThe School of Allied Health Professions, Keele University, Staffordshire, UK; ^cSchool of Sport and Exercise Sciences, Liverpool John Moores University, Liverpool, UK; ^dSchool of Public Health Studies, Faculty of Education, Health and Wellbeing, University of Wolverhampton, UK; ^eWarwickshire Institute for the Study of Diabetes, Endocrinology and Metabolism (WISDEM), University Hospitals Coventry and Warwickshire NHS Trust, Coventry, UK; ^fCentre for Sport, Exercise & Life Sciences, Coventry University, Coventry, UK

KEY WORDS

Meta-analysis
Resistance training
Aging
Physical functional performance
Physical therapy

ABSTRACT

Questions: What is the effect of high-velocity power training (HVPT) compared with traditional resistance training (TRT) on functional performance in older adults? What is the quality of intervention reporting for the relevant literature? **Design:** Systematic review and meta-analysis of randomised controlled trials. **Participants:** Older adults (aged > 60 years), regardless of health status, baseline functional capacity or residential status. **Interventions:** High-velocity power training with the intent to perform the concentric phase as quickly as possible compared with traditional moderate-velocity resistance training performed with a concentric phase of ≥ 2 seconds. **Outcome measures:** Short Physical Performance Battery (SPPB), Timed Up and Go test (TUG), five times sit-to-stand test (5-STST), 30-second sit-to-stand test (30-STST), gait speed tests, static or dynamic balance tests, stair climb tests and walking tests for distance. The quality of intervention reporting was assessed with the Consensus on Exercise Reporting Template (CERT) score. **Results:** Nineteen trials with 1,055 participants were included in the meta-analysis. Compared with TRT, HVPT had a weak-to-moderate effect on change from baseline scores for the SPPB (SMD 0.27, 95% CI 0.02 to 0.53; low-quality evidence) and TUG (SMD 0.35, 95% CI 0.06 to 0.63; low-quality evidence). The effect of HVPT relative to TRT for other outcomes remained very uncertain. The average CERT score across all trials was 53%, with two trials rated high quality and four rated moderate quality. **Conclusion:** HVPT had similar effects to TRT for functional performance in older adults, but there is considerable uncertainty in most estimates. HVPT had better effects on the SPPB and TUG, but it is unclear whether the benefit is large enough to be clinically worthwhile. [Morrison RT, Taylor S, Buckley J, Twist C, Kite C (2023) High-velocity power training has similar effects to traditional resistance training for functional performance in older adults: a systematic review. *Journal of Physiotherapy* ■:■–■]

© 2023 Australian Physiotherapy Association. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

The World Health Organization describes healthy ageing as 'the process of developing and maintaining the functional ability that enables wellbeing in old age'.¹ Age-related functional decline is characterised by a loss of independence in completing activities of daily living,² which has a detrimental impact on quality of life,³ mortality⁴ and the incidence of injurious falls.⁵ Because functional decline falls under the physical domain of frailty syndrome,⁶ functional trajectories influence frailty risk, making people more vulnerable to negative health outcomes.⁷ Whilst acute functional decline can be caused by a medical event, progressive functional decline is often latent and can begin as early as the fifth decade of life.^{8–10}

Muscular strength, defined as the ability to apply force to an external resistance or object,¹¹ is a predictor of functional decline,^{12–14} deteriorating by ~15% every decade beyond 50 years of age¹⁵ and by 3.4% annually in people aged > 75 years.¹⁶ The force-velocity

relationship illustrates how angular velocity and torque determine movement at each anatomical joint.^{17,18} Functional trajectories are influenced by a loss in either force or velocity as people age,^{19,20} with movement velocity considered more essential in determining functional performance.^{21,22} Consequently, muscle power (the product of force and velocity) is a critical component of functional performance in older individuals^{23,24} that declines faster than strength over time,^{25,26} with reductions of up to 6% per year in people aged > 70 years.²⁷ Addressing deficits in strength or power through resistance training may help to mitigate age-related functional decline.^{28,29}

Resistance training refers to exercise where muscular contractions are resisted by external loads.³⁰ High-velocity power training (HVPT) is typically performed with lighter loads (0 to 60% of one-repetition maximum (1RM)) at faster concentric speeds (≤ 1 second); it aims to enhance the rate of force development and peak muscular power.^{31,32} To improve maximum strength, traditional moderate-velocity resistance training (TRT) typically employs loads < 60% of

1RM with a concentric movement duration of 2 to 3 seconds.³³ While multiple studies support the utility of both TRT and HVPT for improving functional performance in older adults,^{29,34–40} the relative effect of each training modality on functional performance is likely to be task specific.⁴¹ This is because health-related domains such as cardiorespiratory fitness, muscular strength, muscular endurance, flexibility and body composition,⁴² as well as skill-related domains such as power, agility, balance, speed, coordination and reaction time, all influence functional performance to varying degrees.⁴³ Although both TRT and HVPT can enhance muscular strength, HVPT is regarded as more helpful for activities of daily living, such as rising from a chair or balance recovery for fall prevention,^{44,45} which are important components of older individuals' quality of life, independence and health.

To date, several systematic reviews have made direct comparisons of the effectiveness of these modalities on older adults' functional performance^{24,37,46–49} but did not fully address the quality of the intervention reporting, which limits the translation and application into practice. Two previous reviews were narrative syntheses that did not quantitatively synthesise study data.^{24,37} Three additional systematic reviews included a small number of studies in their analyses (six studies,⁴⁷ three studies⁴⁶ and four studies⁴⁹); the data were possibly incomplete and the small sample sizes reduced confidence in the findings. A more recent systematic review included 14 trials, but the selection criteria excluded studies where external loading was < 60% of 1RM.⁴⁸ Although heavier loads are superior to improve maximum strength,⁵⁰ lower resistance training loads (< 60% of 1RM) still provide substantial increases in muscular strength and physical function in older adults,^{36,51} meaning that relevant studies may have been omitted. Further, the authors of that review acknowledged that the exclusion of balance outcomes was a limitation of their review. Although balance is not a functional task, it is important to include as it limits movement confidence⁵² and is part of the widely used Short Physical Performance Battery (SPPB).

This systematic review aimed to rate the quality of reporting of the exercise interventions in the included studies using a specific tool. In addition, the Grading of Recommendations Assessment, Development, and Evaluation (GRADE) method was used to rate the quality and strength of the evidence.

Therefore, the research questions for this systematic review and meta-analysis were:

1. What is the effect of HVPT compared with TRT on functional performance in older adults?
2. What is the quality of intervention reporting for the relevant literature?

Method

This systematic review was prospectively registered on PROSPERO, conducted in accordance with the Cochrane guidelines for systematic reviews of interventions,⁵³ and reported according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) 2020 statement.⁵⁴ The PRISMA-2020 reporting checklist and PRISMA-S reporting checklist for searches were utilised (see Tables 1 and 2 on the eAddenda).

Identification and selection of studies

Eligible studies had to be randomised trials (including parallel, cluster and crossover designs) and reported in English. No limits were placed on the geographical location or date of publication. Endnote X9 software^a was used to manage the study records retrieved.

Electronic searches

A comprehensive search was performed of the Cochrane Central Register of Controlled Trials (CENTRAL), PubMed (MEDLINE), Web of

Science, CINAHL and SportDiscus (via EBSCOhost) on 5 February 2021. All databases were searched using a customised algorithm (see Appendix 1 on the eAddenda). Additionally, a search for theses was performed using ProQuest Dissertations and Theses Global^b. Grey literature was identified using Google Scholar^c. To increase search sensitivity, citation tracking for prior and derivative papers was performed using Connected Papers^d. Finally, retraction or errata from the included studies was searched for using PubMed. Search results were imported into the Rayyan tool⁵⁵ for deduplication and study selection.

Each paper retrieved by the searches was evaluated by two reviewers. Trials were included if they met the predetermined eligibility criteria summarised according to the Population, Intervention, Comparator, Outcome, and Study (PICOS) model⁵⁴ (Box 1). The initial screening of title and abstract classified papers as eligible, ineligible or potentially eligible. Where a reviewer could not exclude a paper from the title and abstract or information was missing, two reviewers inspected the full-text version of the paper and discussed relevant points. Any disagreements were resolved through arbitration by a third member of the review team. The selection process was recorded using the PRISMA flow diagram (Figure 1), with the included studies summarised in Table 3; further details are available in Appendix 2 on the eAddenda.

Assessment of characteristics of studies

Risk of bias

The risk of bias for each included study was evaluated using version 2 of the Cochrane risk of bias tool.⁵⁶ Bias was assessed based on five domains: randomisation process (selection bias), deviations from intended interventions (performance bias), missing outcome data (attrition bias), measurement of the outcome (detection bias) and selection of the reported result (reporting bias).

Each potential source of bias was graded as high, low or unclear, along with a justification for each decision in a 'Risk of Bias' table. Summary judgements were categorised as low risk, some concerns or high risk, according to the Cochrane guidelines.⁵⁶ Blinding of participants is unfeasible in exercise interventions and all studies scored 'high' on this component; however, this was excluded when considering overall performance bias risk.

Two reviewers assessed each trial independently with discrepancies arbitrated by a third team member. Cohen's κ was used to determine inter-rater agreement. Where study details were inadequate, study authors were contacted for clarification.

Reporting quality

Intervention reporting quality was assessed using the Consensus on Exercise Reporting Template (CERT).⁵⁷ The CERT was developed to establish a consensus on reporting exercise interventions that would not only improve transparency of the research, but also allow exercise interventions to be replicated and implemented into clinical care.⁵⁸ It provides guidance on 16 items across seven domains that are required to make exercise interventions replicable. Specific domains include materials, provider, delivery, location, dosage, tailoring and compliance. Individual trials were scored on this 16-item checklist with a maximum score of 19. The CERT has previously been implemented by grading studies based on percentage of total score.⁵⁹ Studies satisfying > 75% of criteria were considered to have a high level of reporting, 60 to 74% as moderate and < 60% as low.

Study design

All trials that met the inclusion criteria (Box 1) and reported quantitative data in their analysis were included. Whilst trials must have compared interventions of HVPT with TRT, data were also included where the same additional interventions were completed concurrently in both groups.

Box 1. Inclusion criteria.

Design

- Randomised controlled trial

Participants

- Adults aged > 60 years

Intervention

- High-velocity power training
- Concentric phase performed as quickly as possible

Comparator

- Traditional moderate-velocity resistance training
- Concentric cadence ≥ 2 seconds

Outcome measures

- Objective measures of functional performance

Participants

This review included adults aged > 60 years regardless of health status, baseline functional capacity or residential status (eg, community dwelling or institutionalised).

Interventions

Trials were included where they had an intervention group allocated HVPT and a comparison group allocated TRT. Where other groups existed, only data pertaining to the HVPT and TRT groups were considered. For the purposes of this review, HVPT must have been

used with the intent to perform the concentric portion of an exercise as quickly as possible, whereas TRT was any training protocol without the explicit intent to maximise concentric velocity.

Outcome measures

Trials had to report an objective measure of functional performance. Primary outcome measures were the Short Physical Performance Battery (SPPB), Timed Up and Go test (TUG), five times sit-to-stand test (5-STST), 30-second sit-to-stand test (30-STST), gait speed tests, stair climb tests, walking tests for distance, and static and dynamic balance tests.

Data analysis

Appendix 2 on the eAddenda details the specific items for data extraction. A standardised form was devised in Microsoft Excel^e to extract baseline and post-intervention measures. One reviewer extracted data and a second reviewer independently cross-checked the accuracy of data extraction for all papers. Subsequently, one reviewer entered study data into the Cochrane Collaboration's Review Manager 5.4 software.^f

Where multiple publications existed for the same trial, these were linked together. The initial paper was used as the primary reference; however, data were extracted from all publications to obtain the maximum information possible. Where change-from-baseline data were not reported, the Cochrane Revman calculator^f was used to calculate change scores. In cases where the standard deviation (SD) of the change score was not reported, a standard equation⁵³ was used to calculate them in Microsoft Excel. Where possible, the effects of interventions on all continuous outcome data were presented as a mean difference (MD) and 95% confidence interval (95% CI). Where studies used different scales to measure the same outcome, data were presented as a standardised mean difference (SMD) and 95% CI. A sensitivity analysis was performed in these cases to remove studies with different scales and present the remaining studies as MD. Imperial units were converted to metric units where applicable.

Only data taken directly before and after the intervention were used. For example, de-training phases at follow-up were excluded as the focus of this review was on intervention effectiveness rather than residual effects. In the case of trials reporting multiple time points, the longest period of intervention was used. In the event of missing data, two attempts were made to contact study authors to obtain missing data, with a response waiting time of 6 weeks from the first contact attempt. Where data could not be obtained,^{60,61} a sensitivity analysis was intended to be performed to assess the impact of these studies, but only if it were thought that this would introduce significant bias.

Statistical heterogeneity was assessed by both visual inspection of forest plots and formal statistical tests. For visual inspection, heterogeneity was ascertained by the overlap of CIs for each study, with little or no overlap indicating substantial heterogeneity. Formal assessment used the chi-square test to determine whether differences in results were due to chance alone. Heterogeneity was then quantified by the I^2 statistic, which shows the percentage of variation across the studies resulting from heterogeneity and not chance.⁵³ In accordance with Cochrane guidelines, I^2 values of 0 to 40% were interpreted as low, 30 to 60% as moderate, 50 to 90% as substantial and 75 to 100% as considerable heterogeneity.⁵³ In cases of moderate, substantial or considerable heterogeneity, study effects were inspected, and an attempt was made to explain heterogeneity via removal of outlier studies and subgroup analysis. To explore publication and small study biases, a funnel plot and Egger's test for asymmetry was used where at least 10 different-sized studies were pooled.⁶²

Where studies were judged to be sufficiently similar, a meta-analysis was conducted by grouping continuous outcome data, with results presented as forest plots. A random-effects model was chosen as it was deemed more generalisable to the wider population.⁶³ Change from baseline scores were the primary focus to account for between-group variability. Post-intervention group differences were included to assess whether the intervention produced a




Figure 1. Flow of studies through the review.

Table 3
Characteristics of included studies.

Trial	Participants	Intervention			Outcome measures
		Regimen	HVPT	TRT	
Balachandran 2010 ⁷⁵	n = 21/17 Age: 71 y (SD 11) BMI: > 30 kg/m ² % Female: 100 Healthy	Modality: Pneumatic machines Period: 15 wk Session: 40 to 60 min Frequency: 2/wk	3 x 10 to 12 reps 50 to 80% 1RM AFAP	3 x 10 to 12 reps 70% 1RM 2 s	• SPPB (composite)
Bean 2004 ⁷⁶	n = 21/20 Age: 78 y (SD 10) BMI: < 30 kg/m ² % Female: 100 Healthy	Modality: Bodyweight/vests Period: 12 wk Session: 30 min Frequency: 3/wk	3 x 10 reps RPE < 16 AFAP	3 x 10 reps NR 2 s	• SPPB (5-STS, static balance, gait speed)
Bean 2009 ⁷⁷	n = 138/138 Age: 75 y (SD 10) BMI: < 30 kg/m ² % Female: 69 Mobility limitation	Modality: Bodyweight/vests Period: 16 wk Session: 45 to 60 min Frequency: 3/wk	2 x 10 to 12 reps RPE 11 to 16 AFAP	2 x 10 reps RPE 11 to 16 2 s	• SPPB (composite)
Bottaro 2007 ⁷⁸	n = 24/20 Age: 66 y (SD 8) BMI: < 25 kg/m ² % Female: 0 Healthy	Modality: Machines Period: 10 wk Session: NR Frequency: 2/wk	3 x 8 to 10 reps 40 to 60% 1RM AFAP	3 x 8 to 10 reps 40 to 60% 1RM 2 to 3 s	• TUG (8 ft) • 30-STS
Correa 2012 ⁷⁹	n = 58/58 Age: 67 y (SD 5) BMI: NR ^a % Female: 100 Healthy	Modality: Machines/bodyweight Period: 12 wk ^b Session: NR Frequency: 2/wk	3 to 4 x 8 to 12 reps 8RM to 12RM AFAP	3 to 4 x 8 to 12 reps 8RM to 12RM 2 s	• 30-STS
Drey 2011 ^{69,70}	n = 69/69 Age: 77 y (SD 10) BMI: < 30 kg/m ² % Female: 70 Pre-frail	Modality: Bodyspider machine Period: 12 wk ^c Session: 25 min Frequency: 2/wk	2 x 6 to 15 reps RPE 10 to 16 AFAP	2 x 6 to 15 reps RPE 10 to 16 2 to 3 s	• SPPB (5-STS, static balance, gait speed)
Englund 2017 ⁷¹	n = 26/26 Age: 65 y (SD 6) BMI: < 30 kg/m ² % Female: 62 Healthy	Modality: Multi-joint dynamometer Period: 6 wk Session: NR Frequency: 3/wk	3 x 8 reps NR 240 deg/s	3 x 8 reps NR 75 deg/s	• SPPB (composite) • TUG
Gray 2018 ⁷³	n = 99/53 Age: 81 y (SD 10) BMI: < 30 kg/m ² % Female: 72 Healthy	Modality: Machines/free-weights Period: 48 wk ^d Session: 60 min Frequency: 2/wk	3 x 10 reps 50% 1RM AFAP	3 x 10 reps 80% 1RM 2 s	• TUG (8ft) • 30-STS
Henwood 2006 ⁶⁶⁻⁶⁸	n = 67/53 Age: 70 y (SD 9) BMI: < 30 kg/m ² % Female: 56 Healthy	Modality: Machines Period: 24 wk Session: 60 min Frequency: 2/wk	3 x 8 reps 40 to 75% 1RM AFAP	3 x 8 reps 75% 1RM 3 s	• 5-STS • Static balance • 6 m walk (3 variation) • 400 m walk • Stair climb
Kelly 2016 ⁸⁰	n = 38/38 Age: 72 y (SD 10) BMI: > 30 kg/m ² % Female: 63 Total knee arthroplasty	Modality: Machines/bodyweight Period: 7 wk Session: NR Frequency: 2/wk	3 x 10 reps 50 to 80% 1RM AFAP ^e	3 x 10 reps 50 to 80% 1RM 2 s	• TU • 6-minute walk • Stair climb
Lopes 2016 ⁷²	n = 55/37 Age: 67 y (SD 11) BMI: < 30 kg/m ² % Female: 100 Healthy	Modality: Machines Period: 12 wk Session: 60 min Frequency: 3/wk	3 to 4 x 6 to 8 reps 40% 1RM AFAP	3 x 6 to 8 reps 60% 1RM 2 to 3 s	• 6-minute walk • 30-STS • TUG • Postural control
Marsh 2009 ⁸¹	n = 45/36 Age: 75 y (SD 10) BMI: > 30 kg/m ² % Female: 70 Healthy	Modality: Machines Period: 12 wk Session: 60 min Frequency: 2/wk	3 x 8 to 10 reps 40 to 70% 1RM AFAP	3 x 8 to 10 reps 40 to 70% 1RM 2 to 3 s	• SPPB (composite)
Mierzwicki 2020 ⁶⁰	n = 18/14 Age: 86 y (SD 10) BMI: NR % Female: 71 (Pre-)frail	Modality: Pneumatic machines Period: 10 wk Session: 30 to 40 min Frequency: 2/wk	2 x 15 to 20 reps 50% 1RM AFAP	2 to 3 x 5 reps 87 to 93% 1RM Slow and controlled	• TUG • 5-STS • 30-STS • 10 m fast walk
Miszko 2003 ⁸²	n = 65/39 Age: 73 y (SD 6) BMI: < 30 kg/m ² % Female: 56 Healthy	Modality: Machines Period: 16 wk ^f Session: NR Frequency: 3/wk	3 x 6 to 8 reps 40% 1RM AFAP	3 x 6 to 8 reps 80% 1RM ~ 4 s	• CS-PFP balance and endurance tests

Table 3 (Continued)

Trial	Participants	Intervention			Outcome measures
		Regimen	HVPT	TRT	
Monteiro 2019 ⁸³	N = 80/80 Age: 75 y (SD 11) BMI: < 30 kg/m ² % Female: 100 Healthy	Modality: Machines Period: 16 wk Session: NR Frequency: 3/wk	3 to 4 x 3 to 6 reps 40 to 60% 1RM < 1 s	2 to 3 x 8 to 12 reps 60 to 80% 1RM Slow to moderate	• TUG (8 ft)
Pamukoff 2014 ⁸⁴	n = 20/15 Age: 71 y (SD 5) BMI: > 30 kg/m ² % Female: 45 Healthy	Modality: Machines Period: 6 wk Session: 60 min Frequency: 3/wk	2 x 8 to 10 reps 1 x AMRAP 50% 1RM AFAP	2 x 8 to 10 reps 1 x AMRAP 50% 1RM 2 to 3 s	• Forward and lateral lean step recovery (dynamic balance)
Ramírez-Campillo 2014 ⁸⁵	n = 60/45 Age: 67 y (SD 9) BMI: > 30 kg/m ² % Female: 100 Healthy	Modality: Machines/free-weights Period: 12 wk Session: 70 min Frequency: 3/wk	3 x 8 reps 45 to 75% 1RM AFAP	3 x 8 reps 75% 1RM 3 s	• TUG (8 ft) • 30-STS • 10 m fast walk
Richardson 2019 ⁷⁴	n = 54/50 Age: 67 y (SD 12) BMI: < 30 kg/m ² % Female: 50 Healthy	Modality: Machines Period: 10 wk Session: NR Frequency: 1 and 2 /wk	3 x 14 reps 40% 1RM AFAP	3 x 7 reps 80% 1RM 2 s	• TUG (8 ft) • 30-STS • 6-min walk
Sayers 2003 ⁶¹	n = 30/25 Age: 73 y (SD 2) BMI: < 30 kg/m ² % Female: 100 Healthy	Modality: Pneumatic machines Period: 16 wk Session: NR Frequency: 3/wk	3 x 8 reps 70% 1RM AFAP	3 x 8 reps 70% 1RM 2 s	• Gait speed usual • gait speed fast • 10-STS • Stair climb • Dynamic balance
Tiggemann 2016 ⁸⁶	n = 30/25 Age: 65 y (SD 7) BMI: < 30 kg/m ² % Female: 100 Healthy	Modality: Machines Period: 12 wk Session: NR Frequency: 2/wk	2 to 3 x 8 to 15 reps RPE 13 to 18 AFAP	2 to 3 x 8 to 15 reps RPE 13 to 18 2 s	• TUG (3 m) • 5-STS • 6-min walk • Stair climb
Yoon 2017 ⁸⁷	n = 58/30 Age: 76 y (SD 6) BMI: < 25 kg/m ² % Female: 100 Mild cognitive impairment	Modality: Elastic bands Period: 12 wk Session: NR Frequency: 2/wk	2 to 3 x 12 to 15 reps RPE 12 to 13 AFAP	2 to 3 x 8 to 10 reps RPE 15 to 16 > 2 s	• SPPB (composite) • TUG (8 ft)

AFAP = as fast as possible, AMRAP = as many repetitions as possible, BMI = body mass index, CS-PFP = continuous scale physical functional performance, HVPT = high-velocity power training, NR = not reported, RCT = randomised control trial, RM = repetition maximum, RPE = rating of perceived exertion, SPPB = short physical performance battery, TRT = traditional resistance training, TUG = Timed Up and Go, 5-STS = five times sit-to-stand, 10-STS = 10 times sit-to-stand, 30-STS = 30-s sit-to-stand.

^a Assumed > 30 from body composition.

^b First 6 weeks TRT.

^c 8-week vitamin D supplementation.

^d First 24 weeks TRT.

^e Open chain exercises ≤ 1 s.

^f First 8 weeks TRT.

between-group effect where baseline group differences existed that may bias the estimate of the treatment effect.⁶⁴ For the primary comparison, data were pooled from all relevant trials stratified by functional test outcome. The results section includes meta-analyses that revealed statistically significant effects, with all remaining analyses found in Appendix 3 on the eAddenda.

The Grading of Recommendations, Assessment, Development and Evaluation (GRADE) approach was used to assess overall certainty of evidence.⁶⁵ The GRADE included the assessment of study limitations, consistency of effect, indirectness, imprecision and publication bias. All outcomes began as high certainty and were downgraded based on the GRADE criteria. Decisions were justified with notes and comments. The degree of certainty was classified as high, moderate, low or very low (Table 4).

Subgroup analysis

Post hoc subgroup analysis was performed for baseline body mass index (BMI), training frequency (< three versus ≥ three/week), age (< 70 versus ≥ 70 years) and intervention duration (< 12 versus ≥ 12 weeks). See differences between protocol and review in Appendix 4 on the eAddenda.

Sensitivity analysis

To determine the stability of the results in this systematic review, sensitivity analysis was performed by excluding studies

judged as overall high risk of bias or rated low quality by the CERT checklist.

Results

Flow of studies through the review

The search yielded 4,129 records, with 2,937 remaining after removal of duplicates. After title and abstract screening, 40 records remained, reducing to 24 after full-text analysis (Figure 1, Appendices 2 and 5 on the eAddenda). Included studies were based on 21 trials.^{60,61,66–87} Henwood et al^{66–68} and Drey et al^{69,70} presented multiple papers from the same cohort.

The authors of five trials were contacted to obtain missing outcome data.^{60,61,71–73} Unfortunately, after two attempts, there was no response from two authors.^{60,61} Neither trial reported favourable effects of HVPT over TRT on any outcome: TUG test,⁶⁰ dynamic balance, stair-climb and gait speed.⁶¹

The third paper from the Henwood et al trial⁶⁶ included a subset of participants who were re-trained after a wash-out period. As this paper was omitted from the main meta-analyses, a sensitivity analysis was conducted for each relevant outcome. This same trial⁶⁶ was the only one to report the backwards walking and 400 m walk tests. Whilst the effect for the backwards walking test was very uncertain, there was a favourable effect for HVPT in the 400 m walk test (MD

Table 4
Summary of findings table for primary outcomes: high-velocity power training versus traditional lower velocity training.

Outcomes (Change from baseline)	Difference (95% CI)	Number of participants (studies)	Certainty of the evidence (GRADE)	Comments
Short Physical Performance Battery (SPPB)	SMD 0.27 (0.02 to 0.53)	245 (6 RCTs)	⊕⊕⊕⊕ low ^{a,b}	High-velocity power training may improve SPPB performance slightly compared with lower velocity training but with some uncertainty
Timed Up and Go test (TUG)	SMD 0.35 (0.06 to 0.63)	305 (10 RCTs)	⊕⊕⊕⊕ low ^{b,c}	High-velocity power training may improve TUG performance slightly compared with lower velocity training but with some uncertainty
Five times sit-to-stand (5-STs)	SMD 0.00 (-0.70 to 0.69)	127 (4 RCTs)	⊕⊕⊕⊕ low ^{b,d}	High-velocity power training may have little to no effect on 5-STs performance compared with lower velocity training
30-second sit-to-stand test (30-STs)	MD 0.96 (-0.49 to 2.41)	179 (6 RCTs)	⊕⊕⊕⊕ low ^{b,e}	High-velocity power training may have little to no effect on 30-STs performance compared with lower velocity training
Static balance	SMD 0.36 (-0.33 to 1.04)	126 (4 RCTs)	⊕⊕⊕⊕ low ^{b,f}	High-velocity power training may have little to no effect on static balance compared with lower velocity training
Dynamic balance	MD -0.23 (-0.85 to 0.38)	41 (2 RCTs)	⊕⊕⊕⊕ very low ^{b,g}	High-velocity power training may have little to no effect on dynamic balance compared with lower velocity training
Usual gait speed	SMD -0.35 (-1.04 to 0.34)	102 (3 RCTs)	⊕⊕⊕⊕ low ^{b,h}	High-velocity power training may have little to no effect on usual gait speed compared with lower velocity training
Fast gait speed	SMD 0.08 (-0.64 to 0.79)	68 (2 RCTs)	⊕⊕⊕⊕ very low ^{b,i}	High-velocity power training may have little to no effect on fast gait speed compared with lower velocity training
Long walking tests	SMD 0.17 (-0.15 to 0.49)	153 (5 RCTs)	⊕⊕⊕⊕ low ^{b,j}	High-velocity power training may have little to no effect on long walking test performance compared with lower velocity training
Stair climb tests	SMD 0.20 (-0.27 to 0.67)	101 (3 RCTs)	⊕⊕⊕⊕ low ^{b,k}	High-velocity power training may have little to no effect on stair climb performance compared with lower velocity training

High certainty: We are very confident that the true effect lies close to that of the estimate of the effect. Moderate certainty: We are moderately confident in the effect estimate. The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different. Low certainty: Our confidence in the effect estimate is limited. The true effect may be substantially different from the estimate of the effect. Very low certainty: We have very little confidence in the effect estimate. The true effect is likely to be substantially different from the estimate of effect.

RCT = randomised controlled trial.

^a Downgraded once: five trials unclear/high risk of selection bias, four trials unclear/high risk of performance bias, three trials unclear/high risk of detection bias, and all trials unclear risk of reporting bias.

^b Downgraded once: imprecision.

^c Downgraded once: all trials unclear/high risk of selection bias, six trials unclear/high risk of performance bias, seven trials unclear/high risk of detection bias, and all trials unclear risk of reporting bias.

^d Downgraded once: three trials unclear/high risk of selection bias, three studies unclear/high risk of performance bias, two trials unclear/high risk of detection bias, and all trials unclear risk of reporting bias.

^e Downgraded once: all trials unclear/high risk of selection bias, detection bias, and reporting bias; five trials unclear/high risk of performance bias.

^f Downgraded once: three trials high risk of selection bias, three trials unclear/high risk of performance bias, one trial high risk of attrition bias, three trials unclear/high risk of detection bias, all unclear risk of reporting bias.

^g Downgraded twice: unclear or high risk of bias across all domains.

^h Downgraded once: two trials high risk of selection bias, two trials unclear risk of performance bias, two trials unclear/high risk of detection bias, and all trials unclear risk of reporting bias.

ⁱ Downgraded twice: both included trials unclear/high risk of selection bias, performance bias, detection bias, and reporting bias.

^j Downgraded once: all trials unclear/high risk of selection bias and reporting bias; four trials unclear/high risk of performance bias; five trials high risk of detection bias.

^k Downgraded once: all trials unclear/high risk of selection and reporting bias; and two trials unclear risk of performance bias.

11.36 seconds), although it was unclear whether the effect was large enough to be clinically worthwhile (95% CI 0.26 to 22.46).

One study⁷⁴ trained groups at different frequencies; these data were treated separately for meta-analysis. The SPPB data from another study⁷¹ were excluded from meta-analysis as post-intervention scores were maximum (12 points) for both HVPT and TRT groups, meaning that the magnitude of improvement beyond this score could not be ascertained. Also, the static balance outcome reported by one study⁷² was omitted due to having multiple discrete measures on different scales.

Characteristics of the included studies

The characteristics of studies included in this systematic review are summarised in Table 3 with full details in Appendix 2 on the eAddenda. This review included 21 trials with 1,055 participants. All the included studies were randomised controlled trials (RCTs). There was a total of 57 groups; nine trials had two groups (one

HVPT and one TRT),^{60,61,71,75-78,80,84,86} eight trials had three groups (two intervention and one control),^{69,72,73,79,81,82,85,87} two trials had four groups (three intervention and one control),^{66,83} and one trial had five groups (four intervention and one control).⁷⁴ The sample sizes ranged from 18⁶⁰ to 138 participants⁷⁷ randomised. Most trials (k = 19, 90%) were conducted in a university setting (gymnasium/therapy centre). An outpatient physical therapy centre was used in one trial⁸⁰ and one trial was conducted in a community gymnasium.⁸²

Risk of bias

Details of the risk of bias assessment for all included trials are found in Appendix 2 on the eAddenda. Review authors' judgements are shown in Figures 2 and 3. Inter-rater agreement for this assessment was calculated as high (Cohen's κ = 0.788, 95% CI 0.515 to 1.000). Overall, 16 (76%) of the trials were judged as having a high

	Randomisation process	Deviations from intended interventions	Missing outcome data	Measurement of the outcome	Selection of the reported result
Balachandran 2014	+	?	+	+	?
Bean 2004	+	?	+	?	?
Bean 2009	?	+	+	+	?
Bottaro 2007	+	?	+	+	?
Correa 2012	?	?	+	+	?
Drey 2011	+	+	+	+	?
Englund 2017	?	+	+	+	?
Gray 2018	?	+	+	+	?
Henwood 2006	+	?	+	+	?
Kelly 2016	?	+	+	+	?
Lopes 2016	+	+	+	+	?
Marsh 2009	+	+	+	+	?
Mierzwicki 2020	+	+	+	+	?
Miszko 2003	+	+	+	+	?
Monteiro 2019	+	+	+	+	?
Pamukoff 2014	?	+	+	+	?
Ramírez-Campillo 2014	?	+	+	+	?
Richardson 2019	?	?	+	+	?
Sayers 2003	?	?	+	+	?
Tiggemann 2016	?	?	+	+	?
Yoon 2017	?	+	?	+	?

Figure 2. Authors' judgement of methodological quality using the Cochrane Risk of Bias 2 tool.

overall risk of bias and five (24%) were deemed as having some concerns. No trial was judged as having a low overall risk of bias.

One trial was judged as low risk for selection bias,⁶⁹ 11 trials (52%) as unclear risk^{61,71,73,74,77,79,80,84-87} and nine trials (43%) were deemed




Figure 3. Risk of bias graph: review authors' judgements about each risk of bias item presented as percentages across all included studies.

to be high risk.^{60,66,72,75,76,78,81-83} In terms of performance bias, six trials (29%) were judged as low risk,^{69,71,72,77,80,83} eight trials (38%) as unclear risk^{61,66,74-76,78,79,86} and seven trials (33%) as high risk.^{60,73,81,82,84,85,87} Most trials (k = 16, 76%) were judged to be at low risk of attrition bias,^{60,61,66,69,71,72,74-81,83,86} one trial as unclear risk⁸⁷ and four as high risk.^{73,82,84,85} For detection bias, most trials (k = 13, 62%) were judged as high risk^{61,66,72-74,78,79,81-85,87} and seven trials (33%) as low risk.^{60,69,71,75,77,80,86} All trials were judged to have an unclear risk of reporting bias.

Quality of reporting

The average CERT score across all trials was 53% (Figure 4; Appendices 6 and 7 on the eAddenda). Two trials (10%) were rated 'high' quality,^{77,81} whilst four trials (19%) were rated 'moderate'.^{61,76,80,84} Common downgrade criteria included: not reporting qualifications/experience of those delivering the intervention, details for supervision, adherence, replication and study adherence/fidelity. Conversely, almost all trials (k = 18, 86%) provided detail on exercise progression and one trial did not provide an adequate description of the intervention.⁷¹

Participants

Of the 1,055 participants who were randomised, 224 were lost to follow-up, resulting in follow-up data being available for 831 participants. Across studies that reported participant sex (k = 20), 70% of the sample were female. All participants were female in nine trials^{61,72,75,76,79,83,85-87} and all male in one trial.⁷⁸

The mean participant age ranged from 65⁷¹ to 86 years.⁶⁰ Most trials (k = 17, 81%) had participants that were assumed healthy. Three trials incorporated participants who were classified as pre-frail or frail^{60,69,77} and one trial included participants who were post-total knee arthroplasty.⁸⁰

Interventions

The intervention period ranged from 6^{71,84} to 48 weeks,⁷³ with a mean period of 14 weeks. However, only part of the intervention period compared HVPT with TRT in three trials.^{73,79,82} Training frequency was twice per week in 12 trials,^{60,66,69,73,75,78-81,86,87} and three times/week in nine trials.^{61,71,72,76,77,82-85} One trial had groups training both once and twice/week⁷⁴ and these were treated separately for meta-analysis. Where reported (k = 11), session duration ranged from 25⁶⁹ to 70 minutes,⁸⁵ with the most common session duration being 60 minutes (k = 6).^{66,72,73,81,84} Session duration was not reported in 10 trials.^{61,71,74,78-80,82,83,86,87}

Most studies defined HVPT as 'a concentric velocity as fast as possible' (k = 18, 86%) and three trials defined HVPT as a 24 deg/second concentric velocity,⁷¹ < 1 second,⁸³ and ≤ 1 second,⁸⁰ respectively. TRT was performed in nine trials at a concentric velocity of 2 seconds,^{61,73-77,79,80,86} 2 to 3 seconds in five trials,^{69,72,78,81,84} 3 seconds in two trials,^{66,85} 4 seconds in one trial,⁸² > 2 seconds in one trial,⁸⁷ and 75 deg/second in one trial.⁷¹ TRT was defined in two trials



Figure 4. Summary graph of results for the reporting quality evaluation: % of trials rated by each Consensus on Exercise Reporting (CERT) domain.

as 'slow and controlled' and 'slow to moderate' concentric velocities, respectively.^{60,83}

Interventions used weight-stack resistance machines in nine trials,^{66,72,74,78,81–84,86} pneumatic resistance machines in three trials,^{60,61,75} elastic resistance bands in two trials,^{69,87} bodyweight with weighted vests in two trials,^{76,77} a multi-joint dynamometer in one trial,⁷¹ and combined interventions in four trials (Table 3).^{73,79,80,85} To determine training intensity, trials used a wide range of % 1RM or RPE scale values (Table 3). Intensity was not reported in one trial.⁷¹

Outcomes

Reported outcomes (Table 3) included the TUG test (k = 11),^{60,71–74,78,80,83,85–87} SPPB composite test score (k = 7),^{69,71,75–77,81,87} 5-STSTS (k = 5),^{60,66,69,76,86} 30-STSTS (k = 6),^{60,72–74,78,79,85} various balance tests (k = 7),^{61,66,69,72,76,82,84} endurance-oriented walking tests (k = 6),^{66,72,74,80,82,86} gait speed tests (k = 6),^{60,61,66,69,76,85} and stair climb tests (k = 4).^{61,66,80,86}

Table 5

Effect estimates and heterogeneity for change from baseline and post-intervention values.

Outcome (test)	Trials	Change scores from baseline				Post-intervention values					
		N	SMD	95% CI		I ² (%)	N	SMD	95% CI		I ² (%)
				Lower	Upper				Lower	Upper	
SPPB	69,75–77,81,87	245	0.27	0.02	0.53	0	245	0.23	-0.02	0.48	0
TUG	71–74,78,80,83,85–87	305	0.35	0.06	0.63	33	305	0.27	0.04	0.50	3
5-STSTS	66,69,76,86	127	0.00	-0.70	0.69	72	127	0.26	-0.25	0.78	50
30-STSTS	72–74,78,79,85	179	0.96 ^a	-0.49	2.41	73	179	0.86 ^a	0.08	1.64	0
Static balance	66,69,76,82	126	0.36	-0.33	1.04	70	126	0.30	-0.15	0.76	36
Dynamic balance	72,84	15	-0.48	-1.51	0.56	0	15	-0.33	-1.36	0.69	0
Usual gait	66,69,76	102	-0.35	-1.04	0.34	64	102	-0.09	-0.80	0.61	66
Fast gait	66,85	68	0.08	-0.64	0.79	55	68	0.03	-0.63	0.69	47
Long walking tests	72,74,80,82,86	153	0.17	-0.15	0.49	0	153	0.32	-0.00	0.64	0
Stair climb	66,80,86	101	0.20	-0.27	0.67	29	101	-0.24	-0.64	0.15	0

SPPB = short physical performance battery, TUG = Timed Up and Go test, 5-STSTS = five times sit-to-stand, 30-STSTS = 30-second sit-to-stand.

^a mean difference.

Effects of interventions

Ten outcomes were assessed when HVPT was compared with TRT for functional performance in older adults (Tables 4 and 5). The estimates of effect were very uncertain for the 5-STSTS, static balance, dynamic balance, usual gait speed, fast gait speed, long walking, and stair climb outcomes for change from baseline, or comparison of post-intervention values (Table 5, and Figures 5 to 11 in Appendix 3 on the eAddenda). Findings for the 30-STSTS showed comparable change scores although post-intervention values revealed an effect in favour of HVPT (Figure 12 in Appendix 3 on the eAddenda). A sensitivity analysis to include additional data from Henwood et al⁶⁶ for the 5-STSTS, static balance, usual gait speed, fast gait and stair climb outcomes did not influence results (Figures 13 to 17 in Appendix 3 on the eAddenda). The removal of Drey et al,⁶⁹ who used a different 5-STSTS scale, did not influence results when pooled as a MD (MD 0.36 seconds, 95% CI -1.27 to 1.98; three trials, 83 participants, I² = 85%; Figure 18 in Appendix 3 on the eAddenda). Sub-group analyses revealed a favourable effect for HVPT in the TUG, 30-STSTS and static balance outcomes for higher training frequencies (Appendix 8 on the eAddenda).




Figure 19. Forest plot showing the effect of high-velocity power training (HVPT) relative to traditional resistance training (TRT) as change scores for (A) the short physical performance battery (SPPB) and (B) Timed Up and Go test (TUG).

There were weak to moderate effects for change from baseline scores favouring HVPT for the SPPB and TUG outcomes. Figure 19 depicts simplified forest plots for these meta-analyses. Detailed forest plots are found in Figure 20 in Appendix 3 on the eAddenda.

For the SPPB, results were based on pooled SMD as one study utilised a modified scale.⁷⁵ Meta-analysis revealed a weak to moderate effect in favour of HVPT on change from baseline SPPB score (SMD 0.27, 95% CI 0.02 to 0.53; six trials, 245 participants, $I^2 = 0\%$; low-certainty evidence; Table 5; Figure 19A; Figure 20 in Appendix 3 on the eAddenda). In sensitivity analyses, this effect was not retained following removal of trials rated as having a high overall risk of bias or removal of trials rated 'low' (< 60%) on the CERT checklist (Figures 21 and 22 in Appendix 3 on the eAddenda). A sensitivity analysis was performed because Balachandran et al⁷⁵ used a circuit training protocol for HVPT but not TRT. The removal of this trial, with the results pooled as a MD, revealed an uncertain effect (MD 0.33, 95% CI 0.00 to 0.67; five trials, 228 participants, $I^2 = 0\%$; Figure 23 in Appendix 3 on the eAddenda).

Two variants of the TUG test were reported, so results were based on pooled SMD. There was a weak to moderate effect in favour of HVPT for TUG change from baseline scores (SMD 0.35, 95% CI 0.06 to 0.63; 10 trials, 305 participants, $I^2 = 33\%$; low-certainty evidence; Table 5; Figure 19B, Figure 24 in Appendix 3 on the eAddenda). This was supported by a weak to moderate effect for HVPT in comparison of post-intervention values (SMD 0.27, 95% CI 0.04 to 0.50, $I^2 = 3\%$; low-certainty evidence; Figure 24 in Appendix 3 on the eAddenda). Sensitivity analyses revealed no effect for TUG change scores or post-intervention values when trials rated as having a high overall risk of bias or trials rated 'low' on the CERT checklist were removed (Figures 25 and 26 in Appendix 3 on the eAddenda). The largest outlier was removed to explore heterogeneity in the TUG change




Figure 27. Funnel plot for the Timed Up and Go test outcome.

score analysis. This resulted in the I^2 statistic being reduced to 0% with the effect maintained (SMD 0.42, 95% CI 0.18 to 0.66). There was no indication of publication and small studies bias. Visual inspection of a funnel plot did not find marked asymmetry and Egger's test did not show asymmetry (Figure 27).

Discussion

This systematic review and meta-analysis provides evidence that HVPT may be as effective as TRT for functional performance in older adults, but there is still considerable uncertainty. Twenty-one RCTs were included and 19 (1,007 participants) were meta-analysed. However, most trials lacked pre-registration and were judged to have a high overall risk of bias with poor intervention reporting quality, resulting in the quality of evidence being classified as low to very low. Although meta-analyses for two global tests of functional performance, the SPPB and TUG tests, showed weak to moderate effects in favour of HVPT, these estimates were imprecise with some uncertainty.

All of the reported outcomes rely on the integration of several health and skill-related domains,⁸⁸ thus the results may have been influenced by changes in rate of force development, maximal strength or a combination of both.^{88–90} Accordingly, individual components of global tests of functional performance should be reported where effects of muscle power and strength are being investigated, particularly where tests incorporate disparate movements.⁹¹ However, as functional performance is a multidimensional concept, it is not possible to identify the relative contribution of power or strength to each individual test component.^{92,93} The majority of included trials reported positive changes from baseline for both interventions, suggesting that each intervention may provide a distinct benefit. This implies that, in terms of strength or power, functional performance cannot be limited to an either/or strategy and that a combined approach may produce the best results. Notably, this is consistent with the American College of Sport Medicine position stand for resistance training progression in older adults,³³ which recommends the concurrent performance of both HVPT (one to three sets, 30 to 60% 1RM, 12 to 15 repetitions) and TRT (one to three sets, 60 to 80% 1RM, 8 to 12 repetitions). However, current global physical activity guidelines do not recommend a specific dose of power training, although they recognise the role of multicomponent physical activity involving aerobic, balance and resistance training activities.^{94–96}

Training specificity, which is often overlooked in trials comparing HVPT and TRT, was an important factor that limited interpretation of the findings of this systematic review. While training adaptation occurs at the specific speeds performed,⁹⁷ this remains a contentious issue.⁹⁸ Some authors suggest that the intent to move faster, even at slower actual velocities (eg, TRT), may provide comparable changes in rate of force development.^{97,99} Yet, focusing on training that mirrors the demands of a certain task improves functional performance to a greater extent.^{100–102} The specific physiological adaptations that occur during functional movement training can be attributed to the Specific Adaptation to Imposed Demands principle¹⁰³ and are affected by Fitts law.¹⁰⁴ As a result, muscle power is compromised until task mastery has occurred.¹⁰⁵ This concept has been demonstrated by improvements in rate of force development over a single practice session.¹⁰⁶ Only three trials in this review incorporated some outcome practice during the intervention period;^{76,80,85} one of these used specific sit-to-stand training with weighted vests, reporting a large effect size in favour of HVPT.⁷⁶ Moreover, strength can be developed in a generalised manner and transfers the capacity to generate force across tasks, whereas skill transfer (ie, power) is limited.¹⁰⁷ Given the lack of training specificity across all outcomes, it is plausible that the transfer of strength developed from either HVPT or TRT explains their comparable effect. Adaptations to HVPT may have been attenuated by a lack of task-specific practice to maximise rate of force development.¹⁰⁸ Hypothetically, higher velocity training may have a role for specific 'functional training';¹⁰⁹ purpose-driven exercise that mirrors activities of daily living.¹¹⁰ This type of application is important for older adults' quality of life, mobility and independence;¹¹¹ however, data on this topic remain scarce.¹¹²

Strength curves based on the muscles' force-angle (torque) relationship govern the mechanical loading of exercise movements,¹¹³ classified as: ascending, descending and bell-shaped.¹¹⁴ Training protocols used in the trials included pneumatic machines, weight-stack machines, elastic resistance, body weight and free weights. Elastic resistance and free weights have different strength curves than resistance machines and are likely to provide a unique training stimulus,^{115,116} with pneumatic machines being shown to improve movement velocities when compared with free weights.¹¹⁷ Consequently, it remains unclear whether a particular method of training had a greater influence on the results.

There were some limitations to this systematic review and meta-analysis. Study selection criteria meant that other interventions with a power component such as plyometric training or jump training were omitted. Despite a comprehensive search, it is possible that some relevant literature was missed. Only studies published in English were included, meaning that some publications in other languages may have been missed. Version 2 of the Cochrane risk of bias tool has poor to modest inter-rater reliability, meaning that the risk of bias findings should be interpreted with caution. There were some differences between the study protocol and the review related to the definition of HVPT, and pre-defined subgroup and sensitivity analyses. These changes did not influence the overall results (see Appendix 4 on the eAddenda).

Several factors may influence the overall completeness of the evidence. Limited follow-up in primary studies means that this systematic review included only baseline and immediately post-intervention data and did not seek to evaluate the retained effects of either training modality. Factors outside of functional performance outcomes, such as quality of life, were not considered. There is also a paucity of pragmatic studies that examine real-world outcomes outside of a university or therapy clinic. Most studies used supervised group training sessions that may have influenced the participants' effort as per the Hawthorne effect.¹¹⁸ The generalisability of the findings were limited by most participants being female, reportedly healthy and without functional impairment at baseline. Although a precise threshold has not been identified,^{119,120} participants were possibly above the threshold required to see meaningful differences from resistance training, as seen in the relatively small change scores. A lack of functional impairment at baseline across outcomes precluded further subgroup analysis in this regard. For example, the

mean baseline values in all studies reporting the SPPB outcome were > 7, which is considered a threshold for functional impairment.¹²¹ Individuals with greater functional impairment can still increase muscle strength and power and may benefit the most from resistance training.¹²² It is unknown whether TRT and HVPT may have comparable effects in this population.

To reduce the risk of bias in future trials, pre-registration, disclosure of the randomisation process and blinding of assessors should be performed. Moreover, the lack of participants with low physical functioning highlights the need to include a wider range of baseline functional capacity in trials. Given the diverse interventions, there is a need to conduct more studies using similar protocols for machines, elastic bands or task-specific training, combined with more specific training for outcomes. This would ensure a consistency of training stimulus and enable a better comparison between HVPT and TRT. Finally, the value of either modality may not be fully revealed from a focus on specific functional outcomes; therefore, a broader range of outcomes should be considered for future research. Future trials should compare the effects of HVPT and TRT on quality of life, exercise-related perceived exertion and exercise adherence in older adults.

In conclusion, this review discovered that there is low to very low quality evidence that the efficacy of HVPT may be equivalent to TRT protocols for functional performance in older adults, but the true effect remains uncertain due to the high degree of imprecision. There is currently insufficient evidence to recommend HVPT over TRT in practice. Across the included studies, there was a lack of training specificity and diverse protocols, which may have diluted the true effect of either modality. Future research is required to determine whether specific and/or combined protocols may be more favourable with either of these individual approaches for enhancing functional performance in older individuals.

What is already known on this topic: Functional performance is a vital component of quality of life, independence and health in older adults. Both higher and lower velocity resistance training are viable options to improve functional performance.
What this study adds: Resistance training at both higher and lower velocities are similarly effective in improving older adults' functional performance, although the estimated difference in effect was uncertain. High-velocity power training had better effects than traditional resistance training on the Short Physical Performance Battery and Timed Up and Go test, but it is unclear whether these benefits are large enough to be clinically worthwhile.

Footnotes: ^a Endnote. Version Endnote X9. Philadelphia, PA: Clarivate Analytics; 2013.

^b ProQuest Dissertations and Theses Global, www.proquest.com.

^c Google Scholar, <https://scholar.google.com/>.

^d Connected Papers, www.connectedpapers.com.

^e Microsoft Excel, <https://office.microsoft.com/excel>: Microsoft; 2018.

^f Revman. London: The Cochrane Collaboration; 2020.

eAddenda: Tables 1 and 2, Appendices 1 to 8

Ethics approval: N/A.

Competing interests: Nil.

Source(s) of support: Nil.

Acknowledgements: Nil.

Provenance: Not invited. Peer reviewed.

Correspondence: Chris Kite, School of Public Health Studies, Faculty of Education, Health and Wellbeing, University of Wolverhampton, UK. Email: C.Kite2@wlv.ac.uk

References

1. World Health Organization. *World report on ageing and health*. Geneva: World Health Organization; 2015.

2. Jonkman NH, Del Panta V, Hoekstra T, Colpo M, van Schoor NM, Bandinelli S, et al. Predicting trajectories of functional decline in 60- to 70-year-old people. *Gerontology*. 2018;64:212–221.
3. Simone PM, Haas AL. Frailty, leisure activity and functional status in older adults: relationship with subjective well being. *Clin Gerontol*. 2013;36:275–293.
4. Yeh K-P, Lin M-H, Liu L-K, Chen L-Y, Peng L-N, Chen L-K. Functional decline and mortality in long-term care settings: Static and dynamic approach. *J Clin Gerontol Geriatr*. 2014;5:13–17.
5. Ward RE, Leveille SG, Beauchamp MK, Trivison T, Alexander N, Jette AM, et al. Functional performance as a predictor of injurious falls in older adults. *J Am Geriatr Soc*. 2015;63:315–320.
6. Khezrian M, Myint PK, McNeil C, Murray AD. A review of frailty syndrome and its physical, cognitive and emotional domains in the elderly. *Geriatrics*. 2017;2.
7. Morley JE, Vellas B, van Kan GA, Anker SD, Bauer JM, Bernabei R, et al. Frailty consensus: a call to action. *J Am Med Dir Assoc*. 2013;14:392–397.
8. Brown RT, Diaz-Ramirez LG, Boscardin WJ, Lee SJ, Steinman MA. Functional impairment and decline in middle age: a cohort study. *Ann Intern Med*. 2017;167:761–768.
9. Wloch EG, Kuh D, Cooper R. Is the hierarchy of loss in functional ability evident in midlife? Findings from a British Birth Cohort. *PLoS One*. 2016;11:e0155815.
10. Hébert R. Functional decline in old age. *CMAJ*. 1997;157:1037–1045.
11. Siff M. *Biomechanical foundations of strength and power training*. London: Blackwell Scientific; 2001.
12. Dos Santos L, Cyrino ES, Antunes M, Santos DA, Sardinha LB. Sarcopenia and physical independence in older adults: the independent and synergic role of muscle mass and muscle function. *J Cachexia Sarcopenia Muscle*. 2017;8:245–250.
13. Francis P, McCormack W, Toomey C, Lyons M, Jakeman P. Muscle strength can better differentiate between gradations of functional performance than muscle quality in healthy 50–70 y women. *Braz J Phys Ther*. 2017;21:457–464.
14. Visser M, Deeg DJ, Lips P, Harris TB, Bouter LM. Skeletal muscle mass and muscle strength in relation to lower-extremity performance in older men and women. *J Am Geriatr Soc*. 2000;48:381–386.
15. Larsson L. Histochemical characteristics of human skeletal muscle during aging. *Acta Physiol Scand*. 1983;117:469–471.
16. Milanovic Z, Pantelic S, Trajkovic N, Sporis G, Kostic R, James N. Age-related decrease in physical activity and functional fitness among elderly men and women. *Clin Interv Aging*. 2013;8:549–556.
17. Bieryla KA, Anderson DE, Madigan ML. Estimations of relative effort during sit-to-stand increase when accounting for variations in maximum voluntary torque with joint angle and angular velocity. *J Electromyogr Kinesiol*. 2009;19:139–144.
18. Westing S, Seger J. Eccentric and concentric torque-velocity characteristics, torque output comparisons, and gravity effect torque corrections for the quadriceps and hamstring muscles in females. *Int J Sports Med*. 1989;10:175–180.
19. Pojednic RM, Clark DJ, Patten C, Reid K, Phillips EM, Fielding RA. The specific contributions of force and velocity to muscle power in older adults. *Exp Gerontol*. 2012;47:608–613.
20. Alcazar J, Rodriguez-Lopez C, Ara I, Alfaro-Acha A, Rodriguez-Gomez I, Navarro-Cruz R, et al. Force-velocity profiling in older adults: An adequate tool for the management of functional trajectories with aging. *Exp Gerontol*. 2018;108:1–6.
21. Van Roie E, Verschueren SM, Boonen S, Bogaerts A, Kennis E, Coudyzer W, et al. Force-velocity characteristics of the knee extensors: an indication of the risk for physical frailty in elderly women. *Arch Phys Med Rehabil*. 2011;92:1827–1832.
22. Dalton BH, Power GA, Vandervoort AA, Rice CL. Power loss is greater in old men than young men during fast plantar flexion contractions. *J Appl Physiol*. 2010;109:1441–1447.
23. Reid KF, Fielding RA. Skeletal muscle power: a critical determinant of physical functioning in older adults. *Exerc Sport Sci Rev*. 2012;40:4–12.
24. Byrne C, Faure C, Keene DJ, Lamb SE. Ageing, muscle power and physical function: a systematic review and implications for pragmatic training interventions. *Sports Med*. 2016;46:1311–1332.
25. Petrella JK, Kim JS, Tuggle SC, Hall SR, Bamman MM. Age differences in knee extension power, contractile velocity, and fatigability. *J Appl Physiol*. 2005;98:211–220.
26. Foldvari M, Clark M, Laviolette LC, Bernstein MA, Kaliton D, Castaneda C, et al. Association of muscle power with functional status in community-dwelling elderly women. *J Gerontol A Biol Sci Med Sci*. 2000;55:M192–M199.
27. Clark DJ, Pojednic RM, Reid KF, Patten C, Pasha EP, Phillips EM, et al. Longitudinal decline of neuromuscular activation and power in healthy older adults. *J Gerontol A Biol Sci Med Sci*. 2013;68:1419–1425.
28. Valenzuela PL, Morales JS, Pareja-Galeano H, Izquierdo M, Emanuele E, de la Villa P, et al. Physical strategies to prevent disuse-induced functional decline in the elderly. *Ageing Res Rev*. 2018;47:80–88.
29. Papa EV, Dong X, Hassan M. Resistance training for activity limitations in older adults with skeletal muscle function deficits: a systematic review. *Clin Interv Aging*. 2017;12:955.
30. Campbell MD, Rutherford ZH. The Role of Physical Activity and Exercise in Managing Obesity and Achieving Weight Loss. In: Weaver JU, ed. *Practical Guide to Obesity Medicine*. Elsevier; 2018:215–230.
31. Earles DR, Judge JO, Gunnarsson OT. Velocity training induces power-specific adaptations in highly functioning older adults. *Arch Phys Med Rehabil*. 2001;82:872–878.
32. Fielding RA, LeBrasseur NK, Cuoco A, Bean J, Mizer K, Singh MAF. High-velocity resistance training increases skeletal muscle peak power in older women. *J Am Geriatr Soc*. 2002;50:655–662.
33. Ratamess N, Alvar B, Evetoch T, Housh T, Kibler W, Kraemer W. Progression models in resistance training for healthy adults [ACSM position stand]. *Med Sci Sports Exerc*. 2009;41:687–708.
34. Liu CJ, Latham NK. Progressive resistance strength training for improving physical function in older adults. *Cochrane Database Syst Rev*. 2009;CD002759.
35. Latham N, Anderson C, Bennett D, Stretton C. Progressive resistance strength training for physical disability in older people. *Cochrane Database Syst Rev*. 2003;CD002759.
36. Raymond MJ, Bramley-Tzerefos RE, Jeffs KJ, Winter A, Holland AE. Systematic review of high-intensity progressive resistance strength training of the lower limb compared with other intensities of strength training in older adults. *Arch Phys Med Rehabil*. 2013;94:1458–1472.
37. Rice J, Keogh J. Power training: can it improve functional performance in older adults? A systematic review. *Int J Exerc Sci*. 2009;2:6.
38. Hvid LG, Strotmeyer ES, Skjødt M, Magnussen LV, Andersen M, Caserotti P. Voluntary muscle activation improves with power training and is associated with changes in gait speed in mobility-limited older adults - A randomized controlled trial. *Exp Gerontol*. 2016;80:51–56.
39. Pereira A, Izquierdo M, Silva AJ, Costa AM, Bastos E, González-Badillo JJ, et al. Effects of high-speed power training on functional capacity and muscle performance in older women. *Exp Gerontol*. 2012;47:250–255.
40. Reid KF, Martin KI, Doros G, Clark DJ, Hau C, Patten C, et al. Comparative effects of light or heavy resistance power training for improving lower extremity power and physical performance in mobility-limited older adults. *J Gerontol A Biol Sci Med Sci*. 2015;70:372–378.
41. Barry BK, Carson RG. The consequences of resistance training for movement control in older adults. *J Gerontol A Biol Sci Med Sci*. 2004;59:M730–M754.
42. Varela S, Ayán C, Cancela J. Batteries assessing health related fitness in the elderly: a brief review. *Eur Rev Aging Phys Act*. 2008;5:97–105.
43. Corbin CB, Pangrazi RP, Franks BD. Definitions: Health, fitness, and physical activity. *President's Council on Physical Fitness and Sports Research Digest*. 2000. <https://eric.ed.gov/?id=ED470696>
44. Hazell T, Kenno K, Jakobi J. Functional benefit of power training for older adults. *J Aging Phys Act*. 2007;15:349–359.
45. Caserotti P. Strength training in older adults: changes in mechanical muscle function and functional performance. *Open Sports Sci J*. 2010;3.
46. Steib S, Schoene D, Pfeifer K. Dose-response relationship of resistance training in older adults: a meta-analysis. *Med Sci Sports Exerc*. 2010;42:902–914.
47. Tschopp M, Sattelmayer MK, Hilfiker R. Is power training or conventional resistance training better for function in elderly persons? A meta-analysis. *Age Ageing*. 2011;40:549–556.
48. Orssatto LbdR, de la Rocha Freitas C, Shield AJ, Pinto RS, Trajano GS. Effects of resistance training concentric velocity on older adults' functional capacity: A systematic review and meta-analysis of randomised trials. *Exp Gerontol*. 2019;127:110731.
49. Huber A. *The Effects of Power Training for Functional Performance as Compared to Strength Training in Older Adults: A Meta-Analysis*. California State University: Fresno; 2020.
50. Schoenfeld BJ, Wilson JM, Lowery RP, Krieger JW. Muscular adaptations in low-versus high-load resistance training: A meta-analysis. *Eur J Sports Sci*. 2016;16:1–10.
51. Watanabe Y, Madarame H, Ogasawara R, Nakazato K, Ishii N. Effect of very low-intensity resistance training with slow movement on muscle size and strength in healthy older adults. *Clin Physiol Funct Imaging*. 2014;34:463–470.
52. Liu-Ambrose T, Khan KM, Donaldson MG, Eng JJ, Lord SR, McKay HA. Falls-related self-efficacy is independently associated with balance and mobility in older women with low bone mass. *J Gerontol A Biol Sci Med Sci*. 2006;61:832–838.
53. Higgins JP, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, et al. *Cochrane Handbook for Systematic Reviews of Interventions*. NJ, USA: John Wiley & Sons; 2019.
54. Page MJ, Moher D, Bossuyt P, Boutron I, Hoffmann TC, Mulrow CD, et al. PRISMA 2020 explanation and elaboration: updated guidance and exemplars for reporting systematic reviews. *BMJ*. 2021;372.
55. Ouzzani M, Hammady H, Fedorowicz Z, Elmagarmid A. Rayyan—a web and mobile app for systematic reviews. *Syst Rev*. 2016;5:1–10.
56. Sterne JA, Savović J, Page MJ, Elbers RG, Blencowe NS, Boutron I, et al. RoB 2: a revised tool for assessing risk of bias in randomised trials. *BMJ*. 2019;366.
57. Slade SC, Dionne CE, Underwood M, Buchbinder R. Consensus on exercise reporting template (CERT): explanation and elaboration statement. *Brit J Sports Med*. 2016;50:1428–1437.
58. Slade SC, Dionne CE, Underwood M, Buchbinder R, Beck B, Bennell K, et al. Consensus on exercise reporting template (CERT): modified Delphi study. *Phys Ther*. 2016;96:1514–1524.
59. Charlton PC, Drew MK, Mentiplay BF, Grimaldi A, Clark RA. Exercise interventions for the prevention and treatment of groin pain and injury in athletes: a critical and systematic review. *Sports Med*. 2017;47:2011–2026.
60. Mierzwicki JT, Fox MA, Griffith KR, Harrison KM, Holstay DK, Singley NM. Comparison of high-intensity resistance and power training programs in pre-frail and frail older adults. *Phys Occup Ther Geriatr*. 2020;38:271–282.
61. Sayers SP, Bean J, Cuoco A, LeBrasseur NK, Jette A, Fielding RA. Changes in function and disability after resistance training: does velocity matter?: a pilot study. *Am J Phys Med Rehabil*. 2003;82:605–613.
62. Sterne JA, Sutton AJ, Ioannidis JP, Terrin N, Jones DR, Lau J, et al. Recommendations for examining and interpreting funnel plot asymmetry in meta-analyses of randomised controlled trials. *BMJ*. 2011;343.
63. Borenstein M, Hedges LV, Higgins JP, Rothstein HR. A basic introduction to fixed-effect and random-effects models for meta-analysis. *Res Synth Methods*. 2010;1:97–111.
64. Clifton L, Clifton DA. The correlation between baseline score and post-intervention score, and its implications for statistical analysis. *Trials*. 2019;20:1–6.
65. GRADE-Working-Group. Grading quality of evidence and strength of recommendations. *BMJ*. 2004;328(7454):1490.
66. Henwood TR, Taaffe DR. Short-term resistance training and the older adult: the effect of varied programmes for the enhancement of muscle strength and functional performance. *Clin Physiol Funct Imaging*. 2006;26:305–313.

67. Henwood TR, Taaffe DR. Detraining and retraining in older adults following long-term muscle power or muscle strength specific training. *J Gerontol A Biol Sci Med Sci.* 2008;63:751–758.
68. Henwood TR, Riek S, Taaffe DR. Strength versus muscle power-specific resistance training in community-dwelling older adults. *J Gerontol A Biol Sci Med Sci.* 2008;63:83–91.
69. Drey M, Zech A, Freiburger E, Bertsch T, Uter W, Sieber CC, et al. Effects of strength training versus power training on physical performance in prefrail community-dwelling older adults. *J Am Geriatr Soc.* 2011;59:S66–S67.
70. Zech A, Drey M, Freiburger E, Hentschke C, Bauer JM, Sieber CC, et al. Residual effects of muscle strength and muscle power training and detraining on physical function in community-dwelling prefrail older adults: a randomized controlled trial. *BMC Geriatrics.* 2012;12:68–68.
71. Englund DA, Sharp RL, Selsby JT, Ganesan SS, Franke WD. Resistance training performed at distinct angular velocities elicits velocity-specific alterations in muscle strength and mobility status in older adults. *Exp Gerontol.* 2017;91:51–56.
72. Lopes PB, Pereira G, Lodovico A, Bento PCB, Rodacki ALF. Strength and power training effects on lower limb force, functional capacity, and static and dynamic balance in older female adults. *Rejuvenation Res.* 2016;19:385–393.
73. Gray M, Powers M, Boyd L, Garver K. Longitudinal comparison of low- and high-velocity resistance training in relation to body composition and functional fitness of older adults. *Aging Clin Exp Res.* 2018;30:1465–1473.
74. Richardson DL, Duncan MJ, Jimenez A, Juris PM, Clarke ND. Effects of movement velocity and training frequency of resistance exercise on functional performance in older adults: a randomised controlled trial. *Eur J Sport Sci.* 2019;19:234–246.
75. Balachandran A, Krawczyk SN, Potiaumpai M, Signorile JF. High-speed circuit training vs hypertrophy training to improve physical function in sarcopenic obese adults: a randomized controlled trial. *Exp Gerontol.* 2014;60:64–71.
76. Bean JF, Herman S, Kiely DK, Frey IC, Leveille SG, Fielding RA, et al. Increased Velocity Exercise Specific to Task (INVEST) training: a pilot study exploring effects on leg power, balance, and mobility in community-dwelling older women. *J Am Geriatr Soc.* 2004;52:799–804.
77. Bean JF, Kiely DK, LaRose S, O'Neill E, Goldstein R, Frontera WR. Increased velocity exercise specific to task training versus the National Institute on Aging's strength training program: changes in limb power and mobility. *J Gerontol A Biol Sci Med Sci.* 2009;64A:983–991.
78. Bottaro M, Machado SN, Nogueira W, Scales R, Veloso J. Effect of high versus low-velocity resistance training on muscular fitness and functional performance in older men. *Eur J Appl Physiol.* 2007;99:257–264.
79. Correa CS, LaRoche DP, Cadore EL, Reischak-Oliveira A, Bottaro M, Krueger LF, et al. 3 different types of strength training in older women. *Int J Sports Med.* 2012;33:962–969.
80. Kelly MA, Finley M, Lichtman SW, Hyland MR, Edeer AO. Comparative analysis of high-velocity versus low-velocity exercise on outcomes after total knee arthroplasty: a randomized clinical trial. *J Geriatr Phys Ther.* 2016;39:178–189.
81. Marsh AP, Miller ME, Rejeski WJ, Hutton SL, Kritchevsky SB. Lower extremity muscle function after strength or power training in older adults. *J Aging Phys Act.* 2009;17:416–443.
82. Miszko TA, Cress ME, Slade JM, Covey CJ, Agrawal SK, Doerr CE. Effect of strength and power training on physical function in community-dwelling older adults. *J Gerontol A Biol Sci Med Sci.* 2003;58:171–175.
83. Monteiro AM, Bartolomeu RF, Forte P, Carvalho J. The effects of three different types of training in functional fitness and body composition in older women. *J Sport Health Res.* 2019;11:289–304.
84. Pamukoff DN, Haakonssen EC, Zaccaria JA, Madigan ML, Miller ME, Marsh AP. The effects of strength and power training on single-step balance recovery in older adults: a preliminary study. *Clin Interv Aging.* 2014;9:697–704.
85. Ramirez-Campillo R, Castillo A, Carlos I, Campos-Jara C, Andrade DC, Álvarez C, et al. High-speed resistance training is more effective than low-speed resistance training to increase functional capacity and muscle performance in older women. *Exp Gerontol.* 2014;58:51–57.
86. Tiggemann CL, Dias CP, Radaelli R, Massa JC, Bortoluzzi R, Schoenell MC, et al. Effect of traditional resistance and power training using rated perceived exertion for enhancement of muscle strength, power, and functional performance. *Age.* 2016;38:42.
87. Yoon DH, Kang D, Kim HJ, Kim JS, Song HS, Song W. Effect of elastic band-based high-speed power training on cognitive function, physical performance and muscle strength in older women with mild cognitive impairment. *Geriatr Gerontol Int.* 2017;17:765–772.
88. Freiburger E, de Vreede P, Schoene D, Rydwik E, Mueller V, Frändin K, et al. Performance-based physical function in older community-dwelling persons: a systematic review of instruments. *Age Ageing.* 2012;41:712–721.
89. Orsatto LB, Bezerra ES, Schoenfeld BJ, Diefenthaler F. Lean, fast and strong: Determinants of functional performance in the elderly. *Clin Biomech.* 2020;78, 105073.
90. Wages NP, Simon JE, Clark LA, Amano S, Russ DW, Manini TM, et al. Relative contribution of muscle strength, lean mass, and lower extremity motor function in explaining between-person variance in mobility in older adults. *BMC Geriatr.* 2020;20:1–11.
91. Balachandran A, Signorile JF. How to improve reporting of the Short Physical Performance Battery scores. *J Gerontol A Biol Sci Med Sci.* 2015;70:1595–1596.
92. Rutkauskienė L, Lendraitienė E. Batteries assessing functional fitness of older people: A brief review. *J Complex Health Sci.* 2019;2:64–69.
93. Beaudart C, Rolland Y, Cruz-Jentoft AJ, Bauer JM, Sieber C, Cooper C, et al. Assessment of muscle function and physical performance in daily clinical practice. *Calcif Tissue Int.* 2019;105:1–14.
94. Piercy KL, Troiano RP, Ballard RM, Carlson SA, Fulton JE, Galuska DA, et al. The physical activity guidelines for Americans. *JAMA.* 2018;320:2020–2028.
95. Department of Health and Social Care. Physical Activity Guidelines: UK Chief Medical Officers' Report. 2019.
96. Sims J, Hill K, Hunt S, Haralambous B. Physical activity recommendations for older Australians. *Australas J Ageing.* 2010;29:81–87.
97. Behm D, Sale D. Velocity specificity of resistance training. *Sports Med.* 1993;15:374–388.
98. Kawamori N, Newton RU. Velocity specificity of resistance training: Actual movement velocity versus intention to move explosively. *Strength Cond J.* 2006;28:86.
99. Behm D. Velocity specificity in resistance training is determined by intended rather than the actual contraction velocity. Toronto: McMaster University; 1990.
100. Manini T, Marko M, VanArnam T, Cook S, Fernhall B, Burke J, et al. Efficacy of resistance and task-specific exercise in older adults who modify tasks of everyday life. *J Gerontol A Biol Sci Med Sci.* 2007;62:616–623.
101. Morrissey MC, Harman EA, Johnson MJ. Resistance training modes: specificity and effectiveness. *Med Sci Sports Exerc.* 1995;27:648–660.
102. Reilly T, Morris T, Whyte G. The specificity of training prescription and physiological assessment: A review. *J Sports Sci.* 2009;27:575–589.
103. Fox EL, Bowers RW, Foss ML. *The physiological basis of physical education and athletics.* USA: Wm. C. C Brown Pub Dubuque; 1989:204–209.
104. Fitts PM. The information capacity of the human motor system in controlling the amplitude of movement. *J Exp Psychol Gen.* 1954;47:381.
105. Corbin CB, Janz KF, Baptista F. Good health: The power of power. *J Phys Educ Recreat Dance.* 2017;88:28–35.
106. Hinder MR, Schmidt MW, Garry MI, Carroll TJ, Summers JJ. Absence of cross-limb transfer of performance gains following ballistic motor practice in older adults. *J Appl Physiol.* 2011;110:166–175.
107. Issurin VB. Training transfer: scientific background and insights for practical application. *Sports Med.* 2013;43:675–694.
108. Maffuletti NA, Aagaard P, Blazevich AJ, Folland J, Tillin N, Duchateau J. Rate of force development: physiological and methodological considerations. *Eur J Appl Physiol.* 2016;116:1091–1116.
109. Stenger L. What is functional/neuromotor fitness? *ACSM's Health Fit J.* 2018;22:35–43.
110. Thompson WR. Worldwide survey of fitness trends for 2020. *ACSM's Health Fit J.* 2019;23:10–18.
111. De Vriendt P, Peersman W, Florus A, Verbeke M, Van de Velde D. Improving health related quality of life and independence in community dwelling frail older adults through a client-centred and activity-oriented program. A pragmatic randomized controlled trial. *J Nutr Health Aging.* 2016;20:35–40.
112. Hazell T, Kenno K, Jakobi J. Functional benefit of power training for older adults. *J Aging Phys Act.* 2007;15:349–359.
113. Kulig K, Andrews JG, Hay JG. Human strength curves. *Exerc Sport Sci Rev.* 1984;12:417–466.
114. McMaster DT, Cronin J, McGuigan M. Forms of variable resistance training. *Strength Cond J.* 2009;31:50–64.
115. Melchiorri G, Rainoldi A. Muscle fatigue induced by two different resistances: Elastic tubing versus weight machines. *J Electromyogr Kinesiol.* 2011;21:954–959.
116. Flanagan SP, Salem GJ. Lower extremity joint kinetic responses to external resistance variations. *J Appl Biomech.* 2008;24:58–68.
117. Frost DM, Cronin JB, Newton RU. A comparison of the kinematics, kinetics and muscle activity between pneumatic and free weight resistance. *Eur J Appl Physiol.* 2008;104:937–956.
118. Sedgwick P, Greenwood N. Understanding the Hawthorne effect. *BMJ.* 2015;351.
119. Willen C, Sunnerhagen KS, Ekman C, Grimby G. How is walking speed related to muscle strength? A study of healthy persons and persons with late effects of polio. *Arch Phys Med Rehabil.* 2004;85:1923–1928.
120. Rantanen T, Avela J. Leg extension power and walking speed in very old people living independently. *J Gerontol A Biol Sci Med Sci.* 1997;52:M225–M231.
121. Brown JD, Lo-Ciganic W-H, Shao H, Pahor M, Manini TM. Trajectories of short physical performance battery are strongly associated with future major mobility disability: Results from the LIFE Study. *J Clin Med.* 2020;9:2332.
122. Lopez P, Pinto RS, Radaelli R, Rech A, Grazioli R, Izquierdo M, et al. Benefits of resistance training in physically frail elderly: a systematic review. *Aging Clin Exp Res.* 2018;30:889–899.