



OPEN ACCESS

EDITED BY

Gordana Kendel Jovanović,
Teaching Institute of Public Health of
Primorsko Goranska County, Croatia

REVIEWED BY

Diego Fernández Lázaro,
University of Valladolid, Spain
Tonia Vassilakou,
University of West Attica, Greece
Ioanna Kontele,
University of West Attica,
in collaboration with reviewer TV

*CORRESPONDENCE

Hadeel Ghazzawi

✉ h.ghazzawi@ju.edu.jo

RECEIVED 01 November 2023

ACCEPTED 21 December 2023

PUBLISHED 18 January 2024

CITATION

Amawi A, AlKasasbeh W, Jaradat M, Almasri A,
Alobaidi S, Hammad AA, Bishtawi T,
Fataftah B, Turk N, Saoud HA, Jarrar A and
Ghazzawi H (2024) Athletes' nutritional
demands: a narrative review of nutritional
requirements.

Front. Nutr. 10:1331854.

doi: 10.3389/fnut.2023.1331854

COPYRIGHT

© 2024 Amawi, AlKasasbeh, Jaradat, Almasri,
Alobaidi, Hammad, Bishtawi, Fataftah, Turk,
Saoud, Jarrar and Ghazzawi. This is an open-
access article distributed under the terms of
the [Creative Commons Attribution License
\(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or reproduction
in other forums is permitted, provided the
original author(s) and the copyright owner(s)
are credited and that the original publication
in this journal is cited, in accordance with
accepted academic practice. No use,
distribution or reproduction is permitted
which does not comply with these terms.

Athletes' nutritional demands: a narrative review of nutritional requirements

Adam Amawi¹, Walaa AlKasasbeh², Manar Jaradat³,
Amani Almasri³, Sondos Alobaidi³, Aya Abu Hammad³,
Taqwa Bishtawi³, Batoul Fataftah³, Nataly Turk⁴, Hassan Al Saoud¹,
Amjad Jarrar^{5,6} and Hadeel Ghazzawi^{3*}

¹Department of Exercise Science and Kinesiology, School of Sport Science, The University of Jordan, Amman, Jordan, ²Department of Physical and Health Education, Faculty of Educational Sciences, Al-Ahliyya Amman University, Amman, Jordan, ³Department of Nutrition and Food Technology, School of Agriculture, The University of Jordan, Amman, Jordan, ⁴Department of Family and Community Medicine, Faculty of Medicine, The University of Jordan, Amman, Jordan, ⁵Department of Nutrition and Health, College of Medicine and Health Sciences, United Arab Emirates University, Al Ain, United Arab Emirates, ⁶Oxford Brookes Center for Nutrition and Health, Faculty of Health and Life Sciences, Oxford Brookes University, Oxford, United Kingdom

Nutrition serves as the cornerstone of an athlete's life, exerting a profound impact on their performance and overall well-being. To unlock their full potential, athletes must adhere to a well-balanced diet tailored to their specific nutritional needs. This approach not only enables them to achieve optimal performance levels but also facilitates efficient recovery and reduces the risk of injuries. In addition to maintaining a balanced diet, many athletes also embrace the use of nutritional supplements to complement their dietary intake and support their training goals. These supplements cover a wide range of options, addressing nutrient deficiencies, enhancing recovery, promoting muscle synthesis, boosting energy levels, and optimizing performance in their respective sports or activities. The primary objective of this narrative review is to comprehensively explore the diverse nutritional requirements that athletes face to optimize their performance, recovery, and overall well-being. Through a thorough literature search across databases such as PubMed, Google Scholar, and Scopus, we aim to provide evidence-based recommendations and shed light on the optimal daily intakes of carbohydrates, protein, fats, micronutrients, hydration strategies, ergogenic aids, nutritional supplements, and nutrient timing. Furthermore, our aim is to dispel common misconceptions regarding sports nutrition, providing athletes with accurate information and empowering them in their nutritional choices.

KEYWORDS

nutrition, athletes, carbohydrate, protein, micronutrients, fat, hydration, ergogenic aids

1 Introduction

Achieving peak physical condition and optimizing athletic performance significantly hinge on the role of food (1). Optimal nutrition is instrumental in supporting physical activity, enhancing sports performance, and facilitating post-exercise recovery. Professional athletes are expected to comprehensively meet their nutritional requirements by consuming foods in appropriate quantity and quality (2).

A properly balanced diet generally provides adequate energy to sustain individuals with heightened energy needs due to physical activity, ensuring a maintained energy balance in most cases (3). However, meeting the energy needs of larger athletes with increased body weight and height, as well as athletes involved in demanding, high-volume training, can present a significant difficulty. Endurance athletes such as runners, swimmers, as well as those in sports like gymnastics, wrestling, and boxing that use dietary restriction for body composition adjustments, often face a negative energy balance (3, 4).

Sports nutrition refers to the nutrients found in sports-related foods that enhance physical capabilities. These nutrients form the foundation for meeting the metabolic requirements of athletes or bodybuilders engaged in regular physical exercise, contributing to the preservation of physical health and athletic performance (5).

The nutritional needs of athletes are primarily dictated by the demands of their activities and the goals they set to attain peak sports performance and overall health (2, 5). The age of athletes is critical, especially adolescents whose nutritional needs tend to be higher not only to cover the performance demands but due to growth demands also. Adhering to appropriate nutritional practices is crucial as it impacts nearly every bodily process, spanning from energy production to post-exercise recovery (6). Moreover, individual dietary choices may be influenced by factors such as knowledge, attitude, and the availability of nutrition-related information resources (7).

Dietitians need to consider various sport-specific factors when evaluating an athlete's nutrition requirements and objectives. These factors include rules, arena size, competition timing, match frequency, and season length, which comprises macrocycles such as preseason, competition season, and off-season. Moreover, the physical attributes and position-specific responsibilities within the sport significantly impact the dietary requirements of athletes. For instance, rugby union forwards may require greater weight and strength compared to leaner and faster backs (8). Considering these sport-specific variables, variations in physique, and distinctions in positions, individualized dietary guidance becomes crucial for athletes participating in team sports.

Nutritional knowledge is a modifiable aspect that significantly shapes dietary behaviors (9). Athletes' comprehension of sports nutrition can directly impact their food preferences, subsequently influencing their overall athletic performance (10–12). Therefore, athletes should seek reliable and certified sources for nutritional information, as this contributes to their understanding of nutrition (13, 14). A strategic approach to enhancing nutritional awareness among athletes involves the implementation of nutrition education programs (15, 16), including individual nutrition counseling, group workshops, or online educational materials (17–19).

Guidance that assists athletes in attaining adequate energy intake, achieving the right balance of macronutrients and micronutrients, and strategically timing their nutrition to enhance performance and recovery contributes to optimizing training and overall athletic performance (20). A previous review (21) revealed athletes in team sports frequently fail to meet the recommended dietary intake requirements (18, 19). Athletes who do not achieve sufficient energy intake and lack a well-balanced diet of macronutrients may encounter obstacles in both training adaptations and recovery processes (20, 22).

Inadequate energy intake can have adverse effects on an athlete's performance, resulting in a reduction of fat-free mass, compromised

immune function, decreased bone mineral density, heightened susceptibility to injuries, and an elevated occurrence of symptoms related to overtraining (20).

The field of sports nutrition has seen the publication of numerous new research papers, accompanied by the release of 17 fresh consensus statements and recommendation papers from authoritative organizations such as the IOC, ACSM, and ISSN (2, 20, 23–25). However, there is a recurrent observation of suboptimal adherence to sports nutrition guidelines among athletes (26–28).

Achieving optimal performance in sports demands meticulous attention to nutrition, where athletes must strategically examine the timing, amount, and nutritional value of their food consumption, coupled with ensuring sufficient fluid consumption. A synergistic blend of essential elements such as carbohydrates, proteins, fats, vitamins, and minerals plays a pivotal role, acting as the energy source, foundational components, and catalysts for success in the world of sports. This research paper endeavors to comprehensively examine existing literature on the nutritional needs of athletes, aiming to pinpoint and elucidate the specific nutritional demands essential for their peak performance.

Maintaining a healthy and well-balanced diet is of paramount importance for athletes and active individuals to optimize performance and overall well-being. The athlete nutrition guidelines under scrutiny in this manuscript play a pivotal role in guiding individuals to arrange their diet according to established recommendations. These guidelines not only serve the athletes themselves but also provide invaluable assistance to sports nutrition specialists in their work. The primary goal is to prevent disordered eating and eating disorders, which have become increasingly prevalent among the active population. As such, this manuscript seeks to comprehensively review current guidelines, offering a clear and updated overview. By doing so, it aims to empower athletes and active individuals to enhance their dietary practices, mitigating the risk of potential disordered eating.

2 Method

A narrative review was conducted over 6 months, spanning from November 2022 to April 2023. To ensure a comprehensive and relevant collection of studies, four prominent online databases—PubMed, Google Scholar, Scopus, and Web of Science—were meticulously explored. The search aimed to encompass a broad spectrum of articles pertaining to various facets of nutrition and its impact on athletic performance. To facilitate an exhaustive search, a combination of specific search terms and keywords was employed. These included “Intake of Carbohydrates for athletes,” “Intake of Protein for athletes,” “Intake of fat for athletes,” “Vitamins,” “Minerals,” “Hydration,” “Ergogenic Aids,” “Nutritional Supplements,” and “Nutrient Timing.” These keywords were selected to cover key areas of interest within the field of sports nutrition. The search strategy involved using different combinations of the aforementioned keywords to optimize the retrieval of pertinent literature. The review specifically included studies of an experimental nature, comprising randomized controlled trials, observational studies, case studies, and case reports conducted within groups of elite or semi-elite athletes. During the evaluation process, the authors scrutinized the articles to assess how well they addressed and aligned with the overall objectives

of the literature review. Only those articles deemed highly relevant were included in the final selection.

3 Result

3.1 Energy (Kcal) and energy availability

In sports, maintaining peak performance relies heavily on precise nutritional practices and optimal energy intake. This is crucial not only for effective recovery and fatigue prevention but also to reduce the risk of injuries and illnesses (29).

Energy availability (EA) is a key concept, representing the dietary energy available for physiological functions after subtracting exercise energy expenditure. When there's an imbalance, resulting in Low Energy Availability (LEA), athletes may face negative consequences across various physiological systems, including endocrine, cardiovascular, immune, metabolic, reproductive, and gastrointestinal (30).

Athletes persisting in a state of low energy availability may experience a range of issues, from disruptions in menstrual function to heightened focus on food, increased illness susceptibility, diminished mood, compromised performance, decreased libido, and hormonal imbalances (31).

While energy availability and energy balance seem similar, they differ fundamentally. Energy balance considers all components of energy expenditure, particularly relevant in changes induced by diet and exercise on body weight and composition. In contrast, energy availability focuses specifically on exercise energy expenditure, providing a nuanced perspective on the intricate relationship between nutrition and athletic performance (32).

3.2 Carbohydrate

A high proportion of carbohydrates in the diet can significantly enhance performance during endurance and intense training. This is achieved by increasing exogenous carbohydrate availability and storing carbohydrates, known as glycogen, in muscles and the liver (33). During training, there is a gradual depletion of endogenous carbohydrates due to energy expenditure; this dependence is contingent upon the level and length of the exercise. Rapid carbohydrate intake after exercise replenishes carbohydrate stores quickly and enhances the body's training-induced adaptation processes. Among macronutrients, carbohydrates play a particularly crucial role in athletic performance since they can be metabolized aerobically and anaerobically (34, 35). Muscle glycogen and blood glucose are key energy sources for active muscles. Achieving optimal carbohydrate intake aids recovery and optimizes glycogen stores for subsequent training sessions. The recommended carbohydrate requirement varies with training volume and intensity. Emphasize the importance of incorporating foods exhibiting a glycemic index that extends across the spectrum from low to moderate into the meal plan, including complex carbohydrates (3). Nevertheless, during challenging and intense training sessions or when meeting high carbohydrate needs becomes difficult due to the substantial bulk and fiber content of complex carbohydrates, it is permissible to incorporate concentrated, nutrient-dense sources of carbohydrates.

Additionally, low-risk supplements may be considered to meet daily requirements if necessary (3, 22).

The glycemic index functions as a tool to classify carbohydrate-containing foods according to their impact on blood glucose levels relative to glucose or white bread consumption (22). Its application in sports nutrition remains controversial, lacking clear recommendations for athletes. Studies indicate potential improvements in metabolism and substrate utilization during exercise when incorporating low glycemic index carbohydrate-containing foods in the pre-exercise meal (22). However, these findings do not consistently translate to enhanced exercise performance. Notably, the influence of glycemic index on the pre-event meal diminishes when carbohydrates are consumed during exercise (22). Therefore, choosing a low glycemic index meal before exercise might prove beneficial when restricted consuming carbohydrates during physical activity.

Endurance activities of moderate to high levels of intensity, along with resistance-based workouts, heavily depend on carbohydrates as a primary fuel source. Therefore, maintaining essential glycogen stores, around 80–100 g in the liver and 300–400 g in the muscles of the skeletal system, is crucial. Numerous studies confirm the limited nature of glycogen stores (36, 37), highlighting their pivotal role as a predominant fuel source for several hours during moderate engaging in aerobic exercise at a high level of intensity (38).

There is documented evidence that as glycogen stores decrease, it becomes harder to maintain a high pace of intense exercise (23). Consuming a snack or meal high in carbohydrates before exercise ensures optimal muscle glycogen reserves. On the other hand, low pre-exercise glycogen levels result in early fatigue, reduced training intensity, depleted muscle glycogen, impaired muscle contraction, glycogenolysis, and protein degradation (34, 39, 40).

A straightforward suggestion for maximizing internal glycogen reserves in high-performance athletes is to consume adequate carbohydrates based on the intensity and duration of their training. Generally, the suggested daily carbohydrate intake falls within the range of 5–12 g/kg of body weight. This is particularly relevant for athletes engaged during training sessions of moderate to high intensity, lasting over 12 h per week; it is recommended to target the upper limit of this range, precisely 8 to 10 g/kg of body weight daily (41). When substantial muscle damage is not present, this level of carbohydrate intake has demonstrated effectiveness in maximizing glycogen storage. Recommendations based on percentages (for instance, aiming for 60–70% of your total daily caloric intake to come from carbohydrates) have lost favor because they fail to accurately prescribe the required carbohydrate amounts for athletes with elevated caloric intake or those following restricted energy intake (33).

It is important to highlight that athletes frequently fall short of meeting the recommended levels of energy and carbs. Consequently (42), emphasizing approaches to restore glycogen reserves could be essential in preparing for optimal performance in the upcoming competition.

The synthesis of glycogen after exercise is closely tied to factors considering factors like the degree of glycogen reduction, the nature, duration, and intensity of the workout session (43). During the initial 6 h following exercise, there is an increased pace of muscle glycogen replenishment, and consuming sufficient carbohydrates within 24 h after the workout can lead to full restoration of glycogen stores (44).

Current studies suggest that having a recovery meal within 2 h after exercise, as opposed to not eating, proves effective in enhancing

recovery (45). Despite optimal conditions, complete muscle glycogen resynthesis can extend up to 24h, prompting studies to explore techniques for accelerating the pace at which muscle glycogen is restored (46). Customizing the approach based on the degree of glycogen depletion, an effective strategy for adequate glycogen resynthesis involves ingesting 1.0 to 1.5g of carbohydrates per kg of body weight per hour right after exercise and maintaining this intake at 30-min intervals for up to 6h following the workout (2, 47). On the contrary, in instances where carbohydrate (CHO) intake is postponed by 2h, studies indicate a 45% reduction in glycogen resynthesis rates (48). The absorption of glucose in the intestines might serve as a constraint on glycogen resynthesis, especially notable when a significant quantity of carbohydrates is ingested in a single bolus after exercise (49). There is a proposition that carbohydrate (CHO) supplementation administered at a rate of about 1.0g/kg/h, given at consistent intervals (15 to 60-min intervals) post-exercise, is more efficacious than a substantial single bolus in sustaining elevated blood glucose levels. This, consequently, facilitates increased muscle glycogen restoration (48).

Studies investigating glycogen resynthesis rates after varying carbohydrate consumption post-exercise yield inconsistent results (50). Van Loon et al. discovered that increasing post-exercise carbohydrate consumption—raising carbohydrate intake cyclists who consumed carbohydrates ranging from 0.8 to 1.2g per kg per hour, in terms of body weight, at 30-min intervals exhibited increased rates of muscle glycogen synthesis (16.6 vs. 35.4mmol/kg dry weight/h) following a glycogen depletion cycling trial (43). Current studies suggest that 2-h intervals may not be the most effective for promoting muscle glycogen resynthesis, especially given the rapid restoration observed within the initial 2h post-exercise (43, 51). Additional studies have documented higher rates of glycogen resynthesis with more frequent supplement ingestion compared to a single large bolus (52).

Variations in study outcomes may arise from disparities in protocols; the effectiveness of post-exercise recovery meals is influenced by factors such as the timing and intervals of consumption, the subjects' training status, and the type of carbohydrates used. Given these limitations and variations in research protocols, the current evidence indicates that an ingestion of carbohydrates ranging from 1.0 to 1.5 grams per kilogram per hour is recommended and is sufficient to achieve maximal glycogen resynthesis (2, 48, 50). Further research is essential, especially studies investigating the prolonged impacts of post-exercise nourishment and simulating a typical training program.

3.2.1 Recommended daily intake of carbohydrates

To estimate athletes' carbohydrate needs, a recommended guideline involves ingesting 3-12g of carbohydrate consumption per

kilogram of body weight each day, with the precise quantity contingent on the intensity and duration of their physical activity. Individual variations and the comfort of their digestive systems should also be considered. For approximate carbohydrate needs, please refer to Table 1 (35, 53).

The formula for calculating the daily intake of carbohydrates is as follows:

$$\text{Daily Intake of Carbohydrates} = \text{Weight in Kilograms} \times \text{Grams of Carbohydrates}$$

3.3 Protein

Preserving and optimizing skeletal muscle mass are crucial goals for individuals with athletic aspirations, whether aiming for improved performance, increased muscularity, or accelerated recovery. Moreover, the importance of skeletal muscle mass extends beyond active individuals, providing direct clinical applications and benefits. This dynamic interplay—the balance between muscle protein synthesis and breakdown—is particularly crucial for aging adults, as skeletal muscle undergoes continuous regulation (54). The loss of muscle mass occurs when there is a negative balance, indicating a higher breakdown than synthesis, resulting in a net loss. On the flip side, the accumulation of muscle occurs when the rate of synthesis outpaces that of breakdown. The interaction between physical activity and dietary elements, particularly concerning the consumption of protein and indispensable amino acids, is pivotal in governing both the construction and degradation of muscle proteins (55). Recent findings indicate that alterations in the context of physical activity and nutritional intake have a more pronounced effect on muscle protein synthesis rates (56). Therefore, changes in muscle protein synthesis rates are deemed the principal determinants of variations in muscle mass over time as a response to both exercise and nutritional influences (57).

Amino acids are the structural constituents of proteins, providing the building blocks for all tissues. For athletes, the main purpose of consuming protein following vigorous exercise or competitions is the rebuilding and restoration involving both skeletal muscle and connective tissues (25). The amount, timing, and type of protein intake all affect the extent to which muscles remodel after training. It's crucial to emphasize that protein is not the primary source of fuel for athletes (25).

Various factors, including the quantities of overall amino acids, indispensable amino acids, and BCAA concentrations, impact the anabolic effectiveness of a protein source. Additionally, factors such as protein digestibility, digestion rate, and absorption kinetics are taken into account. In the assessment of dietary protein quality, attention is often directed toward the indispensable amino acid composition offered by the protein source concerning human nutritional requirements. Additionally, its capacity for digestion, absorption, and assimilation by diverse tissues throughout the body is considered (58).

Two categories of protein determine whether they include essential amino acids: complete and incomplete protein sources. Animal protein is considered complete as it provides a comprehensive source of protein, containing all the required amino acids. In contrast, plant-based proteins are incomplete sources as they lack some essential amino acids (59).

TABLE 1 The approximate requirements for carbohydrates.

Exercise intensity	Minutes of activity per day	Recommended carbohydrate intake (g/kg)
Low	<60	3-5
Moderate	60	5-7
High	60-180	6-10
Very high	>180	8-12

As highlighted in a comprehensive review (60), biological values for prevalent plant sources typically fall within the range of 56–74, whereas various animal sources exhibit a spectrum spanning from 77 to 104 on theoretical scales ranging from 0 to 100 points. A similar discrepancy is noted regarding values of net protein utilization; plant sources typically fall within the range of 53–67, while animal sources tend to be in the range of 73–94 on a 100-point scale. The scores for Protein Digestibility Corrected Amino Acid stand out as one of the commonly utilized benchmarks for assessing protein quality (61) emerge as one of the most frequently employed metrics.

Casein, whey, and eggs achieve scores of 100 in their respective categories, serving as examples of animal protein sources, whereas red meat scores 92. In contrast, typical plant protein sources typically display Protein Digestibility Corrected Amino Acid stands values below 100, typically falling within the reported range of 45–75 (60). Except for soy protein, which has a score of 100, other plant sources generally fall below this threshold. Similarly, employing the DIAAS approach for evaluating protein quality reveals a consistent pattern: animal sources often exceed 100, in contrast, the majority of plant sources tend to be below this threshold. When analyzing the amino acid profiles of different plant-based isolates in comparison to standard proteins derived from animal sources and specimens from muscles in the human skeletal system, it becomes clear that several sources of protein derived from plants do not contain adequate quantities of specific amino acids, such as lysine and methionine levels (62). Moreover, they consistently exhibit reduced levels of indispensable and BCAA in contrast comparing animal protein sources to the amino acid composition present in human skeletal muscle. Furthermore, elements such as vary depending on factors such as the type of nutrient, individual characteristics, and the specific physiological context notably affect the nutritional quality of a protein. Concerning digestibility, compelling evidence suggests that numerous plant sources have a considerably lower digestibility rate (45–80%) compared to diverse animal proteins (>90%) (63). In essence, there is a general agreement that the transportation of amino acids to peripheral tissues from plant-based proteins is typically regarded as less efficient compared to that from animal proteins (64, 65). These distinctions are considered critical factors that influence the postprandial protein synthesis response observed in different tissues.

To promote muscle repair, remodeling, and improve post-exercise strength- and hypertrophy-related responses, it is crucial to ingest protein before, during, and after a workout (66). Consumption of protein during these periods has been associated with a favorable impact on Muscle Protein Synthesis (MPS) (67).

Prior to exercise, ingesting protein with meals within 3–4 h before the workout can assist in maintaining muscle growth and enhancing muscle recovery, especially when combined with resistance training (68). Combining amino acids with carbohydrates before exercise can lead to peak rates of MPS, although the effects of protein and amino acid feedings during this period on exercise performance are not firmly established. However, consuming carbohydrates combined with protein or essential amino acids during endurance and resistance training can have beneficial effects, including an improved anabolic hormonal status, reduced muscle damage, improved muscle cross-sectional area, and extended time to exhaustion (25).

It is important to note that protein has a limited capacity for the body to utilize as an energy source during activity, whereas

carbohydrates are the primary fuel source. Therefore, rehydration and intake of simple carbohydrates (glucose) are most important for athletes during exercise (68).

To enhance myofibrillar protein synthesis after exercise and minimize amino acid degradation, a nutritional recommendation advises incorporating 0.31 g/kg of high-quality and quickly digestible protein, like whey protein. Those pursuing this objective should strive to integrate this amount per meal (69). A mixed meal consisting of carbohydrates and protein after exercise, with a carbohydrate-to-protein ratio of approximately 4 to 1, is recommended to initiate muscle glycogen synthesis (20). Note that exercise has a sustained anabolic impact for at least 24 h. However, the ideal timing for protein ingestion depends on individual tolerance as it may diminish over time after activity (70).

3.3.1 The recommended daily intake of protein

Using the following recommendations in Table 2 may help the athlete to assess their need for protein and prevent excessive intake and keep them on track (74).

3.4 Fat

To achieve optimal performance, athletes need to consume an appropriate amount of energy not only during exercise but also during recovery. Fat oxidation primarily depends on oxygen, while carbohydrate catabolism can occur with or without oxygen. It's worth noting that the complete breakdown of glucose in the mitochondria, involving the presence of oxygen, yields more ATP (75). Consuming an adequate amount of fat is important, but high-fat or fat-loading diets are ineffective (76, 77). It is recommended to consume dietary fat between 20 and 35% of the total calorie intake for athletes, with saturated fat intake being less than 10% (76).

Athletes frequently turn to dietary supplements to boost metabolic capacity, delay the onset of fatigue, enhance muscle hypertrophy, and shorten recovery periods (76). Omega-3, a type of PUFA, acts as a structural component within cell membranes of phospholipids. Omega-3 plays a crucial role in the inflammatory response of the body (78). Among athletes, Omega-3 has been linked to the postponement of onset muscle soreness, enhancement of anaerobic endurance capacity, improvement in oxygen efficiency during aerobic exercise, support for skeletal muscle health, and mitigation of exercise-induced oxidative stress (79). It is recommended to take omega-3 after or with

TABLE 2 Recommendation of protein intake based on the healthy athlete in different types of exercise and goals.

Type of training or exercise	The recommended amount of protein is typically expressed in g/kg of body weight
Preserving and building muscle mass (20)	1.4–2.0 g/kg of body weight
During caloric deficit periods (20)	2.3–3.1 g/kg fat-free mass
Endurance exercise (71, 72)	1.2–2.0 g/kg of body weight
Strength exercise (59)	1.6–2.8 g/kg of body weight
Gain (muscle) strength (73)	1.6–2.2 g/kg of body weight

a high-fat meal for optimal absorption (79). The safe recommended dose for omega-3 is 450–900 mg/day, with a maximum recommended dose of up to 3 g per day (80).

Athletes have employed a dietary strategy of increasing the proportion of dietary fat, primarily aiming to enhance intramuscular triglyceride stores. The theory behind this approach suggests potential benefits for prolonged exercise performance while preserving glycogen stores (81). Endurance athletes, in particular, have considered and applied this strategy to improve their performance in prolonged exercises. Conversely, athletes focused on strength and power have given little thought to modifying fat intake in their training strategies.

Burke and her research team delivered compelling results, revealing that a five-day regimen of adhering to a high-fat diet, where over 65% of total calorie intake came from fat, while concurrently maintaining a low carbohydrate and a daily protein consumption of 2.5 g per kg of body weight, not only enhanced fat utilization but also enabled athletes to effectively partake in high-intensity and high-volume training sessions (82).

Moreover, sustained enhancements in fat utilization were observed even following the implementation of a regimen for carbohydrate loading aimed at replenishing muscle glycogen levels. This dietary strategy suggests that a sequence of high-fat intake followed by carbohydrate loading might establish a conducive environment, enabling skeletal muscle to oxidize more fat while maintaining sufficient muscle glycogen. Nevertheless, subsequent studies did not reveal improvements in exercise performance (83). Certainly, there was a noted decrease in the rates of muscle glycogen utilization throughout the exercise bout (84). Given the expectation that enhanced carbohydrate availability is likely to enhance power generation and exercise intensity, especially during the latter phases of prolonged exercise, these results were considered counterproductive.

While there has been extensive research on the effectiveness of high-fat diets, there is a general agreement that opting for a higher critical determinant, but rather, it depends on individual factors and the overall dietary context, an advisable approach to enhancing sports performance. In an extensive review of the literature, Johnson provided insights into how a high-fat diet affects performance in physical activities. These observations encompass (1): no definitive conclusions supporting the idea that decreasing intramuscular triglycerides adversely affects performance, a hypothesis intended to enhance performance (2); a diet rich in fats, comprising more than 46% of total calories derived from fat and less than 21% from carbohydrates, promotes fat oxidation through mechanistic adaptations. These adaptations encompass increased enzymes involved in the oxidation of fatty acids and improvements in both fatty acid transport and beta-oxidation; and (3) despite these mechanistic changes, improvements in exercise performance were not consistently observed, and in certain cases, a negative impact was evident (81).

While the suggestion of increasing dietary fat intake has been made for a positive impact on the utilization of substrates, the prevailing consensus discourages high-fat diets due to their adverse effects on performance. Instances of reduced carbohydrate utilization and gastrointestinal discomfort further reinforce the argument against such dietary approaches. Whether the negative outcomes arise from the elevated consumption of dietary fat or the probable simultaneous decrease within dietary carbohydrate, the adoption of diets rich in fats is not recommended.

3.5 Micronutrients

Micronutrients play a crucial role in sustaining life, encompassing vitamins and minerals that support well-being, development, and reproductive processes. These essential substances, required in small quantities, must be obtained through dietary intake as the human body cannot synthesize them (85).

Vitamins are categorized based on their solubility, with A, D, E, and K being fat-soluble, and B and C being water-soluble. Minerals, on the other hand, are non-organic compounds contributing to physiological operations (85). Daily physiological requirements classify minerals, with macrominerals requiring around 100 mg per day and trace elements approximately 20 mg per day for individuals' health (85).

While a nutritionally balanced diet generally provides essential micronutrients in recommended doses for regular bodily functions, the appropriateness of these guidelines for athletes is a subject of debate. Athletes often resort to micronutrient supplementation, with close to half of them incorporating vitamin or mineral supplements in their regimen, according to a meta-analysis (86).

Research findings indicate that, similar to the impact of exercise intensity, duration, and character on macronutrient needs, these factors also influence athletes' micronutrient requirements. In sports with elevated energy demands, there's a likelihood of increased micronutrient needs, though precise quantification remains challenging (85).

When athletes have heightened overall energy requirements due to their training regimen, this increased need should reflect in both macro- and micronutrient intake. Meeting this demand through a well-rounded diet aligned with recommended dietary reference intakes for vitamins and minerals is generally achievable (87). However, certain scenarios, such as substantial losses through sweat and urine or specific dietary preferences, may lead to increased vitamin and mineral requirements (85).

In instances where athletes face challenges in meeting their micronutrient needs through diet alone, supplementation may be beneficial. Athletes might consider external supplements to enhance well-being and performance, especially in situations like altitude training, where iron supplementation may be necessary (88, 89). Specific sports may present unique concerns, such as lower concentrations of vitamin D for athletes in winter sports or indoor activities (90).

Athletes consuming a diet rich in energy from nutrient-dense foods typically do not require vitamin and mineral supplements. However, those who struggle to meet their micronutrient needs may benefit from supplementation, guided by a sports nutritionist (91, 92). It's crucial to emphasize that while micronutrients are crucial for health, they do not exert ergogenic effects, and factors like low-energy diets, vegetarianism, illness, and injuries can contribute to micronutrient deficiencies (40, 93).

3.6 Hydration

It is crucial to replenish fluid loss during, before, and after exercise. Researchers have found that losing 2% of body fluids can affect performance and cognitive function. Thirst is often not an effective indicator of dehydration, as 1.5 L can be lost before thirst perception

(94). Athletes are susceptible to losing 0.3 to 2.4 L per hour of sweat, which includes not only water but also salt, potassium, calcium, magnesium, and chloride. Consequently, fluid and electrolyte replacement should be incorporated into their recovery (93).

Fluid balance is fundamental for athletes, as hypohydration, which occurs when body water levels are lower than normal due to excessive sweating during exercise or diarrhea, can have life-threatening consequences and negatively impact performance (95). On the other hand, during endurance exercise, increased consumption of mostly sodium-poor or sodium-free liquids, such as water, can lead to hyponatremia (96). Excessive fluid consumption causes fluid retention in the body, resulting in dilutional hyponatremia (97). Athletes should be aware that the body can lose fluids in various ways through sweating as a natural result of prolonged exercise, urination, and other factors such as temperature and humidity (98).

The physical signs and symptoms of dehydration include dry and poor skin turgor, dark urine color, rapid weight loss, dry and sticky mouth, weakness, fatigue, headache, sunken eyes, muscle cramps, an increased rate of injuries, difficulty in recovery, and a racing heartbeat (99, 100). Additionally, over-hydration can manifest through physical signs and symptoms such as weight gain, swelling of the feet and hands (edema), nausea and vomiting, orthopnea (sensation of breathlessness during sleep), low blood sugar, weakness, seizures, fatigue, headache, and increased urination (99, 100).

Athletes are advised to initiate a pre-training regimen in a well-hydrated state, ingesting approximately 500 to 600 mL of water or a sports beverage 2 to 3 h before engaging in exercise. An additional intake of 200 to 300 mL of water or a sports drink is recommended 10 to 20 min before the onset of exercise (101). Throughout the training session, fluid replenishment should align with sweat and urine losses, endeavoring to sustain hydration levels while minimizing body weight reduction to less than 2%. Typically, this goal can be met by consuming 200 to 300 mL of fluid at intervals of 10 to 20 min. Post-activity, the focus of hydration should be on rectifying any fluid deficits incurred during the practice or competition (102).

Sports drinks are primarily used to rehydrate and replenish essential electrolytes and other important components for athletes, such as magnesium, sodium, calcium, potassium, glucose, and fluids lost during strenuous exercise, to enhance endurance and performance (103). On the other hand, the usage of sports drinks may differ based on the nature of the exercise. The best time to use a sports drink is during high-intensity and long-duration exercise lasting more than 1 h, on hot and humid days, during heavy or salty sweating, and for recovery after training (104).

The beverage hydration index model was introduced to evaluate the hydrating potential of a drink compared to plain water when individuals are at rest. This model operates on the assumption that a beverage inducing greater diuresis than water results in less retained available fluid in the total body water pool, reflected by a beverage hydration index below 1.0. Although a recent addition to beverage metrics, akin to the glycemic index for foods, the beverage hydration index has garnered replication by various research groups (105–107) since its inception (108). Significantly, population-specific factors like body mass and sex seem to have negligible effects, and the reproducibility of the hydration index model is reported to be robust (105). Consequently, the hydration index model has gained recognition as a reliable method for evaluating beverage hydration characteristics in well-controlled conditions, particularly when

individuals are in a state of dehydration, as opposed to rehydration scenarios following exercise.

The addition of electrolytes to water seems to enhance fluid retention according to the beverage hydration index method (105, 109). Nevertheless, the minimum sodium level required to achieve this effect varies across studies, as some indicate that sports drinks with approximately 20 mmol sodium may not necessarily yield a significantly greater hydration index compared to the water control (106, 108). This aspect warrants further exploration, especially considering that sports drinks are commonly recommended for general public use as a suitable beverage for oral rehydration post-dehydration. Beverages with elevated sodium content (usually around 45 mmol), like Pedialyte, tend to exhibit a higher hydration index compared to water. However, whether Pedialyte has a hydration index superior to a sports drink remains uncertain in some studies (105, 107, 108). This aspect requires further investigation, especially considering that sports drinks are often recommended for general public use as a suitable beverage for oral rehydration after dehydration. Beverages with higher sodium content (typically around 45 mmol), such as Pedialyte, generally show a higher hydration index compared to water. However, whether Pedialyte has a hydration index superior to a sports drink remains uncertain in some studies (107, 108).

3.7 Ergogenic aids and nutritional supplements

In recent times, there has been a significant increase in attention devoted to nutritional supplements and ergogenic aids within the sports community. Nutritional supplements are concentrated reservoirs of nutrients or other compounds exhibiting nutritional or physiological qualities beyond what is naturally obtained through a regular diet (110). Ergogenic aids pertain to pharmaceutical substances employed to boost sports performance (111). Recognizing a profitable market, commercial brands cater to high-performance athletes (112), university students (113), and young amateur athletes (114) who enthusiastically supplement their diets with these products. The growing prevalence of athletes using nutritional supplements and ergogenic aids has raised concerns among health and sports authorities. A significant number of these supplements and aids have been found to be contaminated with harmful or banned substances (115). Such contamination could pose a risk to the health of athletes or lead to competition bans if the products contain prohibited doping substances (116). While the utilization of nutritional supplements and ergogenic aids is common among athletes ranging from recreational to elite levels, only a select few ergogenic aids (such as creatine, sodium bicarbonate, and caffeine) have been proven to enhance sports performance (117).

Dietary supplements play a crucial role in building muscle, boosting the immune system, and providing fuel to enhance training or athletic performance. Elite athletes often utilize performance-enhancing agents, with many considering supplements to be an essential component for sports success (118). A Canadian study revealed that 87% of elite male and female athletes used supplements regularly (112), while another study conducted in Canada found that 98% of young athletes between the ages of 11 and 25 used supplements on a regular or intermittent basis (119). In this section, we will discuss

the effectiveness of the most widely used supplements for improving physical performance.

3.7.1 Creatine

Creatine stands out as the most frequently used and scientifically backed ergogenic aid (120, 121). It holds a preferred status over other ergogenic aids due to its proven ability to increase power, enhance muscular strength, and promote an increase in fat-free mass, ultimately improving exercise and sports performance (121).

Recent studies highlight that creatine supplementation, with doses ranging from 0.3 g per kg per day for 3 to 5 days or 20 g per day for 5 to 7 days without interruption, results in a rapid increase in intramuscular creatine, providing ergogenic advantages (122). Additionally, creatine supplements have been shown to accelerate recovery from injury and muscle damage (122, 123). Notably, a study demonstrated that introducing a preload at 0.3 g/kg body weight and a post-maintenance protocol at 0.1 g/kg body weight after a vigorous eccentric resistance training session in young men led to reduced muscle damage and strength loss (124). Another study documented a substantial reduction in oxidative stress and an increase in athletic performance in male athletes who took creatine monohydrate (20 g/day) after 7 days (125). Previous research also suggests that creatine supplements can mitigate muscle damage resulting from prolonged, intense exercise sessions.

Studies on creatine supplements consistently show enhanced performance and increased strength in short-duration, maximal-intensity exercises, as evidenced by improvements in metrics such as single-repetition maximum, muscular strength, repetitions, muscular endurance, speed, and overall strength (126, 127). A meta-analysis examining the impact of creatine supplementation on upper and lower extremity performance revealed a noticeable increase in strength for both extremities (128). Notably, performance improvement was observed in individuals following a creatine supplementation program, particularly in conjunction with resistance training. This effect was particularly pronounced in individuals without a history of prior training, defined as those engaging in exercise less than 3 h per week.

Currently, the scientific literature strongly supports the utilization of creatine supplementation for boosting performance in short-duration, high-intensity resistance training, demonstrating a distinct influence on lean body mass. Nevertheless, it remains unclear whether these effects of creatine supplementation translate into enhanced athletic performance.

3.7.2 Caffeine

Caffeine, a natural derivative stimulant, is associated with several proposed ergogenic effects. Known for its stimulating properties, caffeine not only improves performance but also increases the release of neurotransmitters, enhances intellectual ability, and boosts energy expenditure (129). Studies indicate that caffeine serves as a potent ergogenic aid for both aerobic and anaerobic training, particularly benefiting endurance activities like cycling and running (71).

Research suggests that consuming 2–5 mg/kg of caffeine before engaging in performance-based activities can significantly enhance sports performance (130). Another study demonstrated an approximate 3.2% improvement in athletic performance with similar doses of caffeine (131). It has been observed that moderate doses of caffeine, ranging from 3 to 6 mg/kg, are effective in improving sports

performance, while high doses (≥ 9 mg/kg) do not show any additional benefits (80).

Several proposed mechanisms aim to explain the impact of caffeine supplements on athletic performance, focusing on their effects on endurance, muscle contraction, and perceived exertion (132, 133). The primary mechanism involves caffeine's action within the central nervous system, where it competes with adenosine for its receptors, mitigating the adverse effects of adenosine on neurotransmission, arousal, and pain perception (134). Additionally, the analgesic effect of caffeine reduces the perception of pain and effort during exercise, potentially serving as an additional mechanism, especially in exercises inducing discomfort (132, 133). Consequently, reduced pain perception may contribute to sustained or increased motor unit firing rates, facilitating greater force production (133, 135).

The recommended dosage of 2–5 mg/kg aligns with existing literature, emphasizing the importance of optimal caffeine intake for maximizing benefits. The observation that high doses do not yield additional advantages suggests a dose–response relationship, emphasizing the need for moderation in caffeine consumption.

3.7.3 Protein and amino acid supplements

Amino acid supplements enjoy widespread popularity and are commonly utilized by highly-trained athletes. Beyond the realm of sports, amino acids offer potential therapeutic benefits, such as promoting healing, enhancing the immune system, preventing muscle atrophy in both the elderly and malnourished individuals, and contributing to the treatment of kidney and liver diseases (136).

Critical for maintaining a positive nitrogen balance in the body, amino acid supplements, including branched-chain amino acids and protein powder, play an essential role (71). Approved by the FDA to counteract nitrogen loss, protein supplements are recognized as safe when used in accordance with good manufacturing or feeding practices (REF) (80).

3.7.4 Whey protein

Following resistance training, incorporating whey supplements may contribute to enhanced muscle building. In a study, participants who consumed a 20 g whey supplement before and after resistance exercise exhibited greater gains in muscle mass and strength over a 10-week period compared to those who took a placebo (137). Another investigation revealed that athletes who consumed a whey supplement before and after a training session achieved the ability to perform more repetitions and lift heavier weights 24 and 48 h after the exercise, respectively (138).

It is crucial to emphasize that immediately after resistance training, the consumption of a high-quality protein source promotes muscle growth and aids in recovery (139). While whey supplements may be preferred over casein or soy in the immediate post-exercise period due to their faster absorption, there is no evidence suggesting that they result in greater muscle growth over a 24-h period (140). Additionally, whey protein has been associated with potential immune system benefits. Participants taking whey supplements experienced a smaller drop in glutathione levels, linked to lower immunity, after a 40-kilometer cycling time trial (141).

After intense exercise, additional protein is necessary to build new muscle proteins and repair damaged muscle cells (137). Current recommendations from scientists suggest athletes should consume between 1.3 and 1.8 g of protein per kilogram of body weight per day.

The precise amount of protein required for muscle building has been a subject of debate, with strength and power athletes tending to consume at the higher end of this range (1.6–1.8 g) (103).

The study's results support the idea that the timing and type of protein intake can influence post-exercise outcomes. While the faster absorption of whey is advantageous immediately after training, the overall 24-h impact on muscle growth appears comparable to other high-quality protein sources. Athletes can strategically incorporate whey protein into their post-exercise nutrition, recognizing its advantages while ensuring a balanced overall protein intake.

3.7.5 Branched-chain amino acid

The best way to describe branched-chain amino acids (BCAAs) is as a combination of three out of the nine essential amino acids. Valine, leucine, and isoleucine, the three BCAAs, cannot be synthesized by the body on its own (142). These amino acids collectively constitute one-third of muscle proteins and play a pivotal role in the metabolism of skeletal muscle due to their distinctive properties (143). BCAAs facilitate the absorption of blood sugar by muscle fibers and influence insulin signaling (143). Notably, leucine is of particular importance among the three BCAAs, serving a crucial role in regulating muscle protein synthesis (MPS) and acting as a modulator even in the presence of hyperaminoacidemia (144).

Additionally, BCAA supplements operate through various mechanisms, including reducing soreness and preventing muscle tissue breakdown during resistance and intense training (145). They contribute to the reduction of central fatigue, promote muscle function recovery, and maximize the MPS response (145).

According to some studies, incorporating BCAAs before and after exercise may effectively prevent exercise-induced muscle damage and increase muscle protein synthesis (146). There is evidence suggesting that taking BCAA supplements before resistance training can also reduce delayed-onset muscle soreness and assist athletes in maintaining muscle mass during dieting (147).

However, it seems that endurance athletes may not significantly benefit from BCAA supplementation. A study conducted at Florida State University indicated that while taking a BCAA supplement before and during prolonged endurance exercise reduced muscle damage, similar effects were achieved by consuming a sports drink with carbohydrates (146). Another study involving long-distance runners found that, compared to a placebo, BCAA supplements taken 7 days prior to a marathon did not improve performance or reduce muscle damage (148). In essence, BCAAs do not appear to offer significant performance advantages during endurance exercises.

BCAAs, with a particular emphasis on leucine, play a crucial role in muscle protein synthesis and various aspects of muscle metabolism. The documented advantages of BCAA supplementation, such as reducing soreness, preventing muscle tissue breakdown, and enhancing recovery, align with their well-established role in supporting muscle function. While BCAAs demonstrate potential benefits in situations like resistance training and muscle preservation during dieting, their advantages may not be notably pronounced in the context of endurance exercises.

3.7.6 Arginine

L-arginine, a non-essential amino acid naturally produced in the body, is commonly known by names such as arginine alpha-ketoglutarate (A-AKG) and arginine ketoisocaproate (A-KIC) (80).

Numerous studies suggest that the performance of elite athletes during anaerobic exercise remains largely unaffected by arginine supplements (149). In a study focused on A-AKG supplements, athletes did not exhibit differences in nitric oxide (NO) levels, blood flow, or performance (150). However, a review of multiple studies indicated that arginine supplements might offer a modest benefit to novice athletes but not to more experienced athletes or female athletes (151). The recommended safe dose for arginine is up to 20 g per day (80).

While arginine is a naturally occurring amino acid, its supplementation seems to have a limited impact on elite athletes during anaerobic exercise, as suggested by several studies. The potential modest benefit for novice athletes, highlighted in a review, prompts further investigation into factors such as experience level and gender that may influence the effectiveness of arginine supplementation. The specified safe dose serves as a reference for individuals considering incorporating arginine into their nutritional regimen.

3.7.7 Beta-alanine

Beta-alanine, a non-essential amino acid naturally produced in the body, increases muscle carnosine concentrations when taken as a supplement (152). Elevated muscle carnosine levels enhance buffering capacity, reducing lactic acid buildup during high-intensity exercise, which can improve performance in sprints and short distances by mitigating fatigue (153).

A systematic review of 19 randomized controlled studies has confirmed that beta-alanine supplements enhance performance in short, high-intensity activities (154). Analyzing 15 studies revealed an average performance improvement of 2.85%, translating to a 6-s enhancement over a 4-min event (155). Notably, runners who took beta-alanine supplements for 28 days in an Australian study significantly improved their 800-meter race times (156). In a United Kingdom study, amateur boxers receiving 6 g of beta-alanine per day for 4 weeks experienced a 20-fold increase in punch force and a four-fold increase in punch frequency (80). Many studies utilize daily doses of 3.2–6.4 g for six to 10 weeks (157). However, optimal results seem achievable with approximately 3 g (4 × 800 mg) per day for 6 weeks, followed by a maintenance dose of 1.2 g per day (156).

Beta-alanine supplementation has demonstrated efficacy in enhancing performance in short, high-intensity activities through increased muscle carnosine levels. The systematic review and specific studies provide robust evidence of its positive impact on various athletic parameters. The recommended dosage strategy underscores the significance of both the initial loading phase and the subsequent maintenance dose for optimal results. Athletes and individuals involved in high-intensity activities may consider beta-alanine supplementation as part of their performance enhancement strategy (158).

In summary, sports supplements lack systematic regulation, and there is no guarantee that they fulfill their claims or do not contain prohibited substances. Major sports organizations, including United Kingdom Sport, the US National Collegiate Athletic Association, and the International Olympic Committee (IOC), have policies advising against the use of sports supplements (159). It is recommended to prioritize a healthy diet and consult with your medical team or sports nutritionist before considering any supplements (160). Further research is essential to comprehend the combined effects of various sports supplement intake.

3.8 Nutrient timing

Nutrient timing involves strategically providing the appropriate macronutrients when the body is most primed to utilize them effectively (161). In the context of exercise, nutrient timing can be segmented into three distinct phases: the energy phase, the anabolic phase, and the adaptation phase.

The energy phase encompasses the period right before and during the exercise itself. Following exercise, the anabolic phase occurs immediately and typically spans about 60 to 90 min. This period, often referred to as the anabolic or metabolic window (161), highlights the heightened responsiveness of exercised muscles to nutrient intervention.

Subsequent to the anabolic phase, the adaptation phase unfolds. Consistently incorporating suitable supplements and meals during this period sustains an improved response to nutrient intervention for an extended duration. This fosters quicker recovery and facilitates training adaptation, enhancing overall exercise performance.

3.8.1 Pre-exercise nutrition

The recommended timeframe for pre-exercise nutrition typically extends to the hour leading up to a training session, although some research has investigated the impact of consuming nutrients up to 4 h before engaging in physical activity (162). During this period, the primary objective of nutrient consumption is to ensure an adequate fuel reserve for the muscles, thereby enhancing performance during the exercise.

Explorations into pre-exercise nutrition trace back to the 1930s, when researchers began investigating physiological reactions during exercise in response to the intake of pre-exercise carbohydrates (CHO), such as glucose and fructose (163). As research progressed, studies delved into manipulating exercise performance through pre-exercise nutrition strategies.

An early study involving trained swimmers, employing different nutritional strategies, including supplemental cane sugar, did not reveal significant differences in performance (164). However, this study laid the groundwork for subsequent interventions and explorations in the field. Hargreaves et al. (165) found that consuming 75 g of CHO, either as glucose or fructose, 45 min before exercise did not confer advantages in time-to-exhaustion performance or glycogen utilization during 75% effort of maximal oxygen uptake in trained cyclists compared to a placebo. Conversely, a larger carbohydrate intake of 312 g, consumed 4 h prior to exercise, enhanced time trial performance at a 70% effort level VO_{2max} after 100 min in individuals with recreational cycling training. However, these differences were not statistically significant when compared to meals with equivalent energy content comprising either 45 or 156 grams of carbohydrates (166).

In a separate study, significant improvements in a similar performance task were observed when recreationally trained individuals ingested either 1.1 or 2.2 g/kg of carbohydrates 1 h before exercise, in contrast to a placebo. Interestingly, no significant distinctions were noted between the two CHO doses (167) (33). Additionally, pre-exercise carbohydrate consumption has demonstrated an impact on substrate utilization during exercise, promoting glucose oxidation over free fatty acids even at low-to-moderate intensities (below 60% of maximal oxygen uptake, VO_{2max}) (168) (34).

In summary, the collective influence of pre-exercise carbohydrate intake on endurance performance generally appears favorable, although findings across studies can be inconsistent. Interpretation of results should consider methodological aspects, including factors like the time elapsed since the last intense training session and existing muscle glycogen levels, which are interconnected and can influence the effectiveness of pre-exercise feeding.

The significance and performance-enhancing benefits of pre-exercise carbohydrates may be contingent on muscle glycogen content before feeding. This suggests that individuals with limited rest between training sessions may derive greater benefits compared to those with extended rest periods, provided they adequately consume carbohydrates. While much of the research has focused on aerobic exercises, there is growing evidence that activities involving high-intensity intervals, such as resistance exercise, may also experience advantages. These activities predominantly rely on glycolytic, fast-twitch muscle fibers, which generate force through rapid muscular contractions fueled by stored phosphagens and anaerobic glycolysis, leading to lactate production.

Even with around 40% total glycogen depletion after high-volume resistance exercise, it suggests that carbohydrate availability may not be a limiting factor unless glycogen stores are suboptimal. Importantly, substantial evidence indicates that pre-exercise supplementation with carbohydrates can mitigate glycogen reductions, even if it does not notably impact blood glucose levels (169).

3.8.2 During exercise nutrition

Carbohydrate (CHO) intake during physical activity has been extensively studied since the 1960s (49). It plays a crucial role in mitigating the utilization of muscle and liver glycogen, especially in situations involving high exercise intensity, durations exceeding 60 min, or shorter, supramaximal exertions (170, 171). Insufficient CHO in these scenarios can lead to decreased exercise intensity due to a shortage of efficient fuel, diminished calcium release, and increased fatigue (172, 173). Inadequate carbohydrate intake during such activities may result in decreased exercise intensity due to a shortage of efficient fuel, diminished calcium release, and increased fatigue. Carbohydrate (CHO) oxidation rates from exogenous sources show an exponential increase in the initial 75–90 min of exercise, underlining the significance of commencing CHO ingestion from the start and sustaining it throughout the exercise session for the conservation of muscle and liver glycogen (174). Excessive CHO intake, on the other hand, may lead to gastrointestinal upset, potentially impeding performance goals. Diversifying CHO ingestion with different transporters can enhance CHO uptake and oxidation to approximately 1.5 g/min or 90 g/h (175). This varied CHO consumption not only improves CHO availability without causing gastrointestinal upset (176) but also carries the potential to enhance overall performance (177). For example, a study illustrated an 8% increase in power output during a time trial after 120 min of steady-state cycling when cyclists consumed beverages with a 2:1 ratio of glucose to fructose, as opposed to glucose alone (177). Importantly, fructose ingested at a rate of 1.2 g/min, when combined with an equal amount of glucose, leads to a higher CHO oxidation rate compared to lower quantities, in line with the commonly recommended 60 g/h. These results highlight the potential benefits of integrating varied carbohydrate (CHO) intake to enhance performance (171–174, 176, 177).

An alternative approach to optimizing carbohydrate (CHO) delivery, with the goal of minimizing gastrointestinal distress and potentially boosting performance, involves the simultaneous intake of protein and CHO. Recent findings from a review and meta-analysis indicated positive performance outcomes, especially in time trials or efforts to exhaustion, for groups consuming a combination of CHO and protein compared to CHO alone (178). The noted favorable effect persisted consistently, even with the utilization of non-isocaloric supplements. However, when ensuring that CHO and protein supplements were equivalent in CHO content and subsequent examination of the effects of isocaloric supplementation involving both CHO and protein or CHO alone on time to exhaustion, no notable differences were observed (178).

Although the simultaneous ingestion of protein and carbohydrates may not yield immediate performance improvements, there are indirect advantages. These encompass the capacity to boost caloric intake while reducing carbohydrate consumption to prevent gastrointestinal distress, enhancing amino acid bioavailability to reduce muscle protein breakdown, and improving amino acid availability for gluconeogenesis. Moreover, co-ingestion may play a role in postponing central nervous system fatigue (179).

The effectiveness of intra-exercise nutrition, especially the consumption of carbohydrates (CHO), is highly contingent on variables like pre-exercise feeding, glycogen status, and the type of exercise (178). For aerobic activities lasting more than 2 h, optimal results are achieved by consuming 90–144 g/h of CHO in a 2:1 glucose to fructose solution. This strategy maximizes the uptake and oxidation of CHO while simultaneously preserving muscle glycogen.

In competitive scenarios, where extended endurance events frequently conclude with a sprint to the finish line, relying significantly on anaerobic metabolism and the utilization of endogenous muscle glycogen, the prudent conservation of this fuel source throughout the entire bout becomes paramount.

3.8.3 Post-exercise nutrition

After engaging in physical activity, individuals commonly experience a temporary surge. During this phase, there is an increase in fatigue, muscle soreness, and a decline in performance. In this stage, catabolic processes take precedence, leading to decreased insulin levels, restricted glycogen, and limited substrate availability. Cortisol and catecholamines collectively influence physiological processes in the body, heightening the pace at which muscle protein is being broken down (180).

The intake of carbohydrates and protein post-exercise offers the potential to raise glucose levels in the bloodstream, reduce cortisol levels, and improve substrate availability, enabling the transition from a catabolic state to a more anabolic condition (180). Additionally, activating muscle GLUT4 transporters, increasing glycogen synthase activity, and enhancing insulin sensitivity all contribute to improving how responsive skeletal muscles are to absorbing carbohydrates and amino acids (50, 181). Therefore, the post-exercise period offers a strategic opportunity for nutrient intake to aid in replenishing muscle glycogen, promoting protein synthesis, and reducing the degradation of muscle proteins (180, 181). Integrating the timing of nutrient intake after exercising into a training routine becomes essential for optimizing recovery rates and maximizing the benefits of training.

During moderate-to-high intensity exercise, muscle glycogen assumes a crucial role as the primary source of energy to sustain

physical activity. In light of this situation, precise post-exercise nutrient timing becomes vital, emphasizing the primary goal of replenishing muscle glycogen to hasten the recovery process. After physical activity, there is a decrease in the heightened levels of post-exercise glucose transporters, which are crucial for the absorption of nutrients. This decline brings the transporter levels back to baseline within a two-hour period (182).

Aside from glycogen synthesis, the consumption of protein and essential amino acids following exercise plays a pivotal role in triggering muscle protein synthesis and aiding in the reconditioning of skeletal muscles (183). After exercising, there is a notable increase in muscle damage and protein degradation in the aftermath of exercise (161, 181). Moreover, when glycogen stores are depleted, the pace of protein breakdown increases, as amino acids could potentially undergo gluconeogenesis to be utilized in replenishing levels of glycogen (91). As a result, it is crucial to consume protein after exercise to mitigate the breakdown of proteins and assist in the repair of muscle damage (183).

When aiming to stimulate muscle protein synthesis, proteins that are rapidly digestible and of high quality, containing an adequate amount of essential amino acids, may be more effective than proteins with lower quantities of branched-chain amino acids or those that are slower to digest (69).

Comprehensive training and maintaining a sufficient daily protein intake are crucial for achieving strength and hypertrophy. However, beyond these foundational aspects, there are potential advantages to carefully considering the timing of protein consumption, especially immediately after exercising. The positive impacts on net protein balance and glycogen synthesis underscore the significant benefits of ingesting protein in the post-training period.

Fundamentally, critical factors contributing to optimal performance include not only the quality of the training but also the overall protein intake throughout the day. The strategic timing of protein consumption provides an additional layer of support to boost performance. Even if the resulting benefits are seemingly minor, this aspect becomes a pertinent factor, particularly for competitive athletes who are dedicated to optimizing their performance.

In summary, the significance of nutrient timing is a nuanced matter, and its relevance varies greatly depending on the context. Defined as the delivery of adequate macronutrients precisely when the body is ready to use them (184), nutrient timing represents a dietary approach where specific nutrients are ingested before training to enhance both short-term performance and long-term adaptations (23).

Early research delved into the effects of acute carbohydrate (CHO) consumption on exercise performance, focusing on glycogen depletion and use during moderate to high-intensity aerobic activity (185, 186). Subsequent studies broadened the scope to investigate how acute protein consumption (PRO) impacts endurance and resistance workout performance, as well as recovery and adaptation. Despite these inquiries, it's noteworthy that the effects of nutrient timing on performance, recovery, and adaptation outcome variables have only been explored in a limited number of chronic interventional studies.

The time before exercise, typically within 4 h prior, involves specific dietary tactics. Glucose loading, for instance, aims to boost muscle glycogen reserves in the days preceding an endurance race, resulting in enhanced glycogen storage levels and improved performance in activities lasting over 90 min (167, 187).

Consuming 150–200 grams of carbs 4 h before exercise has shown to significantly increase muscle glycogen reserves, ultimately contributing to improved exercise performance (188–190). Additionally, it is advisable to include healthy fats in the diet 6–5 h before exercise. Conversely, to prevent gastrointestinal discomfort, it's recommended to avoid fiber consumption 2–3 h before engaging in physical activity (67, 190, 191).

The energy phase during a workout is crucial as muscles require sufficient energy for contractions. According to timing science, consuming carbohydrates, proteins, specific amino acids, and vitamins 10 min before and during exercise is necessary for this objective. The benefits encompass a good supply of glycogen, a reduction in cortisol, and assistance in preparing muscle enzymes for faster recovery (192, 193).

Post-workout nutrition timing is widely regarded as the most crucial phase. Intensive resistance training sessions damage muscle fibers and deplete the body's reserves of fuel, including glycogen and amino acids. Consuming the right balance of nutrients during this time initiates the healing process for injured tissue and replenishes energy stores. This occurs in a super-compensated manner, enhancing exercise performance and body composition. Researchers often refer to this limited timeframe following training as the “anabolic window of opportunity,” emphasizing its significance in maximizing muscle adaptations related to the training (194–196).

4 Conclusion

In conclusion, this narrative review offers targeted recommendations for addressing the nutritional needs of the active population, with a specific focus on preventing disordered eating. Given the unique challenges faced by athletes, it is imperative to tailor nutrition plans to individual requirements.

Individualization emerges as a cornerstone in preventing disordered eating among athletes. Recognizing diverse goals, body compositions, metabolic rates, and dietary preferences is essential. Tailoring nutrition plans to accommodate these individual factors can significantly contribute to optimizing performance while mitigating the risk of disordered eating (197).

Macronutrients, which include carbohydrates, proteins, and fats, play a critical role in athletic nutrition. Adequate carbohydrate intake is necessary to support energy production and replenish glycogen stores, thereby reducing the likelihood of restrictive eating behaviors (33). Proteins are indispensable for muscle repair and growth, emphasizing the importance of meeting increased protein needs without resorting to excessive dietary restrictions (198). Meanwhile, healthy fats contribute to sustained energy, hormone production, and overall health, promoting a balanced approach to nutrition.

In addition to macronutrients, micronutrients, encompassing vitamins and minerals, are paramount for energy metabolism and immune function. Promoting a diverse, nutrient-dense diet is crucial to ensuring athletes receive adequate micronutrients, thereby reducing the risk of nutritional deficiencies that might contribute to disordered eating (199).

Hydration emerges as a key factor in preventing disordered eating among the active population. Proper fluid balance is essential for physiological function, and athletes must be attuned to their individual

fluid needs. Maintaining adequate hydration levels before, during, and after exercise is crucial, as dehydration can exacerbate disordered eating behaviors.

While acknowledging the interest in sports nutrition supplements, caution is advised. Athletes should prioritize meeting their nutritional needs through whole foods to minimize the risk of disordered eating patterns (200). Supplements should only be considered when dietary intake falls short or specific deficiencies are identified. Consultation with qualified professionals is essential to ensure safe and appropriate usage.

In summary, implementing these targeted nutritional recommendations can serve as a proactive tool in preventing disordered eating within the active population. By understanding and addressing the unique challenges faced by athletes, promoting individualization, and emphasizing a balanced and informed approach to nutrition, this review contributes to the overarching goal of investigating and preventing disordered eating in the active population.

This review delves into the most recent research findings on nutritional recommendations for athletes, offering readers a comprehensive overview of the current state of the field. By skillfully spotlighting significant patterns, accepted procedures, and novel studies, we provide an invaluable resource for both researchers and practitioners, enhancing the reader's comprehension of the intricate connection between nutrition and athletic performance.

It is important to note that the authors' judgment played a major role in the inclusion of studies. The absence of a systematic search and uniform inclusion criteria may lead to the inclusion of research with methodological flaws or the unintentional exclusion of pertinent studies. Additionally, subjectivity in the interpretation of results may have resulted in the overemphasis of some topics and the omission of others.

In summary, this review underscores the pivotal role of athlete nutrition guidelines in facilitating optimal dietary arrangements for individuals involved in sports and physical activity. By comprehensively reviewing existing guidelines, this manuscript aims to furnish a resource that benefits athletes directly and aids sports nutrition specialists in their vital work. The overarching objective is to cultivate an environment of informed dietary choices, contributing to the prevention of disordered eating and promoting the long-term health and performance of athletes and active individuals. As we navigate the intricacies of sports nutrition, the insights gleaned from this manuscript aspire to guide future research and interventions, ensuring a holistic approach to the well-being of individuals engaged in athletic pursuits.

Author contributions

AdA: Conceptualization, Writing – review & editing. WA: Writing – review & editing. MJ: Writing – review & editing, Data curation, Software. AmA: Methodology, Validation, Writing – original draft. SA: Conceptualization, Data curation, Writing – original draft. AH: Validation, Writing – review & editing. TB: Investigation, Writing – original draft, Writing – review & editing. BA: Supervision, Writing – review & editing. NT: Methodology, Writing – review & editing. HS: Formal analysis, Writing – original draft. HG: Conceptualization,

Investigation, Software, Supervision, Validation, Visualization, Writing – original draft, Writing – review & editing. AJ: _____.

Funding

The author(s) declare that no financial support was received for the research, authorship, and/or publication of this article.

Acknowledgments

The authors express gratitude to the members of the Nutritional Student Research Club at the University of Jordan for their contributions to this research.

References

- Amawi A, Alshuwaier G, Alaqil A, Alkassabeh WJ, Bursais A, Al-Nuaim A, et al. Exploring the impact of dietary factors on body composition in elite Saudi soccer players: a focus on added sugars, salt, and oil consumption. *Int J Hum Mov Sports Sci.* (2023) 11:1305–12. doi: 10.13189/saj.2023.110614
- Thomas DT, Erdman KA, Burke LM. Nutrition and athletic performance. *Med Sci Sports Exerc.* (2016) 48:543–68. doi: 10.1249/MSS.0000000000000852
- Kreider RB, Wilborn CD, Taylor L, Campbell B, Almada AL, Collins R, et al. ISSN exercise and sport nutrition review: research and recommendations. *J Int Soc Sports Nutr.* (2010) 7:7. doi: 10.1186/1550-2783-7-7
- Loucks AB, Kiens B, Wright HH. Energy availability in athletes. *J Sports Sci.* (2011) 29:S7–S15. doi: 10.1080/02640414.2011.588958
- McArdle WD. *Sports and exercise nutrition.* New York: Lippincott Williams and Wilkins (2018).
- Okta PG, Yildiz E. The validity and reliability study of the Turkish version of the general and sport nutrition knowledge questionnaire (GeSNK). *Prog Nutr.* (2021) 1:e2021027. doi: 10.23751/pn.v23i1.10029
- Vázquez-Espino K, Fernández-Tena C, Lizarraga-Dallo MA, Farran-Codina A. Development and validation of a short sport nutrition knowledge questionnaire for athletes. *Nutrients.* (2020) 12:3561. doi: 10.3390/nu12113561
- Bradley WJ, Cavanagh BP, Douglas W, Donovan TF, Morton JP, Close GL. Quantification of training load, energy intake, and physiological adaptations during a rugby preseason: a case study from an elite European rugby union squad. *J Strength Cond Res.* (2015) 29:534–44. doi: 10.1519/JSC.0000000000000631
- Alkassabeh WJ, Amawi AT. The effectiveness of using Mobile learning application on undergraduates' intrinsic motivation and their general nutrition knowledge. *Int J Interact Mob Technol.* (2023) 17:19–37. doi: 10.3991/ijim.v17i17.40959
- Birkenhead KL, Slater G. A review of factors influencing athletes' food choices. *Sports Med.* (2015) 45:1511–22. doi: 10.1007/s40279-015-0372-1
- O'Brien L, Collins K, Amirabdollahian F. Exploring sports nutrition knowledge in elite Gaelic footballers. *Nutrients.* (2021) 13:1081. doi: 10.3390/nu13041081
- Alkassabeh WJ, Amawi AT. Impact of eating habits on the psychological state of Jordanian athletes: a descriptive study. *Food Sci Technol.* (2023) 11:168–81. doi: 10.13189/fst.2023.110305
- Boumosleh JM, El Hage C, Farhat A. Sports nutrition knowledge and perceptions among professional basketball athletes and coaches in Lebanon—a cross-sectional study. *BMC Sports Sci Med Rehabil.* (2021) 13:1–19. doi: 10.1186/s13102-021-00280-6
- Trakman GL, Forsyth A, Devlin BL, Belski R. A systematic review of athletes' and coaches' nutrition knowledge and reflections on the quality of current nutrition knowledge measures. *Nutrients.* (2016) 8:570. doi: 10.3390/nu8090570
- Spendlove JK, Heaney SE, Gifford JA, Prvan T, Denyer GS, O'Connor HT. Evaluation of general nutrition knowledge in elite Australian athletes. *Med Sci Sports Exerc.* (2011) 43:501. doi: 10.1249/01.MSS.0000401383.38771.7e
- Tam R, Beck KL, Manore MM, Gifford J, Flood VM, O'Connor H. Effectiveness of education interventions designed to improve nutrition knowledge in athletes: a systematic review. *Sports Med.* (2019) 49:1769–86. doi: 10.1007/s40279-019-01157-y
- Elias SSM, Saad HA, Taib MNM, Jamil Z. Effects of sports nutrition education intervention on sports nutrition knowledge, attitude and practice, and dietary intake of Malaysian team sports athletes. *Malays J Nutr.* (2018) 2018:24.
- Nascimento M, Silva D, Ribeiro S, Nunes M, Almeida M, Mendes-Netto R. Effect of a nutritional intervention in athlete's body composition, eating behaviour and

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

nutritional knowledge: a comparison between adults and adolescents. *Nutrients.* (2016) 8:535. doi: 10.3390/nu8090535

19. Patton-Lopez MM, Manore MM, Branscum A, Meng Y, Wong SS. Changes in sport nutrition knowledge, attitudes/beliefs and behaviors following a two-year sport nutrition education and life-skills intervention among high school soccer players. *Nutrients.* (2018) 10:1636. doi: 10.3390/nu10111636

20. Kerkick CM, Wilborn CD, Roberts MD, Smith-Ryan A, Kleiner SM, Jäger R, et al. ISSN exercise and sports nutrition review update: research and recommendations. *J Int Soc Sports Nutr.* (2018) 15:38. doi: 10.1186/s12970-018-0242-y

21. Holway FE, Spriet LL. Sport-specific nutrition: practical strategies for team sports. *J Sports Sci.* (2011) 29:S115–25. doi: 10.1080/02640414.2011.605459

22. Burke LM, Hawley JA, Wong SHS, Jeukendrup AE. Carbohydrates for training and competition. *J Sports Sci.* (2011) 29:S17–27. doi: 10.1080/02640414.2011.585473

23. Kerkick CM, Arent S, Schoenfeld BJ, Stout JR, Campbell B, Wilborn CD, et al. International Society of Sports Nutrition position stand: nutrient timing. *J Int Soc Sports Nutr.* (2017) 14:33. doi: 10.1186/s12970-017-0189-4

24. Aragon AA, Schoenfeld BJ, Wildman R, Kleiner S, VanDusseldorp T, Taylor L, et al. International society of sports nutrition position stand: diets and body composition. *J Int Soc Sports Nutr.* (2017) 14:16. doi: 10.1186/s12970-017-0174-y

25. Jäger R, Kerkick CM, Campbell BI, Cribb PJ, Wells SD, Skwiat TM, et al. International society of sports nutrition position stand: protein and exercise. *J Int Soc Sports Nutr.* (2017) 14:20. doi: 10.1186/s12970-017-0177-8

26. Sánchez-Díaz S, Yanci J, Raya-González J, Scanlan AT, Castillo D. A comparison in physical fitness attributes, physical activity behaviors, nutritional habits, and nutritional knowledge between elite male and female youth basketball players. *Front Psychol.* (2021) 12:685203. doi: 10.3389/fpsyg.2021.685203

27. Ali A, Al-Siyabi MS, Waly MI, Kilani NI. Assessment of nutritional knowledge, dietary habits and nutrient intake of university student athletes. *Pak J Nutr.* (2015) 14:293. doi: 10.3923/pjn.2015.293.299

28. Sunuwar DR, Singh DR, Bohora MP, Shrestha V, Karki K, Pradhan PMS. Association of nutrition knowledge, practice, supplement use and nutrient intake with athletic performance among Taekwondo players in Nepal. *Front Nutr.* (2022) 9:1004288. doi: 10.3389/fnut.2022.1004288

29. Papadopoulou SK. Rehabilitation nutrition for injury recovery of athletes: the role of macronutrient intake. *Nutrients.* (2020) 12:2449. doi: 10.3390/nu12082449

30. Wasserfurth P, Palmowski J, Hahn A, Krüger K. Reasons for and consequences of low energy availability in female and male athletes: social environment, adaptations, and prevention. *Sports Med Open.* (2020) 6:44. doi: 10.1186/s40798-020-00275-6

31. Logue DM, Madigan SM, Melin A, Delahunty E, Heinen M, Donnell SJM, et al. Low energy availability in athletes 2020: an updated narrative review of prevalence, risk, within-day energy balance, knowledge, and impact on sports performance. *Nutrients.* (2020) 12:835. doi: 10.3390/nu12030835

32. Jurov I, Keay N, Hadžić V, Spudić D, Rauter S. Relationship between energy availability, energy conservation and cognitive restraint with performance measures in male endurance athletes. *J Int Soc Sports Nutr.* (2021) 18:1–10. doi: 10.1186/s12970-021-00419-3

33. Henselmans M, Bjørnsen T, Hedderman R, Vårvik FT. The effect of carbohydrate intake on strength and resistance training performance: a systematic review. *Nutrients.* (2022) 14:856. doi: 10.3390/nu14040856

34. Podlogar T, Wallis GA. New horizons in carbohydrate research and application for endurance athletes. *Sports Med.* (2022) 52:5–23. doi: 10.1007/s40279-022-01757-1

35. König D, Braun H, Carlsohn A, Großhauser M, Lampen A, Mosler S, et al. Position of the working group sports nutrition of the German Nutrition Society (DGE): carbohydrates in sports nutrition. *Dtsch Z Sportmed.* (2020) 71:185–91. doi: 10.5960/dzsm.2020.455
36. Hargreaves M, Spriet LL. Skeletal muscle energy metabolism during exercise. *Nat Metab.* (2020) 2:817–28. doi: 10.1038/s42255-020-0251-4
37. Richter EA, Sylow L, Hargreaves M. Interactions between insulin and exercise. *Biochem J.* (2021) 478:3827–46. doi: 10.1042/BCJ20210185
38. Tarnopolsky MA, Gibala M, Jeukendrup AE, Phillips SM. Nutritional needs of elite endurance athletes. Part I: carbohydrate and fluid requirements. *Eur J Sport Sci.* (2005) 5:3–14. doi: 10.1080/17461390500076741
39. Burk L, Deakin V. *Clinical sport nutrition*. 4th ed. New York: McGraw Hill (2012).
40. Ghazzawi HA, Hussain MA, Raziq KM, Alsendi KK, Alaamer RO, Jaradat M, et al. Exploring the relationship between micronutrients and athletic performance: A comprehensive scientific systematic review of the literature in sports medicine. *Sports.* (2023) 11:109. doi: 10.3390/sports11060109
41. Howarth KR, Moreau NA, Phillips SM, Gibala MJ. Coingestion of protein with carbohydrate during recovery from endurance exercise stimulates skeletal muscle protein synthesis in humans. *J Appl Physiol.* (2009). doi: 10.1152/jappphysiol.90333.2008
42. Burke LM. Guidelines for daily carbohydrate intake: do athletes achieve them? *Sports Med.* (2001) 31:267–99. doi: 10.2165/00007256-200131040-00003
43. Van Loon LJC, Saris WHM, Kruijshoop M, Wagenmakers AJM. Maximizing postexercise muscle glycogen synthesis: carbohydrate supplementation and the application of amino acid or protein hydrolysate mixtures. *Am J Clin Nutr.* (2000) 72:106–11. doi: 10.1093/ajcn/72.1.106
44. Ryan M. *Sports nutrition for endurance athletes*. United States: VeloPress (2012).
45. Romano-Ely BC, Todd MK, Saunders MJ, Laurent TST. Effect of an isocaloric carbohydrate-protein-antioxidant drink on cycling performance. *Med Sci Sports Exerc.* (2006) 38:1608–16. doi: 10.1249/01.mss.0000229458.11452.e9
46. Van Hall G, Shirreffs SM, Calbet J. Muscle glycogen resynthesis during recovery from cycle exercise: no effect of additional protein ingestion. *J Appl Physiol.* (2000) 88:1631–6. doi: 10.1152/jappphysiol.2000.88.5.1631
47. Saunders MJ, Luden ND, Herrick JE. Consumption of an oral carbohydrate-protein gel improves cycling endurance and prevents postexercise muscle damage. *J Strength Cond Res.* (2007) 21:678–84. doi: 10.1519/R-20506.1
48. Bishop PA, Jones E, Woods AK. Recovery from training: a brief review: brief review. *J Strength Cond Res.* (2008) 22:1015–24. doi: 10.1519/JSC.0b013e31816eb518
49. Saunders MJ, Kane MD, Todd MK. Effects of a carbohydrate-protein beverage on cycling endurance and muscle damage. *Med Sci Sports Exerc.* (2004) 36:1233–8. doi: 10.1249/01.MSS.0000132377.66177.9F
50. Jentjens R, Jeukendrup AE. Determinants of post-exercise glycogen synthesis during short-term recovery. *Sports Med.* (2003) 33:117–44. doi: 10.2165/00007256-200333020-00004
51. Ivy JL, Goforth HW, Damon BM, McCauley TR, Parsons EC, Price TB. Early postexercise muscle glycogen recovery is enhanced with a carbohydrate-protein supplement. *J Appl Physiol.* (2002) 93:1337–44. doi: 10.1152/jappphysiol.00394.2002
52. Van Hall G, Saris WHM, Wagenmakers AJM. Effect of carbohydrate supplementation on plasma glutamine during prolonged exercise and recovery. *Int J Sports Med.* (1998) 19:82–6. doi: 10.1055/s-2007-971886
53. Reinhard C, Galloway SDR. Carbohydrate intake practices and determinants of food choices during training in recreational, amateur, and professional endurance athletes: a survey analysis. *Front Nutr.* (2022) 9:862396. doi: 10.3389/fnut.2022.862396
54. Burd NA, Tang JE, Moore DR, Phillips SM. Exercise training and protein metabolism: influences of contraction, protein intake, and sex-based differences. *J Appl Physiol.* (2009) 106:1692–701. doi: 10.1152/jappphysiol.91351.2008
55. Koopman R, van Loon LJC. Aging, exercise, and muscle protein metabolism. *J Appl Physiol.* (2009) 106:2040–8. doi: 10.1152/jappphysiol.91551.2008
56. Biolo G, Tipton KD, Klein S, Wolfe RR. An abundant supply of amino acids enhances the metabolic effect of exercise on muscle protein. *Am J Physiol.* (1997) 273:E122–9. doi: 10.1152/ajpendo.1997.273.1.E122
57. Yang Y, Churchward-Venne TA, Burd NA, Breen L, Tarnopolsky MA, Phillips SM. Myofibrillar protein synthesis following ingestion of soy protein isolate at rest and after resistance exercise in elderly men. In: Y Yang. *Clinical nutrition and aging*. New Jersey: Apple Academic Press, 105–126. (2017).
58. Moore DR, Soeters PB. The biological value of protein In: RF Meier, BR Reddy and PB Soeters, editors. *The importance of nutrition as an integral part of disease management*. Basel, Switzerland: Karger Publishers (2015). 39–51.
59. Szedlak C, Robins A. Protein requirements for strength training. *Strength Cond J.* (2012) 34:85–91. doi: 10.1519/SSC.0b013e31826dc3c4
60. Berrazaga I, Micard V, Gueugneau M, Walrand S. The role of the anabolic properties of plant-versus animal-based protein sources in supporting muscle mass maintenance: a critical review. *Nutrients.* (2019) 11:1825. doi: 10.3390/nu11081825
61. Schaafsma G. The protein digestibility-corrected amino acid score. *J Nutr.* (2000) 130:1865S–7S. doi: 10.1093/jn/130.7.1865S
62. Gorissen SHM, Crombag JJR, Senden JMG, Waterval WAH, Bierau J, Verdijk LB, et al. Protein content and amino acid composition of commercially available plant-based protein isolates. *Amino Acids.* (2018) 50:1685–95. doi: 10.1007/s00726-018-2640-5
63. Consultation FE. Dietary protein quality evaluation in human nutrition. *FAO Food Nutr Pap.* (2011) 92:1–66.
64. Millward DJ, Fereday A, Gibson NR, Cox MC, Pacy PJ. Efficiency of utilization of wheat and milk protein in healthy adults and apparent lysine requirements determined by a single-meal [1-13 C] leucine balance protocol. *Am J Clin Nutr.* (2002) 76:1326–34. doi: 10.1093/ajcn/76.6.1326
65. Fouillet H, Bos C, Gaudichon C, Tomé D. Approaches to quantifying protein metabolism in response to nutrient ingestion. *J Nutr.* (2002) 132:3208S–18S. doi: 10.1093/jn/131.10.3208S
66. Ivy JL. Regulation of muscle glycogen repletion, muscle protein synthesis and repair following exercise. *J Sports Sci Med.* (2004) 3:131–8.
67. Wirth J, Hillesheim E, Brennan L. The role of protein intake and its timing on body composition and muscle function in healthy adults: a systematic review and Meta-analysis of randomized controlled trials. *J Nutr.* (2020) 150:1443–60. doi: 10.1093/jn/nxaa049
68. Moore DR. Nutrition to support recovery from endurance exercise: Optimal carbohydrate and protein replacement. *Curr Sports Med Rep.* (2015) 14:294–300. doi: 10.1249/JSR.0000000000000180
69. Moore DR. Maximizing post-exercise anabolism: The case for relative protein intakes. *Front Nutr.* (2019) 6:147. doi: 10.3389/fnut.2019.00147
70. Morton RW, Murphy KT, McKellar SR, Schoenfeld BJ, Henselmans M, Helms E, et al. A systematic review, meta-analysis and meta-regression of the effect of protein supplementation on resistance training-induced gains in muscle mass and strength in healthy adults. *Br J Sports Med.* (2018) 52:376–84. doi: 10.1136/bjsports-2017-097608
71. Vitale K, Getzin A. Nutrition and supplement update for the endurance athlete: Review and recommendations. *Nutrients.* (2019) 11:1289. doi: 10.3390/nu11061289
72. Casazza GA, Tovar AP, Richardson CE, Cortez AN, Davis BA. Energy availability, macronutrient intake, and nutritional supplementation for improving exercise performance in endurance athletes. *Curr Sports Med Rep.* (2018) 17:215–23. doi: 10.1249/JSR.0000000000000494
73. Stokes T, Hector AJ, Morton RW, McGlory C, Phillips SM. Recent perspectives regarding the role of dietary protein for the promotion of muscle hypertrophy with resistance exercise training. *Nutrients.* (2018) 10:180. doi: 10.3390/nu10020180
74. LaPelusa A, Kaushik R. *Physiology, proteins*. Treasure Island, FL: StatPearls Publishing (2023).
75. Callahan A, Leonard H, Powell T. *Nutrition: Science and everyday application*. 2nd ed. New York: Springer (2022).
76. Kerksick CM, Kulovitz M. Requirements of energy, carbohydrates, proteins and fats for athletes In: CM Kerksick, editor. *Nutrition and enhanced sports performance: Muscle building, endurance, and strength*. Amsterdam, Netherlands: Elsevier Inc. (2014). 355–66.
77. Ghazzawi HA, Alshuwaier GO, Alaql AI, Bursais AK, Al-Nuaim AA, Alhaji JH, et al. Correlation of consumption vegetables, fruit, and nuts with body mass index and fat deposition in Saudi elite male soccer players. *Int J Hum Mov Sports Sci.* (2023) 11:350–9. doi: 10.13189/saj.2023.110212
78. D'Angelo S, Madonna G, Di Palma D. Effects of fish oil supplementation in the sport performance. *J Phys Educ Sport.* (2020) 20:2322–9. doi: 10.7752/jpes.2020.s4313
79. Jeromson S, Gallagher IJ, Galloway SDR, Hamilton DL. Omega-3 fatty acids and skeletal muscle health. *Mar Drugs.* (2015) 13:6977–7004. doi: 10.3390/md13116977
80. Bean A. *Sport supplements: Which nutritional supplements really work*. 2nd ed. Karnataka: Bloomsbury (2015).
81. Genton L, Melzer K, Pichard C. Energy and macronutrient requirements for physical fitness in exercising subjects. *Clin Nutr.* (2010) 29:413–23. doi: 10.1016/j.clnu.2010.02.002
82. Burke LM, Angus DJ, Cox GR, Cummings NK, Febbraio MA, Gawthorn K, et al. Effect of fat adaptation and carbohydrate restoration on metabolism and performance during prolonged cycling. *J Appl Physiol.* (2000) 89:2413–21. doi: 10.1152/jappphysiol.2000.89.6.2413
83. Stellingwerff T, Spriet LL, Watt MJ, Kimber NE, Hargreaves M, Hawley JA, et al. Decreased PDH activation and glycogenolysis during exercise following fat adaptation with carbohydrate restoration. *Am J Physiol.* (2006) 290:E380–8. doi: 10.1152/ajpendo.00268.2005
84. Burke LM, Kiens B. "Fat adaptation" for athletic performance: the nail in the coffin? *J Appl Physiol.* (2006) 100:7–8. doi: 10.1152/jappphysiol.01238.2005
85. Fogelholm M, Burke L, Deakin V. Micronutrients: vitamins, minerals and antioxidants. In: L Burke and V Deakin. *Clinical Sports Nutrition, 5th Edn*. New York: McGraw Hill, pp. 310–345. (2015).
86. Knapik JJ, Steelman RA, Hoedebecke SS, Austin KG, Farina EK, Lieberman HR. Prevalence of dietary supplement use by athletes: systematic review and meta-analysis. *Sports Med.* (2016) 46:103–23. doi: 10.1007/s40279-015-0387-7

87. Volpe SL. Micronutrient requirements for athletes. *Clin Sports Med.* (2007) 26:119–30. doi: 10.1016/j.csm.2006.11.009
88. Deakin V, Peeling P. Prevention, detection and treatment of iron depletion and deficiency in athletes. In: L Burke and V Deakin, editors. *Clinical sports nutrition*. New York: McGraw-Hill Education (2015). 266–309.
89. Garvican-Lewis LA, Govus AD, Peeling P, Abbiss CR, Gore CJ. Iron supplementation and altitude: decision making using a regression tree. *J Sports Sci Med.* (2016) 15:204.
90. Beck KL, Mitchell S, Foskett A, Conlon CA, Von Hurst PR. Dietary intake, anthropometric characteristics, and iron and vitamin D status of female adolescent ballet dancers living in New Zealand. *Int J Sport Nutr Exerc Metab.* (2015) 25:335–43. doi: 10.1123/ijsnem.2014-0089
91. Jordan SL, Albracht-Schulte K, Robert-McComb JJ. Micronutrient deficiency in athletes and inefficiency of supplementation: Is low energy availability a culprit? *PharmaNutrition.* (2020) 14:100229. doi: 10.1016/j.phanu.2020.100229
92. Ghazzawi HA, Amawi AT, Alduraidi H, Juweid M, Alhawari HH, Al-Abbadi MA, et al. The preventable effect of taekwondo sport among cadets and junior Bone Mineral Density: DEXA Assessment. *Children.* (2023) 10:170. doi: 10.3390/children10010170
93. Bytowski JR. Fueling for performance. *Sports Health.* (2018) 10:47–53. doi: 10.1177/1941738117743913
94. The team physician and the return-to-play decision. A consensus statement - 2012 update. *Med Sci Sports Exerc.* (2012) 44:2446–8. doi: 10.1249/MSS.0b013e3182750534
95. Barley OR, Chapman DW, Abbiss CR. Reviewing the current methods of assessing hydration in athletes. *J Int Soc Sports Nutr.* (2020) 17:52. doi: 10.1186/s12970-020-00381-6
96. Knechtle B, Chlíbková D, Papadopoulou S, Mantzorou M, Rosemann T, Nikolaidis PT. Exercise-associated hyponatremia in endurance and ultra-endurance performance—aspects of sex, race location, ambient temperature, sports discipline, and length of performance: A narrative review. *Medicina (Lithuania).* (2019) 55:537. doi: 10.3390/medicina55090537
97. Montain SJ, Chevront SN, Sawka MN. Exercise associated hyponatremia: quantitative analysis to understand the aetiology. *Br J Sports Med.* (2006) 40:98–105. doi: 10.1136/bjism.2005.018481
98. Périard JD, Eijssvogels TMH, Daanen HAM. Exercise under heat stress: thermoregulation, hydration, performance implications, and mitigation strategies. *Physiol Rev Am Physiol Soc.* (2021) 101:1873–979. doi: 10.1152/physrev.00038.2020
99. National Strength and Conditioning Association. *NSCA's guide to sport and exercise nutrition*. 2nd ed. United States: National Strength and Conditioning Association (2020).
100. Channel Protein. *Channel Protein.* (2014).
101. Maughan RJ, Shirreffs SM. Development of hydration strategies to optimize performance for athletes in high-intensity sports and in sports with repeated intense efforts. *Scand J Med Sci Sports.* (2010) 20:59–69. doi: 10.1111/j.1600-0838.2010.01191.x
102. Ayotte D, Corcoran MP. Individualized hydration plans improve performance outcomes for collegiate athletes engaging in in-season training. *J Int Soc Sports Nutr.* (2018) 15:27. doi: 10.1186/s12970-018-0230-2
103. Campbell C, Prince D, Braun M, Applegate E, Casazza GA. Carbohydrate-supplement form and exercise performance. *Int J Sport Nutr Exerc Metab.* (2008) 18:179–90. doi: 10.1123/ijsnem.18.2.179
104. Nutritional Assessment of Athletes. *Nutritional Assessment of Athletes.* (2011).
105. Pence J, Bloomer RJ. Impact of nuun electrolyte tablets on fluid balance in active men and women. *Nutrients.* (2020) 12:1–11. doi: 10.3390/nu12103030
106. Berry CW, Wolf ST, Murray B, Kenney WL. Hydration efficacy of a milk permeate-based oral hydration solution. *Nutrients.* (2020) 12:502. doi: 10.3390/nu12051502
107. Wolf ST, Stanhewicz AE, Clarke MM, Chevront SN, Kenefick RW, Kenney WL. Age-related differences in water and sodium handling after commercial hydration beverage ingestion. *J Appl Physiol.* (2019) 126:1042–8. doi: 10.1152/jappphysiol.01094.2018
108. Maughan RJ, Watson P, Cordery PAA, Walsh NP, Oliver SJ, Dolci A, et al. A randomized trial to assess the potential of different beverages to affect hydration status: development of a beverage hydration index. *Am J Clin Nutr.* (2016) 103:717–23. doi: 10.3945/ajcn.115.114769
109. Maughan RJ, Watson P, Cordery PAA, Walsh NP, Oliver SJ, Dolci A, et al. Sucrose and sodium but not caffeine content influence the retention of beverages in humans under euhydrated conditions. *Int J Sport Nutr Exerc Metab.* (2019) 29:51–60. doi: 10.1123/ijsnem.2018-0047
110. Piccardi N, Manissier P. Nutrition and nutritional supplementation. *Dermatoendocrinol.* (2009) 1:271–4. doi: 10.4161/derm.1.5.9706
111. López-Samanes A, Ortega Fonseca JF, Fernández Elías VE, Borreani S, Maté-Muñoz JL, Kovacs MS. Nutritional ergogenic aids in tennis: a brief review. *Strength Cond J.* (2015) 37:60. doi: 10.1519/SSC.0000000000000141
112. Lun V, Erdman KA, Fung TS, Reimer RA. Dietary supplementation practices in Canadian high-performance athletes. *Int J Sport Nutr Exerc Metab.* (2012) 22:31–7. doi: 10.1123/ijsnem.22.1.31
113. Colls Garrido C, Gómez-Urquiza JL, La Fuente GA, Fernández-Castillo R. Use, effects, and knowledge of the nutritional supplements for the sport in university students. *Nutr Hosp.* (2015) 32:837–44. doi: 10.3305/nh.2015.32.2.8057
114. Parnell JA, Wiens K, Erdman KA. Evaluation of congruence among dietary supplement use and motivation for supplementation in young, Canadian athletes. *J Int Soc Sports Nutr.* (2015) 12:49. doi: 10.1186/s12970-015-0110-y
115. Maughan RJ. Contamination of dietary supplements and positive drug tests in sport. *J Sports Sci.* (2005) 23:883–9. doi: 10.1080/02640410400023258
116. Pallarés JG, López-Samanes A, Fernández-Elías VE, Aguado-Jiménez R, Ortega JF, Gómez C, et al. Pseudoephedrine and circadian rhythm interaction on neuromuscular performance. *Scand J Med Sci Sports.* (2015) 25:e603. doi: 10.1111/sms.12385
117. Bishop D. *This material is the copyright of the original publisher.* Unauthorized copying and distribution is prohibited. Dietary Supplements and Team-Sport Performance. (n.d.).
118. Amawi AT, Moualla DS, Alshuwaier GO, Al-Nuaim AA, Bursais AK, Aljaloud KS, et al. Knowledge and attitude of dietary supplements among Arab Olympic athletes and coaches in preparation program for Tokyo 2020 Olympic games. *Int J Hum Mov Sports Sci.* (2023) 11:368–77. doi: 10.13189/saj.2023.110214
119. Wiens K, Erdman KA, Stadnyk M, Parnell JA. Dietary supplement usage, motivation, and education in young Canadian athletes. *Int J Sport Nutr Exerc Metab.* (2014) 24:613–22. doi: 10.1123/ijsnem.2013-0087
120. Kreider RB, Kalman DS, Antonio J, Ziegenfuss TN, Wildman R, Collins R, et al. International Society of Sports Nutrition position stand: Safety and efficacy of creatine supplementation in exercise, sport, and medicine. *J Int Soc Sports Nutr.* (2017) 14:18. doi: 10.1186/s12970-017-0173-z
121. Maughan RJ, Burke LM, Dvorak J, Larson-Meyer DE, Peeling P, Phillips SM, et al. IOC consensus statement: dietary supplements and the high-performance athlete. *Br J Sports Med.* (2018) 52:439–55. doi: 10.1136/bjsports-2018-099027
122. Wax B, Kerksick CM, Jagim AR, Mayo JJ, Lyons BC, Kreider RB. Creatine for exercise and sports performance, with recovery considerations for healthy populations. *Nutrients.* (2021) 13:1915. doi: 10.3390/nu13061915
123. Cooke MB, Rybalka E, Williams AD, Cribb PJ, Hayes A. Creatine supplementation enhances muscle force recovery after eccentrically-induced muscle damage in healthy individuals. *J Int Soc Sports Nutr.* (2009) 2:6. doi: 10.1186/1550-2783-6-13
124. Rahimi R. Creatine supplementation decreases oxidative DNA damage and lipid peroxidation induced by a single bout of resistance exercise. *J Strength Cond Res.* 25:3448–55. doi: 10.1519/JSC.0b013e3182162f2b
125. Antonio J, Ciccone V. The effects of pre versus post workout supplementation of creatine monohydrate on body composition and strength. *J Int Soc Sports Nutr.* (2013) 10:36. doi: 10.1186/1550-2783-10-36
126. Volek JS, Ratamess NA, Rubin MR, Gomez AL, French DN, McGuigan MM, et al. The effects of creatine supplementation on muscular performance and body composition responses to short-term resistance training overreaching. *Eur J Appl Physiol.* (2004) 91:628–37. doi: 10.1007/s00421-003-1031-z
127. Aedma M, Timpmann S, Lätt E, Ööpik V. Short-term creatine supplementation has no impact on upper-body anaerobic power in trained wrestlers. *J Int Soc Sports Nutr.* (2015) 12:45. doi: 10.1186/s12970-015-0107-6
128. Lanhers C, Pereira B, Naughton G, Trousselard M, Lesage FX, Dutheil F. Creatine supplementation and upper limb strength performance: a systematic review and meta-analysis. *Sports Med.* (2017) 47:163–73. doi: 10.1007/s40279-016-0571-4
129. Jenkinson DM, Harbert AJ. *Supplements and sports.* (2008). Available at: www.aafp.org/afp.
130. Shearer J, Graham TE. Performance effects and metabolic consequences of caffeine and caffeinated energy drink consumption on glucose disposal. *Nutr Rev.* (2014) 72:121–36. doi: 10.1111/nure.12124
131. Goldstein ER, Ziegenfuss T, Kalman D, Kreider R, Campbell B, Wilborn C, et al. International society of sports nutrition position stand: Caffeine and performance. *J Int Soc Sports Nutr.* (2010) 18:1. doi: 10.1186/s12970-020-00383-4
132. Tallis J, Duncan MJ, James RS. What can isolated skeletal muscle experiments tell us about the effects of caffeine on exercise performance? *Br J Pharmacol.* (2015) 172:3703–13. doi: 10.1111/bph.13187
133. Davis JK, Green JM. Caffeine and anaerobic performance: ergogenic value and mechanisms of action. *Sports Med.* (2009) 39:813–32. doi: 10.2165/11317770-000000000-00000
134. Tarnopolsky MA. Effect of caffeine on the neuromuscular system—potential as an ergogenic aid. *Appl Physiol Nutr Metab.* (2008) 33:1284–9. doi: 10.1139/H08-121
135. Stojanović E, Stojilković N, Scanlan AT, Dalbo VJ, Stanković R, Antić V, et al. Acute caffeine supplementation promotes small to moderate improvements in performance tests indicative of in-game success in professional female basketball players. *Appl Physiol Nutr Metab.* (2019) 44:849–56. doi: 10.1139/apnm-2018-0671
136. Khan IA, Nasiruddin M, Haque SF, Khan RA. Comparative evaluation of efficacy and safety profile of rhabarb and α -keto analogs of essential amino acids supplementation in patients with diabetic nephropathy. *Saudi J Kidney Dis Transpl.* (2016) 27:710–6. doi: 10.4103/1319-2442.185227

137. Phillips SM, van Loon LJC. Dietary protein for athletes: from requirements to optimum adaptation. *J Sports Sci.* (2011) 29:29–38. doi: 10.1080/02640414.2011.619204
138. Hoffman JR, Ratamess NA, Faigenbaum AD, Ross R, Kang J, Stout JR, et al. Short-duration β -alanine supplementation increases training volume and reduces subjective feelings of fatigue in college football players. *Nutr Res.* (2008) 28:31–5. doi: 10.1016/j.nutres.2007.11.004
139. Tipton KD, Elliott TA, Cree MG, Wolf SE, Sanford AP, Wolfe RR. Ingestion of casein and whey proteins result in muscle anabolism after resistance exercise. *Med Sci Sports Exerc.* (2004) 36:2073–81. doi: 10.1249/01.mss.0000147582.99810.c5
140. Middleton N, Jelen P, Bell G. Whole blood and mononuclear cell glutathione response to dietary whey protein supplementation in sedentary and trained male human subjects. *Int J Food Sci Nutr.* (2004) 55:131–41. doi: 10.1080/096374080410001666504
141. Tang JE, Moore DR, Kujbida GW, Tarnopolsky MA, Phillips SM. Ingestion of whey hydrolysate, casein, or soy protein isolate: effects on mixed muscle protein synthesis at rest and following resistance exercise in young men. *J Appl Physiol.* (2009) 107:987–92. doi: 10.1152/jappphysiol.00076.2009
142. Wu G, Wu Z, Dai Z, Yang Y, Wang W, Liu C, et al. Dietary requirements of “nutritionally non-essential amino acids” by animals and humans. *Amino Acids.* (2013) 44:1107–13. doi: 10.1007/s00726-012-1444-2
143. Rieu I, Balage M, Sornet C, Giraudet C, Pujos E, Grizard J, et al. Leucine supplementation improves muscle protein synthesis in elderly men independently of hyperaminoacidemia. *J Physiol.* (2006) 575:305–15. doi: 10.1113/jphysiol.2006.110742
144. Yoon MS. The emerging role of branched-chain amino acids in insulin resistance and metabolism. *Nutrients.* (2016) 8:405. doi: 10.3390/nu8070405
145. Burd NA, Holwerda AM, Selby KC, West DWD, Staples AW, Cain NE, et al. Resistance exercise volume affects myofibrillar protein synthesis and anabolic signalling molecule phosphorylation in young men. *J Physiol.* (2010) 588:3119–30. doi: 10.1113/jphysiol.2010.192856
146. Shimomura Y, Shimomura Y, Inaguma A, Watanabe S, Yamamoto Y, Muramatsu Y, et al. Branched-chain amino acid supplementation before squat exercise and delayed-onset muscle soreness. *Int J Sport Nutr Exerc Metab.* (2010) 20:236–44. doi: 10.1123/ijsnem.20.3.236
147. Nosaka K, Sacco PN, Mawatari KN. Effects of amino acid supplementation on muscle soreness and damage. *Int J Sport Nutr Exerc Metab.* (2006) 16:620–35. doi: 10.1123/ijsnem.16.6.620
148. Areces F, Salinero JJ, Abian-Vicen J, González-Millán C, Gallo-Salazar C, Ruiz-Vicente D, et al. A 7-day oral supplementation with branched-chain amino acids was ineffective to prevent muscle damage during a marathon. *Amino Acids.* (2014) 46:1169–76. doi: 10.1007/s00726-014-1677-3
149. Liu TH, Wu CL, Chiang CW, Lo YW, Tseng HF, Chang CK. No effect of short-term arginine supplementation on nitric oxide production, metabolism and performance in intermittent exercise in athletes. *J Nutr Biochem.* (2009) 20:462–8. doi: 10.1016/j.jnutbio.2008.05.005
150. Willoughby DS, Boucher T, Reid J, Skelton G, Clark M. Effects of 7 days of arginine-alpha-ketoglutarate supplementation on blood flow, plasma L-arginine, nitric oxide metabolites, and asymmetric dimethyl arginine after resistance exercise. *Int J Sport Nutr Exerc Metab.* (2011) 21:291–9. doi: 10.1123/ijsnem.21.4.291
151. Bescós RL, Sureda A, Tur JA, Pons A. The effect of nitric-oxide-related supplements on human performance. *Sports Med.* 42:99–117. doi: 10.2165/11596860-000000000-00000
152. Santana JO, de Freitas MC, dos Santos DM, Rossi FE, Lira FS, Rosa-Neto JC, et al. Beta-alanine supplementation improved 10-km running time trial in physically active adults. *Front Physiol.* (2018) 9:1105. doi: 10.3389/fphys.2018.01105
153. Harris RC, Tallon MJ, Dunnett M, Boobis L, Coakley J, Kim HJ, et al. The absorption of orally supplied β -alanine and its effect on muscle carnosine synthesis in human vastus lateralis. *Amino Acids.* (2006) 30:279–89. doi: 10.1007/s00726-006-0299-9
154. Quesnele JJ, Laframboise MA, Wong JJ, Kim P, Wells GD. The effects of beta-alanine supplementation on performance: a systematic review of the literature. *Int J Sport Nutr Exerc Metab.* (2014) 24:14–27. doi: 10.1123/ijsnem.2013-0007
155. Hobson RM, Saunders B, Ball G, Harris RC, Sale C. Effects of β -alanine supplementation on exercise performance: a meta-analysis. *Amino Acids.* (2012) 43:25–37. doi: 10.1007/s00726-011-1200-z
156. Ducker KJ, Dawson B, Wallman KE. Effect of Beta-alanine supplementation on 800-m running performance. *Int J Sport Nutr Exerc Metab.* (2013) 23:554–61. doi: 10.1123/ijsnem.23.6.554
157. Stegen S, Bex T, Vervaet C, Vanhee L, Achten E, Derave W. β -Alanine dose for maintaining moderately elevated muscle carnosine levels. *Med Sci Sports Exerc.* (2014) 46:1426–32. doi: 10.1249/MSS.0000000000000248
158. Ostfeld I, Hoffman JR. The effect of β -alanine supplementation on performance, cognitive function and resiliency in soldiers. *Nutrients.* (2023) 15:1039. doi: 10.3390/nu15041039
159. Maughan RJ, Burke LM, Dvorak J, Larson-Meyer DE, Peeling P, Phillips SM, et al. IOC consensus statement: dietary supplements and the high-performance athlete. *Int J Sport Nutr Exerc Metab.* (2018) 28:104–25. doi: 10.1123/ijsnem.2018-0020
160. Buell JL, Franks R, Ransone J, Powers ME, Laquale KM, Carlson-Phillips A. National Athletic Trainers’ association position statement: evaluation of dietary supplements for performance nutrition. *J Athl Train.* (2013) 48:124–36. doi: 10.4085/1062-6050-48.1.16
161. Ivy J. *Nutrient timing: The future of sports nutrition.* Laguna Beach, CA: Basic Health Publications, Inc. (2004).
162. Ormsbee MJ, Bach CW, Baur DA. Pre-exercise nutrition: the role of macronutrients, modified starches and supplements on metabolism and endurance performance. *Nutrients.* (2014) 6:1782–808. doi: 10.3390/nu6051782
163. Carpenter TM, Fox EL. The effect of muscular work upon the respiratory exchange of man after the ingestion of glucose and of fructose. 1. The respiratory quotient and the mechanics of breathing. *Arbeitsphysiologie.* (1931) 4:532–69.
164. Haldi J, Wynn W. Observations on efficiency of swimmers as related to some changes in pre-exercise nutriment. *J Nutr.* (1946) 31:525–32. doi: 10.1093/jn/31.5.525
165. Hargreaves M, Costill DL, Fink WJ, King DS, Fielding RA. Effect of pre-exercise carbohydrate feedings on endurance cycling performance. *Med Sci Sports Exerc.* (1987) 19:33–6.
166. Sherman WM, Brodowicz G, Wright DA, Allen WK, Simonsen J, Dernbach A. Effects of 4 h preexercise carbohydrate feedings on cycling performance. *Med Sci Sports Exerc.* (1989) 21:598–604.
167. Sherman WM, Peden C, Wright DA. Carbohydrate feedings 1 h before exercise improves cycling performance. *Am J Clin Nutr.* (1991) 54:866–70. doi: 10.1093/ajcn/54.5.866
168. Achten J, Jeukendrup AE. The effect of pre-exercise carbohydrate feedings on the intensity that elicits maximal fat oxidation. *J Sports Sci.* (2003) 21:1017–25. doi: 10.1080/02640410310001641403
169. Escobar KA, VanDusseldorp TA, Kerksick CM. Carbohydrate intake and resistance-based exercise: are current recommendations reflective of actual need? *Br J Nutr.* (2016) 116:2053–65. doi: 10.1017/S0007114516003949
170. Vøllestad NK, Tabata I, Medbø JI. Glycogen breakdown in different human muscle fibre types during exhaustive exercise of short duration. *Acta Physiol Scand.* (1992) 144:135–41. doi: 10.1111/j.1748-1716.1992.tb09278.x
171. Bergström J, Hermansen L, Hultman E, Saltin B. Diet, muscle glycogen and physical performance. *Acta Physiol Scand.* (1967) 71:140–50. doi: 10.1111/j.1748-1716.1967.tb03720.x
172. Karlsson J, Saltin B. Diet, muscle glycogen, and endurance performance. *J Appl Physiol.* (1971) 31:203–6. doi: 10.1152/jappphysiol.1971.31.2.203
173. Ørtenblad N, Nielsen J, Saltin B, Holmberg H. Role of glycogen availability in sarcoplasmic reticulum Ca²⁺ kinetics in human skeletal muscle. *J Physiol.* (2011) 589:711–25. doi: 10.1113/jphysiol.2010.195982
174. Jeukendrup AE, Jentjens R. Oxidation of carbohydrate feedings during prolonged exercise: current thoughts, guidelines and directions for future research. *Sports Med.* (2000) 29:407–24. doi: 10.2165/00007256-200029060-00004
175. Jentjens RLPG, Moseley L, Waring RH, Harding LK, Jeukendrup AE. Oxidation of combined ingestion of glucose and fructose during exercise. *J Appl Physiol.* (2004) doi: 10.1152/jappphysiol.00974.2003
176. Cermak NM, van Loon LJC. The use of carbohydrates during exercise as an ergogenic aid. *Sports Med.* (2013) 43:1139–55. doi: 10.1007/s40279-013-0079-0
177. Currell K, Jeukendrup A. Superior endurance performance with ingestion of multiple transportable carbohydrates. *Med Sci Sports Exerc.* (2008) 40:275–81. doi: 10.1249/mss.0b013e31815adff9
178. Kloby Nielsen LL, Tandrup Lambert MN, Jeppesen PB. The effect of ingesting carbohydrate and proteins on athletic performance: a systematic review and meta-analysis of randomized controlled trials. *Nutrients.* (2020) 12:1483. doi: 10.3390/nu12051483
179. Betts JA, Stevenson E. Should protein be included in CHO-based sports supplements? *Med Sci Sports Exerc.* (2011) 43:1244–50. doi: 10.1249/MSS.0b013e31821b0dce
180. Ivy JL, Ferguson-Stegall LM. Nutrient timing: the means to improved exercise performance, recovery, and training adaptation. *Am J Lifestyle Med.* (2014) 8:246–59. doi: 10.1177/1559827613502444
181. Beelen M, Burke LM, Gibala MJ, Van Loon LJC. Nutritional strategies to promote postexercise recovery. *Int J Sport Nutr Exerc Metab.* (2010) 20:515–32. doi: 10.1123/ijsnem.20.6.515
182. Goodyear LJ, Hirshman MF, King PA, Horton ED, Thompson CM, Horton ES. Skeletal muscle plasma membrane glucose transport and glucose transporters after exercise. *J Appl Physiol.* (1990) 68:193–8. doi: 10.1152/jappphysiol.1990.68.1.193
183. Hawley JA, Tipton KD, Millard-Stafford ML. Promoting training adaptations through nutritional interventions. *J Sports Sci.* (2006) 24:709–21. doi: 10.1080/02640410500482727
184. Ivy J. *Nutrient timing the future of sports nutrition part I: nutrient timing principles.* 1: Nutrient Timing 2: Muscle energy systems and fuel utilization 3: The influence of hormones on muscle growth and development Part II: Nutrient timing system 4: NTS energy phase 5: NTS anabolic phase 6: NTS growth phase Part III: Incorporating nutrient timing into your training 7: Making nutrient timing work for you 8: The NTS nutrition program.

185. Ivy JL, Katz AL, Cutler CL, Sherman WM, Coyle EF. Muscle glycogen synthesis after exercise: Effect of time of carbohydrate ingestion. *J Appl Physiol* (1985). (1988) 64:1480–5. doi: 10.1152/jappl.1988.64.4.1480
186. Hawley JA, Burke LM. Effect of meal frequency and timing on physical performance. *Br J Nutr*. (1997) 77:S91–S103. doi: 10.1079/BJN19970107
187. Walker JL, Heigenhauser F, Hultman E, Spriet LL. Dietary carbohydrate, muscle glycogen content, and endurance performance in well-trained women. *J Appl Physiol*. (2000) 88:2151–8. doi: 10.1152/jappl.2000.88.6.2151
188. Coyle EF, Coggan AR, Hemmert K, Lowe RC, Walters TJ, Hem-Mert MK, et al. (2018). Available at: <https://www.physiology.org/journal/jappl>.
189. Pettersson S, Edin F, Bakkman L, McGawley K. Effects of supplementing with an 18% carbohydrate-hydrogel drink versus a placebo during whole-body exercise in -5 °C with elite cross-country ski athletes: a crossover study. *J Int Soc Sports Nutr*. (2019) 16:46. doi: 10.1186/s12970-019-0317-4
190. Tokmakidis SP, Karamanolis IA. Effects of carbohydrate ingestion 15 min before exercise on endurance running capacity. *Appl Physiol Nutr Metab*. (2008) 33:441–9. doi: 10.1139/H08-012
191. Burdon CA, Spronk I, Cheng HL, O'Connor HT. Effect of glycemic index of a pre-exercise meal on endurance exercise performance: A systematic review and Meta-analysis. *Sports Med*. (2017) 47:1087–101. doi: 10.1007/s40279-016-0632-8
192. *Nutrition and Athletic Performance*. (2000). Available at: <http://www.acsm-msse.org>.
193. Venkatraman JT, Pendergast DR. Effect of dietary intake on immune function in athletes. *Sports Med*. 32:323–37. doi: 10.2165/00007256-200232050-00004
194. Candow DG, Chilibeck PD. Timing of creatine or protein supplementation and resistance training in the elderly. *Appl Physiol Nutr Metab*. (2008) 33:184–90. doi: 10.1139/H07-139
195. Hulmi JJ, Lockwood CM, Stout JR. Effect of protein/essential amino acids and resistance training on skeletal muscle hypertrophy: a case for whey protein. *Nutr Metab*. (2010) 7:51. doi: 10.1186/1743-7075-7-51
196. Kukuljan S, Nowson CA, Sanders K, Daly RM. Effects of resistance exercise and fortified milk on skeletal muscle mass, muscle size, and functional performance in middle-aged and older men: an 18-mo randomized controlled trial. *J Appl Physiol*. (2009) 107:1864–73. doi: 10.1152/japplphysiol.00392.2009
197. Guest NS, Horne J, Vanderhout SM, El-Sohemy A. Sport nutrigenomics: Personalized nutrition for athletic performance. *Front Nutr*. (2019) 6:8. doi: 10.3389/fnut.2019.00008
198. Carbone JW, Pasiakos SM. Dietary protein and muscle mass: translating science to application and health benefit. *Nutrients*. (2019) 11:1136. doi: 10.3390/nu11051136
199. Brancaccio M, Mennitti C, Cesaro A, Fimiani F, Vano M, Gargiulo B, et al. The biological role of vitamins in athletes' muscle, heart and microbiota. *Int J Environ Res Public Health*. (2022) 19:1249. doi: 10.3390/ijerph19031249
200. Beck KL, Thomson JS, Swift RJ, Von Hurst PR. Role of nutrition in performance enhancement and postexercise recovery. *Open Access J Sports Med*. (2015) 6:259–67. doi: 10.2147/OAJSM.S33605