

American College of Sports Medicine Position Stand. Resistance Training Prescription for Muscle Function, Hypertrophy, and Physical Performance in Healthy Adults: An Overview of Reviews

BRAD S. CURRIER¹, ALYSHA C. D'SOUZA¹, MARIA A. FIATARONE SINGH^{2,3}, CAROLINE V. LOWISZ¹, ERIC S. RAWSON⁴, BRAD J. SCHOENFELD⁵, ABBIE E. SMITH-RYAN⁶, JEREMY P. STEEN⁷, GWENDOLYN A. THOMAS⁸, N. TRAVIS TRIPLETT⁹, TYRONE A. WASHINGTON¹⁰, TIMOTHY J. WERNER¹¹, and STUART M. PHILLIPS¹

¹Department of Kinesiology, Faculty of Science, McMaster University, Hamilton, ON, CANADA; ²School of Health Sciences and Sydney Medical School, Faculty of Medicine and Health, University of Sydney, Sydney, NSW, AUSTRALIA; ³Hinda and Arthus Marcus Institute for Aging Research, Hebrew SeniorLife, Roslindale, MA; ⁴Department of Health, Nutrition and Exercise Science, Messiah University, Mechanicsburg, PA; ⁵Department of Exercise Science and Recreation, CUNY Lehman College, Bronx, NY; ⁶Department of Exercise and Sport Science, University of North Carolina, Chapel Hill, NC; ⁷Institute of Health Policy, Management and Evaluation, Dalla Lana School of Public Health, University of Toronto, Toronto, ON, CANADA; ⁸Department of Kinesiology, The Pennsylvania State University, University Park, PA; ⁹Department of Kinesiology, Appalachian State University, Boone, NC; ¹⁰Department of Health, Human Performance and Recreation, University of Arkansas, Fayetteville, AR; ¹¹Department of Exercise Science, Salisbury University, Salisbury, MD

SUMMARY

CURRIER, B. S., A. C. D'SOUZA, M. A. F. SINGH, C. V. LOWISZ, E. S. RAWSON, B. J. SCHOENFELD, A. E. SMITH-RYAN, J. P. STEEN, G. A. THOMAS, N. T. TRIPLETT, T. A. WASHINGTON, T. J. WERNER, and S. M. PHILLIPS. American College of Sports Medicine Position Stand. Resistance Training Prescription for Muscle Function, Hypertrophy, and Physical Performance in Healthy Adults: An Overview of Reviews. *Med. Sci. Sports Exerc.*, Vol. 58, No. 4, pp. 851-872, 2026. **Purpose:** The aim of this overview of reviews was to determine the impact of resistance training (RT) prescription on muscle function and hypertrophy, utilizing evidence synthesis methods. It updates the American College of Sports Medicine 2009 Position Stand, "Progression models in resistance training for healthy adults." **Data sources:** Ovid MEDLINE(R) ALL, Ovid Emcare, Ovid Embase, Cochrane Database of Systematic Reviews, EBSCOhost SPORTDiscus, and Web of Science Core Collection current to October 2024. **Eligibility criteria:** Eligible systematic reviews synthesized randomized trials of healthy adults (≥ 18 yr) who completed RT (≥ 6 wk; range: 6–52 wk), compared with a group that completed no exercise or an alternative RT program, and reported the change in muscle function, size, or physical performance. **Results:** We synthesized data from 137 systematic reviews ($>30,000$ participants). Compared with no exercise (control), RT significantly improved muscle strength, size (hypertrophy), power, endurance, contraction velocity, gait speed, balance, and multiple physical function outcomes. Few RT prescription (RTx) variables affected primary adaptations. However, voluntary strength was enhanced by lifting heavier loads ($\geq 80\%$ one-repetition maximum), through a complete range of motion, for 2–3 sets, at the beginning of training sessions, and ≥ 2 sessions/wk. Muscle hypertrophy was enhanced by higher volumes (≥ 10 sets/wk) and eccentric overload. Power was enhanced by moderate loads (30%–70% one-repetition maximum), low-to-moderate volume (≤ 24 repetitions-sets), Olympic-style weightlifting, and power RT (fast concentric phase). Power RT enhanced physical function. Training to momentary muscle fatigue, equipment type, exercise complexity, set structure, time under tension, blood flow restriction, and periodization did not consistently impact training outcomes. **Conclusions:** Healthy adults should perform progressive RT, with variable prescription consistent with our findings, to improve muscle function, size, and physical performance. Muscle strength, hypertrophy, power, and certain components of physical function can be enhanced by manipulating the RT variables highlighted. **Key Words:** HYPERTROPHY, PHYSICAL FUNCTION, RESISTANCE TRAINING, SKELETAL MUSCLE, STRENGTH

Address for correspondence: Stuart M. Phillips, PhD, Department of Kinesiology, McMaster University, 1280 Main Street West, Hamilton, ON L8S 4K1, Canada; E-mail: phillis@mcmaster.ca.

Submitted for publication February 2025

Accepted for publication September 2025

Supplemental digital content is available for this article. Direct URL citations appear in the printed text and are provided in the HTML and PDF versions of this article on the journal's Web site (www.acsm-msse.org).

0195-9131/26/584-851/872

MEDICINE & SCIENCE IN SPORTS & EXERCISE®

Copyright © 2025 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American College of Sports Medicine. This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

DOI: 10.1249/MSS.0000000000003897

Exercise is crucial for health throughout the human lifespan, and muscle strength and function are essential components of fitness. Resistance training (RT), also known as strength or weight training, is a specialized method of physical conditioning in which muscles are exercised by contracting against external resistance, such as free weights, machines, resistance bands, water, or body weight, through isometric, isotonic, or isokinetic actions, progressively increasing force output to improve muscular strength, power, endurance, and overall health and sports performance. The benefits of RT are being increasingly appreciated as health-promoting behavior (1,2), but were perhaps first documented to be beneficial beyond personal strength development by Captain Thomas DeLorme, who recognized the benefits of RT in wounded soldiers (3). Beyond the hallmark improvements in skeletal muscle mass and function, the benefits of engaging in RT include reduced mortality and risk for and management of cardiovascular disease, cancer, and diabetes (4–6), reduced depression (7,8), and improved sleep quality (9).

Most guidelines call for healthy adults to complete “muscle-strengthening activities at moderate or greater intensity that involve all major muscle groups on two or more days a week” (2). Exercise programs are generally constructed by manipulating six factors comprising the framework FITT-VP: Frequency, Intensity, Time, Type, Volume, Pattern, and Progression (10). Prescription of variable RT (RTx), however, involves several variables within each category that are inherent to any practice of RT. However, a barrier to engaging in RT is that people and practitioners often lack understanding of how to prescribe RT (5). Thus, RTx guidelines are required to support healthcare practitioners and exercise professionals when designing RT programs.

The American College of Sports Medicine (ACSM) 2009 Position Stand, “Progression Models in Resistance Training for Healthy Adults” (11), summarized, at the time, the available evidence for RTx variables, providing guidelines to enhance RT adaptations. Research on the topic of RT has expanded significantly since the publication of that Position Stand; in fact, a simple PubMed search for “resistance training” yields over 30,000 new results since 2009, indicating a need for an update. Evidence synthesis methods have also advanced considerably, and both the current (11) and prior (12) Position Stands were criticized (13,14) for lacking evidence-based rigor. Hence, to provide contemporary, evidence-based guidance to minimize bias, an updated RTx Position Stand was required, utilizing contemporary search and evidence grading methodologies.

Provided ample systematic reviews and meta-analyses, an overview of reviews can systematically summarize an abundance of information (13). An overview of reviews (i.e., umbrella review: a review of systematic reviews)

is a systematic collection and assessment of available evidence that provides a comprehensive, user-friendly summary of a research topic, enabling practitioners and professionals to make evidence-based decisions without assimilating the results of numerous systematic reviews and meta-analyses (14,15).

The purpose of this overview of systematic reviews was to provide an updated, evidence-based summary of the impact of RTx variables on various outcomes relevant to RT in healthy adults. The outcomes of interest included muscle hypertrophy, strength, power, endurance, contraction velocity, and physical function (e.g., gait speed, balance, and stair climbing). The current document updates the ACSM 2009 Position Stand entitled “Progression Models in Resistance Training for Healthy Adults” (11).

METHODS

Protocol and Registration

This review was prospectively registered on the International Platform of Registered Systematic Review and Meta-analysis Protocols (INPLASY202360071; <https://inplasy.com/inplasy-2023-6-0071/>) and conducted in alignment with the Preferred Reporting Items for Overviews of Reviews (PRIOR) (16). The completed PRIOR checklist is contained in Supplemental Appendix 1, Supplemental Digital Content, <https://links.lww.com/MSS/D323>.

Eligibility Criteria

The complete eligibility criteria for systematic reviews to be included are shown in Table 1. Briefly, systematic reviews of randomized trials were included when healthy adults completed an RT program for at least 6 wk and were compared with a group that completed no exercise (control [CTRL]) or a distinctly different RT program (Table 2; i.e., FITT-VP component different between groups). In most cases, reviews examined people with minimal or no RT experience (i.e., novice); however, some reviews included more advanced trainees. Previously, we showed that training experience had a minimal impact on strength and hypertrophy outcomes (17); nonetheless, much of the evidence synthesized here is from inexperienced trainees. “Healthy adults” was operationalized as humans (≥ 18 -yr old) with no defined disease(s), including obesity, sarcopenia, and physical frailty. Body weight-supported RT was a portion of some reviews, but was not studied as a stand-alone intervention except in the case of the Nordic hamstring curl. Standard, or non-specialized, RT was defined as any RT in eligible reviews that was not specifically defined and thus excluded specific forms of RT, including power RT (explicitly stated concentric phase performed as quickly as possible),

TABLE 1. Eligibility criteria for inclusion.

	Inclusion Criteria
Population	<ul style="list-style-type: none"> • Healthy humans ≥ 18-yr old with no defined disease(s) • Any training status (novice or trained)
Intervention	<ul style="list-style-type: none"> • Resistance training interventions spanning at least 6 wk with a minimum of 12 exposures • If a supplement, nutritional, or other cointervention (e.g., behavioral therapy, medication, counseling) is applied, it must be received by intervention and comparator groups
Comparator	<ul style="list-style-type: none"> • Distinct resistance training prescription, as defined by FITT-VP (Frequency, Intensity, Time, Type, Volume, Pattern, Progression) principle, and/or • Nonexercise control group (performing no RT) and/or alternative exercise control conceived as a sham intervention (e.g., stretching)
Outcome	<ul style="list-style-type: none"> • Reported pre- and postintervention change in muscle function (strength, power, endurance, contraction velocity, physical function) or hypertrophy between intervention and comparator arm(s)
Study design	<ul style="list-style-type: none"> • A systematic review (including overviews of reviews) of randomized trials with or without statistical synthesis (e.g., meta-analysis, network meta-analysis, meta-regression)

Olympic-style weightlifting (RT with Olympic-style lifting movements), and velocity-based RT (movement velocity thresholds used to prescribe RT). We acknowledge that this is a broad definition that encompasses several variables, but we have used the definitions of the RTx provided within the reviews we compared. Eligible reviews needed to report the change of at least one muscle function (e.g., strength, power, and physical function) or muscle size (hypertrophy) outcome from pre- to postintervention. Unless otherwise specified, strength was voluntary isotonic one-repetition maximum (1RM) in the same mode as that in which the training was performed. Records were defined as a “systematic review” if they were titled as an evidence synthesis (e.g., systematic review, meta-analysis, and umbrella review) or used a specific systematic search strategy and eligibility criteria. The included reviews were not separated by participant age due to the number of eligible systematic reviews that included participants with participants who were both younger and older than the cutoff of 55-yr old. Thus, our recommendations are evidence-based across all ages.

Information Sources and Search Strategy

The systematic search strategy was executed in October 2024 in Ovid MEDLINE(R) ALL (1946 to current), Ovid Emcare (1995 to current), Ovid Embase (1974 to current), Cochrane Database of Systematic Reviews (2005 to current), EBSCOhost SPORTDiscus, and Web of Science Core Collection. Trained librarians developed the search strategies. Searches conducted on the Ovid platform were limited to English-language records, and no additional limits or filters were applied. The complete search strategy is reported in Supplemental Appendix 2, Supplemental Digital Content, <https://links.lww.com/MSS/D323>.

Record Selection and Data Collection

Four reviewers (B. S. C., C. V. L., A. C. D., and J. P. S.) independently screened all records (title/abstract and full text) and extracted data from eligible reviews in duplicate, with discrepancies resolved by group consensus. Relevant data from eligible overlapping records were included and extracted. Authors of reviews with missing data were contacted via email with a request for the missing data, and

TABLE 2. Resistance training prescription variables.

RTx Variable	Definition
Frequency	The number of days per week RT is performed.
Load	The amount of weight lifted per repetition and often prescribed as a proportion of maximal strength (e.g., %1RM) or maximal number of repetitions possible with a given load (RM)
Failure	Completing repetitions until volitional failure occurs when concentric movement is no longer possible.
Time under tension	The amount of time supporting weight. Time under tension per repetition is the total time required to complete one repetition.
Time of day	When RT is performed in the day (e.g., morning)
Inter-set rest	The amount of rest time between sets.
Exercise order	The order of different training forms (e.g., aerobic and resistance) or specific RT exercises are performed within a training session.
Type	<p>Blood flow restriction: Locally reducing blood flow to and from exercising muscles.</p> <p>Free-weight RT: Weights that can be moved in space freely (e.g., barbell).</p> <p>Machine RT: Machines permitting fixed movement (e.g., leg press).</p> <p>Unstable RT: RT performed on an unstable surface (e.g., pressurized ball).</p> <p>Variable load: Altering load mid-repetition.</p>
Contraction type	<p>Eccentric overload: Increased load or time under tension during eccentric movement.</p> <p>Traditional: Proportionally completing both concentric and eccentric phases of a movement.</p>
Technique	<p>Olympic-style weightlifting: Snatch and clean-and-jerk movements.</p> <p>Partial range of motion: Performing RT through an incomplete range of motion, compared with RT as typically performed through the complete range of motion for the joint(s) involved in a movement.</p> <p>Power RT: Intentionally performing the concentric phase of each lift at maximal volitional speed.</p>
Volume (sets)	The number of sets (a group of repetitions without resting) completed per exercise.
Set structure	<p>Drop sets: Performing sets to failure then reducing load and minimal inter-set rest.</p> <p>Cluster: Traditional inter-set rest with preplanned intraset rest periods.</p> <p>Complex: Sets with heavy loads followed by sets with lighter loads.</p> <p>Contrast: Alternating heavy and light loads set-to-set.</p> <p>Rest redistribution: Short rest periods between each repetition.</p> <p>Traditional: Performing a set without rest until all repetitions are completed.</p>
Periodization	<p>Periodized: Manipulating RTx during a program to maximize adaptations.</p> <p>Block: Dividing program into multiweek blocks with distinct training goals.</p> <p>Linear (traditional): Increasing load and reducing volume during the program.</p> <p>Nonperiodized: Not manipulating RTx during the program.</p> <p>Undulating (nonlinear): Daily or weekly RTx manipulation.</p>

%1RM, percentage of one-repetition maximum; RM, repetition maximum; RT, resistance training; RTx, resistance training prescription.

WebPlotDigitizer (version 4; <https://automeris.io/>) was used if data needed to be extracted from figures. Record screening and data extraction were completed using the systematic review software Covidence (<https://www.covidence.org/>). The complete list of data items sought is reported in Supplemental Appendix 3, Supplemental Digital Content, <https://links.lww.com/MSS/D323>.

Methodological Quality Assessment and Evidence Synthesis

The methodological quality of each included review was assessed independently by two reviewers using the AMSTAR (A Measurement Tool to Assess Systematic Reviews) tool, which yields a score ranging from 1 to 11 that incorporates assessment of publication bias (Supplemental Appendix 4, Supplemental Digital Content, <https://links.lww.com/MSS/D323>) (18,19). Outcome data were tabulated as collected, and no sensitivity analyses were conducted. Heterogeneity was reported as the I^2 statistic (meta-analyses) or the fraction of reviews showing a significant effect (systematic reviews). An outcome-level (bottom-line) statement and standardized effectiveness statement (Supplemental Appendix 5, Supplemental Digital Content, <https://links.lww.com/MSS/D323>) were produced by considering the methodological quality and extracted data (20). Outcome-level quality of evidence (QoE) was calculated using a method based on the Grading of Recommendations Assessment, Development and Evaluation approach for primary evidence (21). This method incorporates the design (meta-analysis: yes/no) and methodological quality (AMSTAR score) of each included review (Supplemental Appendix 6, Supplemental Digital Content, <https://links.lww.com/MSS/D323>). There remains no standardized method to evaluate the certainty of evidence in overviews of reviews (15,22), so a summary percentage scores were calculated to demonstrate the quality of evidence contributing to each conclusion. The summary percentage was calculated within each prescription variable for all three directions of evidence (impactful, not impactful, and cannot determine) by dividing the average QoE by four (the maximum QoE score) to yield a percentage score ranging from 0% (lowest possible QoE) to 100% (highest possible QoE). This method has been used in previous overviews of reviews (18,20,23). The effectiveness (impact) of RTx variables for each outcome was assessed based on standardized effectiveness statements and the quantity of evidence (24). The reviews, which contributed evidence for the impact of a prescription variable, were independently scrutinized to comment on favorable RTx parameters for improving each outcome.

Overlap of Papers

An issue with overviews of reviews is the possibility of “double counting” (or more) papers that are included

in more than one review (25). Such counting may unduly affect the results of the analysis, resulting in spurious levels of precision and confidence in the outcomes (26). We employed the corrected covered area (CCA) index to quantify the degree of overlap between systematic reviews to be pooled in an overview of reviews (27). To obtain an estimate of the degree of publication overlap, we calculated the CCA for strength using the ccaR package (28), arguably the most relevant outcome of our analysis.

RESULTS

Included Reviews

The systematic search yielded 5751 records following duplicate removal, and 137 systematic reviews were included in this overview of reviews (Fig. 1). The AMSTAR scores for included reviews ranged from 1 (lowest) to 9 (highest) out of a possible 11 (Supplemental Appendix 7, Supplemental Digital Content, <https://links.lww.com/MSS/D323>). The conflict-of-interest statement for each included review is reported in Supplementary Appendix 8, Supplemental Digital Content, <https://links.lww.com/MSS/D323>. The details, bottom-line statement, standardized effectiveness statement, and QoE for each review are reported in the Supplemental Appendices, Supplemental Digital Content, <https://links.lww.com/MSS/D323>, according to the outcome.

Strength

CCA analysis of strength across all included reviews showed only “moderate” overlap, defined as between 6% and 10% of primary papers appearing in two or more reviews (27). So the results of our analyses for this variable (for which there were more reviews than for any other variable) were not unduly affected by primary data overlap.

The effects of RT versus CTRL on strength and the impact of distinct RTx variables are summarized in Tables 3 and 4, respectively. The results for each review are reported in Supplemental Appendix 11, Supplemental Digital Content, <https://links.lww.com/MSS/D323>. Compared with CTRL, strength was impacted by engaging in standard RT (17,24,29–56), circuit RT (57–59), elastic band RT (60–62), home-based RT (63,64), and velocity-based RT (65,66). There were insufficient data to determine if eccentric flywheel RT (67), Nordic hamstring RT (68), Olympic-style weightlifting (69), and unstable surface RT (70) impacted strength compared with CTRL. In comparisons between distinct RTx, strength was positively affected by training session frequency (24,71–76), load (24,77–84), eccentric flywheel RT versus standard RT (85), range of motion (86,87), volume (24,88–93), and exercise order (24,94–96), but strength was not affected in programs with lifting to fatigue/failure (24,97–99), machines versus free-weight RT (100), unstable surfaces (70), time under tension (24,101), time




FIGURE 1—Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flow diagram of review selection.

of day (24,102), inter-set rest (24,103), contraction type (24,104), power training (82,105,106), or set structure (24,107–109). There were insufficient data to determine if strength was affected by blood flow restriction (110–112), Olympic-style weightlifting (69), intrasession autoregulation (113,114), varied exercise selection (115), periodization (24,116–118), or concurrent training (i.e., aerobic and RT in the same training session) (119–124).

The impact of unilateral RT on the untrained, contralateral limb (i.e., cross-education) was only reported for the strength outcome. The impact of RT versus CTRL on strength in the untrained contralateral limb is summarized in Table 3, and results for each review are reported in Supplemental Appendix 11, Supplemental Digital Content, <https://links.lww.com/MSS/D323>. Unilateral RT impacted strength (125,126) such that strength improved in the untrained, contralateral limb. Comparing RTx, strength was enhanced by RT performed with higher frequency (≥ 2 d/wk, though the upper limit cannot, from our analysis, be undetermined), higher loads (dose-response), eccentric flywheel devices compared with standard RT, full range of motion, higher

volume (multiple sets), and at the beginning of training sessions (compared with the end of training sessions). Strength was not affected by RT performed with contractions to muscle failure/fatigue, free weights versus machines, unstable versus stable surfaces; fast (< 2 s) versus moderate-slow (> 2 s) contractions, morning versus evening training sessions, short (< 1 min) versus long (> 1 min) between-set rest intervals, eccentric versus concentric contractions, power training techniques, or different set structures (cluster and complex).

Hypertrophy

The impact of RT versus CTRL on muscle hypertrophy is summarized in Table 3, and the impact of distinct RTx variables is summarized in Table 4. The results for each review are reported in Supplemental Appendix 12, Supplemental Digital Content, <https://links.lww.com/MSS/D323>. Compared with CTRL, muscle hypertrophy was positively affected by standard RT (17,24,30,32,33,37,38,40,42–47,50,127), circuit RT (57,58), and elastic band RT (60,62). There were insufficient data

TABLE 3. Resistance training forms that impact strength, hypertrophy and power when compared with CTRL.

	Strength	Hypertrophy	Power
Standard RT vs CTRL	✓ ↑ 26 reviews; <i>n</i> = 23,204 QoE = 73% ? 3 reviews; <i>n</i> = 138 QoE = 50% ↓ 1 review; <i>n</i> = 488 QoE = 50%	✓ ↑ 12 reviews; <i>n</i> = 14,924 QoE = 79% ? 3 reviews; <i>n</i> = 437 QoE = 83% ↓ 2 reviews; <i>n</i> = 775 QoE = 63%	✓ ↑ 4 reviews; <i>n</i> = 1,001 QoE = 63% ? 1 review; <i>n</i> = 152 QoE = 25%
Circuit RT vs CTRL	✓ ↑ 3 reviews; <i>n</i> = 843 QoE = 92%	✓ ↑ 1 review; <i>n</i> = 236 QoE = 100% ? 1 review; <i>n</i> = 190 QoE = 100%	? ND
Elastic band RT vs CTRL	✓ ↑ 2 reviews; <i>n</i> = 1,921 QoE = 63% ? 1 review; <i>n</i> = 51 QoE = 75%	✓ ↑ 1 review; <i>n</i> = 236 QoE = 50% ? 1 review; <i>n</i> = 133 QoE = 75%	? ND
Flywheel RT vs CTRL	? ? 1 review; <i>n</i> = 140 QoE = 50%	? ? 1 review; <i>n</i> = 104 QoE = 50%	? ND
Home-based RT vs CTRL	✓ ↑ 2 reviews; <i>n</i> = 892 QoE = 50%	? ND	? ? 1 review; <i>n</i> = 88 QoE = 75%
Nordic hamstring RT vs CTRL	? ↑ 1 review; <i>n</i> = 112 QoE = 100%	? ND	? ND
Olympic-style weightlifting vs CTRL olympic-style	? ? 1 review; <i>n</i> = 44 QoE = 75%	? ND	? ? 1 review; <i>n</i> = 112 QoE = 75%
Unilateral RT, cross-education vs CTRL	✓ ↑ 2 reviews; <i>n</i> = 1,194 QoE = 88%	? ND	? ND
Velocity-based RT vs CTRL	✓ ↑ 2 reviews; <i>n</i> = 870 QoE = 50%	? ND	? ND
Unstable surface RT vs CTRL	? ↑ 1 review; <i>n</i> = 172 QoE = 75%	? ND	? ND

✓: positively impacts outcome; ? : cannot determine impact on outcome; ✖: does not impact outcome. Within each box, the small symbol and text represent details of reviews for each respective decision, and the enlarged symbol represents the overall decision. The QoE is the quotient of the average QoE and the highest (best) possible quality of evidence, expressed as a percentage. The number of participants was calculated from the most common measurement in reviews reporting outcome-specific sample sizes.
CTRL, nonexercising control; ND, no data; QoE, quality of evidence.

to determine if eccentric flywheel RT (67) impacted muscle hypertrophy compared with CTRL. In comparisons between distinct RTx, muscle hypertrophy was positively affected by contraction type (24,104,128,129) and volume (24,91,128,130–133), but muscle hypertrophy was not affected by frequency (24,73,74,128,131,134,135), load (24,78–81,83,84,128,131,136), absolute fatigue/failure (24,98,99,128,137), blood flow restriction (110–112), variable loading (138), time under tension (24,131,139,140), power training (105), periodization (24,117,131,141), or exercise order (24,94,96,128,131). There were insufficient data to determine if muscle hypertrophy was influenced by eccentric flywheel versus standard RT (85), machine versus free-weight RT (100), single-joint versus multi-joint RT (128,142), time of day (24,102,131), inter-set rest (24,131,143,144), range of motion (86,87,128,145), set structure (24,107,108,146), intrasession autoregulation (113), varied exercise selection (115,128), or concurrent training (119,120,122,124,147).

Improving hypertrophy. Compared with CTRL, hypertrophy was improved by RT, including circuit RT and elastic band RT (Table 6). Between RTx, hypertrophy was enhanced by RT performed with eccentric-only contractions (versus concentric-only) and higher volume (≥10 sets/muscle group/wk). RT did not influence hypertrophy performed with low (1 d/wk) versus high (>5 d/wk) frequency when total volume was equated; low (30% 1RM) to high (100% 1RM) loads; contractions to muscle failure/fatigue; blood flow restriction; varying loads during repetitions; fast (0.5 s) versus slow (8 s) repetitions; power training techniques; periodization (linear, undulating, and nonperiodized); or exercise order within training sessions.

Power

The impact of RT versus CTRL on power is summarized in Table 3, and the impact of distinct RTx variables is summarized in Table 4. The results for each review are

TABLE 4. Resistance training prescription variables that impact strength, hypertrophy, and power when compared with standard (nonspecialized) RT.

	Strength	Hypertrophy	Power
Frequency	✓ ↑ 4 reviews; <i>n</i> = 3,531 QoE = 69% ↓ 3 reviews; <i>n</i> = 608 QoE = 75%	✗ ↑ 1 review; <i>n</i> = 200 QoE = 50% ? 2 reviews; <i>n</i> = 166 QoE = 38% ↓ 4 reviews; <i>n</i> = 3,177 QoE = 63%	? ND
Intensity: Load	✓ ↑ 6 reviews; <i>n</i> = 6,574 QoE = 79% ? 1 review; <i>n</i> = 151 QoE = 100% ↓ 2 reviews; <i>n</i> = 734 QoE = 63%	✗ ↑ 1 review; <i>n</i> = 231 QoE = 75% ? 1 review; <i>n</i> = 108 QoE = 75% ↓ 8 reviews; <i>n</i> = 5,340 QoE = 66%	✓ ↑ 2 reviews; <i>n</i> = 1,980 QoE = 75% ? 1 review; <i>n</i> = 151 QoE = 100%
Intensity: Fatigue/failure	✗ ↑ 1 review; <i>n</i> = 199 QoE = 75% ↓ 3 reviews; <i>n</i> = 1,371 QoE = 83%	✗ ↑ 1 review; <i>n</i> = 189 QoE = 100% ↓ 4 reviews; <i>n</i> = 800 QoE = 69%	? ? 1 review; <i>n</i> = 150 QoE = 100%
Type: Blood flow restriction	? ↑ 1 review; <i>n</i> = 460 QoE = 75% ? 2 reviews; <i>n</i> = 364 QoE = 88%	✗ ? 1 review; <i>n</i> = 72 QoE = 100% ↓ 2 reviews; <i>n</i> = 587 QoE = 75%	? ND
Type: Eccentric flywheel	✓ ↑ 1 review; <i>n</i> = 332 QoE = 75%	? ↑ 1 review; <i>n</i> = 160 QoE = 75%	✓ ↑ 1 review; <i>n</i> = 235 QoE = 75%
Type: Machine vs free-weight	✗ ↓ 1 review; <i>n</i> = 683 QoE = 75%	? ? 1 review; <i>n</i> = 123 QoE = 75%	? ND
Type: Single-joint vs multi-joint	? ND	? ? 1 review; <i>n</i> = 65 QoE = 25% ↓ 1 review; <i>n</i> = 193 QoE = 100%	? ND
Type: Unstable vs stable surface	✗ ↓ 1 review; <i>n</i> = 438 QoE = 75%	? ND	? ? 1 review; <i>n</i> = 145 QoE = 75%
Type: Variable load	? ND	✗ ↓ 1 review; <i>n</i> = 408 QoE = 50%	? ND
Time: Time-under-tension	✗ ? 1 review; <i>n</i> = 140 QoE = 75% ↓ 1 review; <i>n</i> = 509 QoE = 75%	✗ ? 1 review; <i>n</i> = 107 QoE = 50% ↓ 3 reviews; <i>n</i> = 806 QoE = 58%	? ND
Time: Time of day	✗ ↓ 2 reviews; <i>n</i> = 430 QoE = 75%	? ? 3 reviews; <i>n</i> = 336 QoE = 75%	? ND
Inter-set rest	✗ ↓ 2 reviews; <i>n</i> = 982 QoE = 63%	? ? 4 reviews; <i>n</i> = 265 QoE = 44%	? ND
Technique: Contraction type	✗ ? 1 review; <i>n</i> = 38 QoE = 75% ↓ 1 review; <i>n</i> = 1,051 QoE = 75%	✓ ↑ 1 review; <i>n</i> = 868 QoE = 75% ? 2 reviews; <i>n</i> = 92 QoE = 50% ↓ 1 review; <i>n</i> = 356 QoE = 75%	? ND
Technique: Range of motion	✓ ↑ 2 reviews; <i>n</i> = 1,262 QoE = 50%	? ↑ 1 review; <i>n</i> = 90 QoE = 75% ? 3 reviews; <i>n</i> = 276 QoE = 33%	? ND
Technique: Power training	✗ ? 1 review; <i>n</i> = 140 QoE = 75% ↓ 2 reviews; <i>n</i> = 670 QoE = 75%	✗ ↓ 1 review; <i>n</i> = 336 QoE = 75%	✓ ↑ 2 reviews; <i>n</i> = 863 QoE = 75% ? 1 review; <i>n</i> = 99 QoE = 75%

(Continued)

TABLE 4. Continued

	Strength	Hypertrophy	Power
Volume: Sets	✓ ↑ 7 reviews; n = 5,633 QoE = 71%	✓ ↑ 5 reviews; n = 2,267 QoE = 50% ? 1 review; n = 181 QoE = 75% ↓ 1 review; n = 555 QoE = 75%	✓ ↑ 1 review; n = 454 QoE = 75%
Set structure	✗ ? 1 review; n = 199 QoE = 75% ↓ 3 reviews; n = 1,837 QoE = 83%	? ? 3 reviews; n = 401 QoE = 83% ↓ 1 review; n = 189 QoE = 75%	✗ ? 1 review; n = 142 QoE = 100% ↓ 1 review; n = 445 QoE = 75%
Progression: Intrasession autoregulation	? ↑ 1 review; n = 308 QoE = 75% ↓ 1 review; n = 356 QoE = 25%	? ? 1 review; n = 243 QoE = 75%	? ND
Progression: Varied exercise selection	? ? 1 review; n = 198 QoE = 50%	? ? 1 review; n = 95 QoE = 25% ↓ 1 review; n = 218 QoE = 50%	? ND
Progression: Periodization	? ↑ 1 review; n = 616 QoE = 75% ? 1 review; n = NR QoE = 50% ↓ 2 reviews; n = 1,075 QoE = 75%	✗ ↓ 4 reviews; n = 1,414 QoE = 63%	? ND
Exercise order	✓ ↑ 4 reviews; n = 941 QoE = 88%	✗ ? 1 review; n = 71 QoE = 25% ↓ 4 reviews; n = 759 QoE = 81%	? ND
Concurrent training	? ↑ 2 reviews; n = 725 QoE = 75% ? 1 review; n = 119 QoE = 25% ↓ 3 reviews; n = 354 QoE = 67%	? ↑ 1 review; n = NR QoE = 50% ? 3 reviews; n = 351 QoE = 67% ↓ 1 review; n = NR QoE = 25%	? ↑ 2 reviews; n = NR QoE = 50% ? 1 review; n = 18 QoE = 75%

✓: positively impacts outcome; ? : cannot determine impact on outcome; ✗: does not impact outcome. Within each box, the small symbol and text give the details of reviews for each respective decision, and the enlarged symbol represents the overall decision. The QoE is the quotient of the average QoE and the highest (best) possible quality of evidence, expressed as a percentage. The number of participants was calculated from the most common measurement in reviews reporting outcome-specific sample sizes. CTRL, nonexercising control; ND, no data; QoE, quality of evidence.

reported in Supplemental Appendix 13, Supplemental Digital Content, <https://links.lww.com/MSS/D323>. Compared with CTRL, power was enhanced by standard RT (41,48–50,148). There were insufficient data to determine whether home-based RT (63) and Olympic-style weightlifting (69) impacted power compared with CTRL. In comparisons between distinct RTx, power was affected by load (77,149,150), eccentric flywheel training versus standard RT (85), Olympic-style weightlifting versus standard RT (69), power training versus standard RT (82,105,148), and volume (148), but power was not affected by set structure (107,108). There were insufficient data to determine if power was affected by fatigue/failure (99), unstable surfaces (70), or concurrent training (123,124,151).

Improving power. RT improved power compared with CTRL (Table 6). Between RTx and power training,

power was enhanced by RT performed with moderate loads (30%–70% 1RM), an eccentric flywheel device, Olympic-style weightlifting, and power training techniques, versus standard RT, and low-to-moderate volume (repetitions-set < 24). Power was not affected by RT performed with different set structures (cluster or rest redistribution).

Muscular Endurance

The impact of RT versus CTRL and distinct RTx variables on muscular endurance is summarized in Supplemental Appendix 9, Supplemental Digital Content, <https://links.lww.com/MSS/D323>. The results for each review are reported in Supplemental Appendix 14, Supplemental Digital Content, <https://links.lww.com/MSS/D323>. Compared with CTRL, muscular endurance was impacted by standard RT (41,54) and home-based

RT (63). There were insufficient data to determine whether unstable surface RT (70) or velocity-based RT (66) impacted muscular endurance compared with CTRL. There were insufficient data to determine if muscular endurance was affected by load (82), unstable surfaces (70), power training (82), volume (82), or set structure (107,108).

Improving muscular endurance. RT improved muscular endurance compared with CTRL.

Gait Speed

The impact of RT versus CTRL and distinct RTx variables on gait speed is summarized in Table 5. The results for each review are reported in Supplemental Appendix 15, Supplemental Digital Content, <https://links.lww.com/MSS/D323>. Compared with CTRL, gait speed was

impacted by standard RT (17,30,32,33,36–38,47,53,152–155). There were insufficient data to determine whether elastic band RT (62) or power training (156) impacted gait speed compared with CTRL. In comparisons between distinct RTx, there were insufficient data to determine if gait speed was influenced by load (82) or power training versus standard RT (82).

Improving gait speed. RT improved gait speed compared with CTRL.

Timed Up-and-Go

The impacts of RT versus CTRL and distinct RTx variables on timed up-and-go (TUG) performance are summarized in Table 5. The results for each review are reported in Supplemental Appendix 16, Supplemental Digital Content, <https://links.lww.com/MSS/D323>.

TABLE 5. Resistance training forms and prescription variables that impact physical function outcomes.

	Stair Climbing	Gait Speed	Balance	Chair Stand Performance	Timed Up-and-Go
Standard RT vs CTRL	? ND	✓ ↑ 8 reviews; <i>n</i> = 3,407 QoE = 81% ? 3 reviews; <i>n</i> = 115 QoE = 50% ↓ 2 reviews; <i>n</i> = 1,660 QoE = 100%	✓ ↑ 1 review; <i>n</i> = 406 QoE = 75% ? 2 reviews; <i>n</i> = 227 QoE = 63%	✓ ↑ 2 reviews; <i>n</i> = 954 QoE = 88% ? 3 reviews; <i>n</i> = 291 QoE = 75% ↓ 1 review; <i>n</i> = 657 QoE = 100%	✓ ↑ 5 reviews; <i>n</i> = 1,568 QoE = 90% ? 5 reviews; <i>n</i> = 479 QoE = 70%
Elastic band RT vs CTRL	? ND	? ? 1 review; <i>n</i> = 160 QoE = 75%	? ND	? ND	? ↑ 1 review; <i>n</i> = 154 QoE = 75%
Home-based RT vs CTRL	? ND	? ND	✓ ↑ 1 review; <i>n</i> = 1,484 QoE = 75%	? ND	? ND
Power training RT vs CTRL	? ND	? ? 1 review; <i>n</i> = 165 QoE = 75%	? ND	? ND	? ND
Unstable surface RT vs CTRL	? ND	? ND	? ? 1 review; <i>n</i> = 138 QoE = 75%	? ND	? ND
Intensity: Load	? ? 1 review; <i>n</i> = 84 QoE = 75%	? ? 1 review; <i>n</i> = 38 QoE = 75%	? ND	? ? 1 review; <i>n</i> = 23 QoE = 75%	? ? 1 review; <i>n</i> = 38 QoE = 75%
Intensity: Fatigue/failure Type: Unstable vs stable surface	? ND ? ND	? ND ? ND	? ND ? ? 1 review; <i>n</i> = 142 QoE = 75%	? ND ? ND	? ND ? ND
Technique: Power training	? ? 2 reviews; <i>n</i> = 142 QoE = 88%	? ? 1 review; <i>n</i> = 79 QoE = 75%	? ND	? ? 3 reviews; <i>n</i> = 442 QoE = 83%	? ? 2 reviews; <i>n</i> = 119 QoE = 75% ↓ 1 review; <i>n</i> = 227 QoE = 100%
Concurrent training	? ND	? ND	? ? 1 review; <i>n</i> = 45 QoE = 50%	? ND	? ND

✓: positively impacts outcome; ?; cannot determine impact on outcome; ✖: does not impact outcome. Within each box, the small symbol and text represent details of reviews for each respective decision, and the enlarged symbol represents the overall decision. The QoE is the quotient of the average QoE and the highest (best) possible quality of evidence, expressed as a percentage. The number of participants was calculated from the most common measurement in reviews reporting outcome-specific sample sizes.

CTRL, nonexercising control; ND, no data; QoE, quality of evidence.

TABLE 6. Resistance training prescriptions to improve muscle function and hypertrophy.

Outcome	RT vs CTRL	RTx to Enhance Adaptation
Strength	Strength is improved by RT, including circuit RT, elastic band RT, home-based RT, and velocity-based RT.	Frequency: ≥ 2 sessions/wk Intensity: $\geq 80\%$ 1RM (dose-response) Type: Eccentric flywheel RT Technique: Full range of motion Volume: 2–3 sets/session Exercise order: Beginning of training session
Hypertrophy	Hypertrophy is improved by RT, including circuit RT and elastic band RT.	Type: Eccentric contractions/overload Volume: ≥ 10 sets/wk (dose-response)
Power	Power is improved by RT.	Intensity: Loads = 30%–70% 1RM Type: Eccentric flywheel RT Technique: Olympic-style weightlifting; Power RT Volume: Low-moderate (repetitions \cdot sets ≤ 24)
Muscular endurance	Muscular endurance is improved by RT.	ND
Gait speed	Gait speed is improved by RT.	ND
Timed up-and-go	Timed up-and-go is improved by RT.	ND
Chair stand test	Chair stand test performance is improved by RT.	ND
Balance	Balance is improved by RT.	ND
Stair climbing	ND	ND
Multicomponent function	Multicomponent function is improved by RT, including elastic band RT and home-based RT.	Technique: Power RT
SPPB	SPPB is not improved by RT.	Technique: Power RT
Walking performance	ND	Technique: Power RT
Running performance	ND	Type: Velocity-based RT
Jumping performance	Jumping performance is improved by flywheel RT and velocity-based RT.	Type: Velocity-based RT
Contraction velocity	Contraction velocity is improved by RT.	ND
Change of direction (agility)	ND	ND
Functional reach	ND	ND

^aEnhanced compared with standard RT and based on included meta-analyses showing a significant effect.

1RM, one-repetition maximum; CTRL, no exercise; ND, insufficient data to form conclusion; RT, resistance training; RTx, resistance training prescription; SPPB, Short Physical Performance Battery.

Compared with CTRL, TUG was impacted by standard RT (30,32,33,36,38,47,50,152,154,156). There were insufficient data to determine if elastic band RT (62) impacted TUG compared with CTRL. In comparisons between distinct RTx, there were insufficient data to determine if TUG was affected by load (82) or power training versus standard RT (82,106,157).

Improving timed up-and-go. RT improved TUG compared with CTRL.

Chair Stand Test

The impacts of RT versus CTRL and distinct RTx variables on chair stand test performance are summarized in Table 5. The results for each review are reported in Supplemental Appendix 17, Supplemental Digital Content, <https://links.lww.com/MSS/D323>. Compared with CTRL, chair stand performance was impacted by standard RT (30,32,33,36,50,152). In comparisons between distinct RTx, there were insufficient data to determine if chair stand performance was affected by load (82) or power training versus standard RT (82,106,157).

Improving chair stand test. RT improved chair stand performance compared with CTRL.

Balance

The impacts of RT versus CTRL and distinct RTx variables on balance are summarized in Table 5. The results for each review are reported in Supplemental Appendix 18, Supplemental Digital Content, <https://links.lww.com/MSS/D323>. Compared with CTRL, balance was impacted by standard RT (52,54,154) and home-based RT (63). There were insufficient data to determine whether unstable surfaces (70) impacted balance compared with CTRL. In comparisons between distinct RTx, there were insufficient data to determine if unstable surfaces (70) or concurrent training (158) affected balance.

Improving balance. RT improved balance compared with CTRL.

Stair Climbing

The impacts of RT versus CTRL and distinct RTx variables on stair climbing are summarized in Table 5.

The results for each review are reported in Supplemental Appendix 19, Supplemental Digital Content, <https://links.lww.com/MSS/D323>. In comparisons between distinct RTx, there were insufficient data to determine if stair climbing was affected by load (82) or power training versus standard RT (82,157).

Improving stair climbing. There were insufficient data to determine the impact of RT on stair climbing performance.

Multicomponent Function

The impacts of RT versus CTRL and distinct RTx variables on multicomponent function—the aggregate performance on several physical function assessments—are summarized in Supplemental Appendix 9, Supplemental Digital Content, <https://links.lww.com/MSS/D323>. The results for each review are reported in Supplemental Appendix 20, Supplemental Digital Content, <https://links.lww.com/MSS/D323>. Compared with CTRL, multicomponent function was impacted by standard RT (17,24,42,45,159), elastic band RT (60,61), and home-based RT (64). In comparisons between distinct RTx, multicomponent function was affected by elastic band RT (61) and power training (105,157,160), but was not affected by time under tension (24) or volume (91). There were insufficient data to determine if the multicomponent function was positively affected by load (24), range of motion (87), or concurrent training (119).

Improving multicomponent function. Compared with CTRL, multicomponent function was improved by RT, including elastic band RT and home-based RT (Table 6). Comparing between RTx, multicomponent function was enhanced by RT performed with power training techniques, and standard RT enhanced multicomponent function compared with elastic band RT.

Short Physical Performance Battery

The impacts of RT versus CTRL and distinct RTx variables on the Short Physical Performance Battery (SPPB) are summarized in Supplemental Appendix 10, Supplemental Digital Content, <https://links.lww.com/MSS/D323>. The results for each review are reported in Supplemental Appendix 21, Supplemental Digital Content, <https://links.lww.com/MSS/D323>. Compared with CTRL, SPPB was not affected by standard RT (30). There were insufficient data to determine whether power training (156) had an impact on SPPB compared with CTRL. In comparing different RTx, SPPB was positively affected by power training (106,157).

Improving short physical performance battery. RT did not improve SPPB compared with CTRL. Comparing RTx, SPPB was enhanced by RT performed with power training.

Walking Performance

The impacts of RT versus CTRL and distinct RTx variables on walking test performance (e.g., 6-min walk test) are summarized in Supplemental Appendix 10, Supplemental Digital Content, <https://links.lww.com/MSS/D323>. The results for each review are reported in Supplemental Appendix 22, Supplemental Digital Content, <https://links.lww.com/MSS/D323>. Compared with CTRL, there were insufficient data to determine if standard RT (30) or elastic band RT (62) impacted walking performance compared with CTRL. In comparisons between distinct RTx, walking performance was positively affected by power training (106,157).

Improving walking performance. Between RTx, walking performance was enhanced only by RT performed with power training techniques.

Running Performance

The impacts of RT versus CTRL and distinct RTx variables on running performance are summarized in Supplemental Appendix 9, Supplemental Digital Content, <https://links.lww.com/MSS/D323>. The results for each review are reported in Supplemental Appendix 23, Supplemental Digital Content, <https://links.lww.com/MSS/D323>. Compared with CTRL, there were insufficient data to determine whether standard RT (48,69,161), eccentric flywheel RT (162), and velocity-based RT (66) impacted running performance compared with CTRL. In comparisons between distinct RTx, running performance was positively affected only by velocity-based RT versus standard RT (65). There were insufficient data to determine if running performance was affected by eccentric flywheel RT versus standard RT (85), Olympic-style weightlifting versus standard RT (69), set structure (109), or concurrent training (123).

Improving running performance. Comparing RTx, running performance was enhanced only by velocity-based RT.

Jumping Performance

The impacts of RT versus CTRL and distinct RTx variables on jumping performance are summarized in Supplemental Appendix 9, Supplemental Digital Content, <https://links.lww.com/MSS/D323>. The results for each review are reported in Supplemental Appendix 24, Supplemental Digital Content, <https://links.lww.com/MSS/D323>. Compared with CTRL, jumping performance was impacted by eccentric flywheel RT (162) and velocity-based RT (66). In comparisons between distinct RTx, jumping performance was positively affected by velocity-based RT versus standard RT (65), but jumping performance was not affected by set structure (108,109). There were insufficient data to determine if jumping performance was affected by load (81), eccentric flywheel RT versus standard RT (85), or machine versus free-weight

RT (100). We note that jumping performance should not be considered as a proxy for muscle power.

Improving jumping performance. Compared with CTRL, jumping performance was improved by eccentric flywheel RT and velocity-based RT (Table 6). Comparing RTx, velocity-based RT enhanced jumping performance; however, jumping performance was not affected by RT performed with different set structures (complex vs cluster).

Contraction Velocity

The impact of RT versus CTRL and distinct RTx variables on contraction velocity is summarized in Supplemental Appendix 10, Supplemental Digital Content, <https://links.lww.com/MSS/D323>. The results for each review are reported in Supplemental Appendix 25, Supplemental Digital Content, <https://links.lww.com/MSS/D323>. Compared with CTRL, contraction velocity was positively impacted by standard RT (163). In comparisons between distinct RTx, there were insufficient data to determine if contraction velocity was affected by cluster or rest-redistribution set structures (107,108).

Improving contraction velocity. RT improved contraction velocity compared with CTRL.

Change of Direction (Agility)

The impacts of RT versus CTRL and distinct RTx variables on the ability to change direction are summarized in Supplemental Appendix 9, Supplemental Digital Content, <https://links.lww.com/MSS/D323>. The results for each review are reported in Supplemental Appendix 26, Supplemental Digital Content, <https://links.lww.com/MSS/D323>. Compared with CTRL, there were insufficient data to determine whether standard RT (69) or eccentric flywheel RT (162) impacted the ability to change direction. In comparisons between distinct RTx, there were insufficient data to determine if agility was affected by Olympic-style weightlifting (69).

Improving change of direction (agility). There were insufficient data to determine the impact of RT on agility.

Functional Reach Test

The impact of RT versus CTRL on functional reach test performance is summarized in Supplemental Appendix 10, Supplemental Digital Content, <https://links.lww.com/MSS/D323>. The results for each review are reported in Supplemental Appendix 27, Supplemental Digital Content, <https://links.lww.com/MSS/D323>. Compared with CTRL, there were insufficient data to determine whether standard RT (154) impacted functional reach performance.

Improving functional reach test. There were insufficient data to determine the impact of RT on functional reach test performance.

DISCUSSION

Resistance training is a central component of exercise programs. It should be a core component of physical fitness programming, as it has broad-reaching benefits for muscular health and physical function. This overview of reviews summarized 137 systematic reviews to determine the impact of RTx variables on muscle function and hypertrophy in healthy adults. Compared with no exercise, RT improves muscle strength, hypertrophy, power, endurance, contraction velocity, and performance on several physical function tests (balance, gait speed, chair stand, and timed up-and-go; Table 3). Additionally, other forms of RT improve multiple outcomes compared with CTRL, including home-based RT, elastic band RT, power RT (which involves performing the concentric phase quickly), velocity-based RT, and circuit RT (Table 3). Based on these results, it is apparent that many forms of RT work to promote the primary hallmark outcomes of RT: increased strength, hypertrophy, and power. Our primary recommendation is that healthy adults perform RT with high effort (effort can be measured through various scales, but can be achieved with various loads and sets per the FITT-VP principle) at least twice weekly, with all major muscle groups being engaged.

Between distinct RT programs, only some RTx variables can be altered to optimize increases in muscle function and hypertrophy (Tables 4 and 6). Significant and optimal improvements are two different goals; however, participants and facilitators of RT must distinguish between them. Compared with CTRL, significant improvements in muscle function, hypertrophy, and physical performance can be accomplished with the adoption of many RT programs. This overview of reviews enhances the quality of recommendations for healthy adults to significantly increase muscle function, hypertrophy, and physical performance by providing current, comprehensive, and evidence-based insights into the impact of RTx variables.

Progressive overload is a concept considered fundamental to RT programming principles. Progressive overload refers to the need to increase the stimulus (stressor) placed upon the muscle throughout a training program. Progressive overload is often proposed as essential for continued adaptive progress with any form of exercise training as physiological systems adapt to reduce the stress of the exercise and build an increasing adaptive response (i.e., strength). Such adaptation would require increasing the stimulus, such as load, volume, training frequency, exercise selection, or duration (even if only slightly), as the muscle continues to adapt to produce further adaptations (164). Modifying an RTx (e.g., increasing relative load [percentage of one-repetition maximum, %1RM]) can support progressive overload. Still, the same relative load can also be sustained when regular strength testing is performed or perceived exertion scales are used

to increase absolute load commensurate with strength gains (113,165). Notably, as the results in Table 4 highlight, progression is not necessary to achieve beneficial outcomes, and overload, or more accurately, increasing the stimulus in some manner, is likely a requirement only for those seeking continued longer term progress. We note that in some populations, due to inexperience (and potential safety concerns), RTx necessitates that loads are necessarily low and progression is a requirement to achieve a meaningful benefit. Continued progression could be a personal decision and part of the individualization of RTx (see below). A similar commentary could be made around the “need” for variation of RTx variables (see below).

The principle of specificity states that training adaptations are specific to the training stimulus applied. Some adaptations can be affected by modifying RTx variables, although there is a considerable carryover of training effects on general muscular performance in various domains in nonadvanced trainees (17,24). Individualization involves modifying RT programs to meet the unique goals, needs, and characteristics of each individual, such as their experience and performance level. Individualized programs can increase exercise adoption and adherence (10), but individualization has been scarcely discussed in previous Position Stands (11,12). With the accumulation of evidence demonstrating that many forms of RT are effective for healthy adults to improve muscle function and health, RT programs would ideally be individualized to maximize adherence, enjoyment, safety, and effectiveness specific to training goals.

Variation is the systematic modification of RTx variables over time to facilitate continued adaptation. One of the most common forms of variation is periodization—intentionally modifying prescription variables (e.g., load, volume, frequency) throughout an RT program. The impact of periodization on strength could not be determined, although one review found that periodized programs were slightly favored over nonperiodized programs to maximize strength gains under volume-equated conditions (117). With appropriate progressive overload, periodization is not significantly superior to nonperiodized programs; thus, periodization is less important than previously hypothesized (11,12) for healthy adults to improve muscle function and hypertrophy. We note, however, that definitions of periodization vary (166,167), but broadly involve the systematic manipulation of RTx variables with the goal of optimizing performance adaptations, managing fatigue, preventing overtraining/injury, and peaking for specific goals. Thus, it may be that the ordering of variables within a given RTx to achieve certain goals is understudied compared with the outcomes contained in the reviews we examined. We also note that periodization and programming are often conflated (168), which may have been the case in some reviews.

Resistance training improves numerous measures of muscle function and can result in hypertrophy compared with no exercise (Fig. 2). Provided the strikingly low RT participation levels (169–173), individuals and practitioners alike should acknowledge the tremendous benefits of completing RT of various forms compared with no exercise. Additionally, nontraditional forms of RT also yield marked benefits—for example, elastic band RT has been shown to increase strength, hypertrophy, and certain components of physical function. Similarly, home-based RT improved strength, muscular endurance, and balance. Nontraditional forms of RT may provide alternative, perhaps more accessible or approachable, strategies for completing RT with appreciable physical benefits.

We build on several aspects of the 2009 ACSM Position Stand (11), which aimed to provide evidence-based RTx guidelines for healthy adults wishing to progress beyond the first 3–4 months of training. Previous guidelines have recommended healthy adults use free weights and machines to complete two to three RT sessions per week, with eight to ten exercises involving major muscle groups per session, one to four sets per exercise, eight to twenty repetitions per set, 2–3 min rest between sets, loads 40%–70% 1RM, and follow the principles of progressive overload, specificity, and variation (10,11). Current estimates (from self-report) are that only ~30% of American adults complete some muscle-strengthening activities at least 2 d/wk, and nearly 60% complete no muscle-strengthening exercise (169–173). Estimates for older persons vary; however, data from the UK, United States, and Australia put participation rates from as low as 1% to as high as 40% (174). We suspect that rates of participation among older persons for the types of RT programs we outline here are likely 10%–15% (175). Notably, others have estimated that developing general muscular fitness (strength, hypertrophy, and power) according to previous ACSM guidelines may, in some cases, require training for 20 h or more per week (176). Thus, the recommendations of the previous Position Stand (10,11) may be less relevant for most adults, particularly older adults, many of whom do not engage in RT. The specific program details outlined in previous Position Stands may be an appropriate starting point for some individuals; however, several other RT programs, as evidenced here, can also be effective.

Deviating from previous guidelines, we propose that individualizing programs to increase RT participation is, from our perspective, more important than conforming to specific RTx criteria outlined in previous Position Stands. Research on RTx reveals that “minimal doses” of RT are able to bring about substantial strength, hypertrophy, and physical functional gains (177). We propose that our stance around encouraging participation in RT is underscored by the expanding knowledge of health benefits, some of which are unique to RT (4–6). The recommendations presented here apply to healthy adults,




FIGURE 2—Schematic representation of the modes of RT and the outcomes that are positively influenced by engaging in regular RT.

ranging from complete novices (with no experience with RT) to experienced trainees, and support the adoption of a much broader range of RT programs to improve overall muscular strength, function, and hypertrophy. Advances enable a more comprehensive understanding of how RTx variable manipulation affects outcomes. For example, the previous Position Stand recommended novice lifters complete two to three full-body sessions per week for muscle hypertrophy, and this recommendation was graded as the highest possible quality of evidence (11). However, this recommendation was informed by only three studies (178–180) ($n = 59$ total participants), and two of the studies did not investigate frequency (178,179). In contrast, we present guidelines suitable for many healthy adults, supported by ample primary evidence, that were formulated with leading evidence synthesis methodologies.

Resistance training is safe for healthy adults of all ages (5), but many adults, particularly older adults, avoid RT due to misperceptions about safety and injury risks (181). In an analysis of >38,000 participants (>6700 RT participants), of which >11,000 were older adults, exercise did not increase the risk of serious adverse events. The risk of nonserious adverse events (e.g., pain, fatigue, bursitis, and edema) was not different than aerobic exercise in terms of injury rate or risk (182). Nonfatal cardiovascular complications also occur much less frequently during RT than aerobic training (5,183). In 23 studies ($n = 1174$) of adults with coronary heart disease, all 63 nonfatal

cardiovascular complications occurred during aerobic training, and the 20 musculoskeletal complications that occurred during RT were caused by preexisting conditions (knee arthritis) and were resolved by changing RT intensity or body position (183). Clearly, RT can be safe and effective for healthy adults of all ages.

This overview of reviews summarized RTx variable-level evidence, an approach that prevents statistical insights into the comparative effectiveness (e.g., dose-response) of individual RT variables. Nuanced details of RT protocols and dose-response data require discussion for practitioners designing RT programs to enhance adaptations, since some features of RTx are not reflected in variable-level summaries. For example, RT frequency was found to impact strength; however, there was insufficient evidence to conclude a dose-response relationship, and the impact is diminished when volume is equated (73,76,90).

Load (often used synonymously with intensity) has repeatedly been shown to impact strength more than other RT adaptations (17,83,184). Load has typically been quantified as a %1RM or repetition maximum. Several methods have been used to convert loads between %1RM and repetition maximum (17,184,185), including a recent analysis that considered exercise-specific conversions and between-subject variability (186). The number of repetitions performed within a given set is inherently and inversely related to load, provided a given load and near-fatiguing effort are applied; however, not all

adaptations are enhanced by fatiguing efforts (e.g., power and strength). Practitioners should consider the principle of specificity when designing RT programs to achieve specific outcomes.

Increasing the number of sets (volume) per exercise positively impacted strength (88–93) and hypertrophy (24,128,131–133), which aligns with a recent analysis by Swinton et al. (184). Clearly, one set is superior to zero sets (CTRL), and two sets are superior to one set (17,24,88–93,128,131–133), but the exact number of sets required to optimize adaptations cannot be ascertained. A meta-regression (187) of the effect of volume (weekly) sets showed progression, but as expected, a dose-response that plateaued and showed diminishing returns beyond ~2–3 sets/exercise for strength and ~18–20 weekly sets for hypertrophy with varying loads; thus, healthy adults are advised to complete at least two sets per exercise. Completing more than two sets per exercise may provide additional benefits, but we hypothesize that these benefits diminish with each subsequent set. Weekly volume load—the total amount of load lifted in each week of training, defined by load/rep • reps/set • sets/exercise • exercises/session • sessions/week—which would presumably represent the complete representation of “dose” of RT—was rarely considered in included reviews.

Completing sets to fatigue (momentary muscular failure) does not enhance gains in strength, hypertrophy, and power, and so is not necessary for benefits to occur. It may also be that lifting to fatigue is inadvisable for certain populations (e.g., older individuals) due to risks to vascular health and an increased risk of injury resulting from poor form (24,97–99). We propose that an adequate stimulus (the effort to meet the minimum stressor) is required to induce adaptations. Sufficient effort (assessed using various scales) can be accomplished by completing sets with various RTx and completion of “near-failure” or a target of 2–3 repetitions in reserve (RIR) (24,97–99,128,137,188–190).

Blood flow restriction protocols often compare different loads (e.g., higher load RT versus BFR with lower load), so the individual effect of BFR cannot be distinguished from the load in many cases. Similarly, flywheel RT typically involves eccentric overload, so the individual effect of flywheel RT and contraction type cannot be distinguished. Evidence-based conclusions on these matters cannot be determined with the available data, but consideration is warranted when designing RT programs and conducting future research. Notably, these considerations can be largely overlooked when designing RT programs for most adults who are inexperienced with RT. Untrained individuals will benefit from various RT programs, provided that progression, the variables outlined here as affecting outcomes, and adherence are core principles.

Limitations should be considered when interpreting the results presented in this analysis. Overviews of reviews

provide strong (larger or consistent effects) pooled, group-level evidence, so that appropriate deviations from the reported recommendations will be required on various occasions, in accordance with the principle of individualization. We acknowledge that any RT program would need to be tailored to an individual’s needs and goals. Not discussed here, but of paramount importance, is the understanding and acknowledgment of the critical role adherence plays in any RT program. Such considerations are beyond the scope of this review, but work has been done to examine this important area (191,192).

The totality of evidence in overviews of reviews is limited to interventions and outcomes summarized by systematic reviews. For example, the Short Physical Performance Battery (SPPB) is a common assessment used in randomized trials; however, only two eligible reviews have synthesized the results of the SPPB. When evaluating the results of this review, it would lead one to conclude that RT does not improve SPPB performance; however, this conclusion likely reflects the sparseness of evidence, rather than a true null effect. We note that there was ample evidence for individual components of the SPPB—gait speed, balance, and chair stand tests—and RT improved all these components. Overviews of reviews are also limited by overlapping evidence. While we estimated this overlap using the CCA method for strength, we did not do so for other variables. We acknowledge that caution is needed when examining other outcomes for which we provide no CCA analysis. Individual studies included in multiple eligible reviews could inflate the evidence supporting or refuting conclusions; for example, three reviews were included on the impact of RT time of day on hypertrophy (24,102,131), but two reviews (24,131) synthesized one original review (102). Overlapping data are an issue in overviews of reviews, and methods are being developed to improve exercise-related overviews of reviews (193,194). Overviews of reviews do not permit inferences about the comparative effectiveness of different interventions (14). This review summarizes the individual impact of RTx variables; however, alternative data synthesis methods are required to formulate evidence-based conclusions on the relative effects of multiple RTx. For example, network meta-analysis can be leveraged to statistically compare and rank the efficacy of unique RT programs (17). Reviews included in this overview of reviews were limited to healthy adults. Provided sufficient evidence, future guidelines can be developed for additional subpopulations (e.g., older adults and clinical populations). There was insufficient evidence to form recommendations on emerging RTx strategies; for example, various minimal-dose RT programs warrant further investigation and consideration when individualizing RT programs (195). The method of testing muscular endurance (e.g., absolute vs relative, pre- vs postintervention strength) was not accounted for in the data synthesis and should be considered when designing future studies (196).

Methods to quantify load based on proximity to muscular failure (e.g., RIR, perceived exertion) are appealing because they could translate to various RT forms (e.g., elastic band or body weight RT). While training to failure is not obligatory for optimizing results, there is insufficient evidence to quantify exact RIR and perceived exertion targets (189,190,197,198). The available evidence did not compare distinct body regions or muscle groups. In the view of the authors, applying the recommendations herein to the body regions “upper” and “lower” for “push” and “pull” exercises (i.e., four body regions) is sufficient to target major muscle groups. The movement directions “horizontal” and “vertical” may also be considered for upper body exercises (i.e., six body regions). The regions and movements we recommend should align with the FITT-VP principle in terms of RTx, which could be a per-training-session requirement or a per-week requirement, depending on individual goals, training status, age, and other relevant variables.

We also need to acknowledge that randomized trials in sports and exercise medicine-related fields are often poor quality with small sample sizes, poor randomization methodology, lack of preregistration, poor reporting of adverse events, and other shortcomings (199), and much remains to be done in terms of the conduct of evidence syntheses (200,201). Guidelines have been developed to improve the reporting of exercise trials, and we would encourage their adoption (202). Consequently, cohesive QoE evaluations, such as pooling Grading of Recommendations Assessment, Development and Evaluation assessments (14), were not possible. Field-specific guidelines for evidence syntheses have emerged and will continue to be developed (203). Nonetheless, to improve the quality of evidence that informs guidelines and best practices as a field, a higher, more rigorous standard of trial conduct and evidence syntheses are required in the field of RT.

This overview of reviews represents, to date, the most comprehensive summary of the impact of RTx variables on muscle function and hypertrophy in healthy adults. RT greatly enhances overall muscular health compared with no exercise, and only a few RTx variables can be manipulated to enhance adaptations in experienced trainees. Therefore, individuals should prioritize completing any form of RT to improve muscle function and hypertrophy. Exercise and healthcare practitioners can provide invaluable guidance by leveraging these conclusions to promote RT participation and adherence, helping adults of all ages improve their health and fitness.

This article is being published as an official pronouncement of the American College of Sports Medicine. This pronouncement was reviewed for the American College of Sports Medicine by members-at-large and the Pronouncements Committee. Care has been taken to confirm the accuracy of the information presented and to describe generally accepted practices. However, the authors, editors, and publisher are not responsible for errors or omissions or for any consequences from the application of the information in this

publication and make no warranty, expressed or implied, with respect to the currency, completeness, or accuracy of the contents of the publication. The application of this information in a particular situation remains the professional responsibility of the practitioner; the clinical treatments described and recommended may not be considered absolute and universal recommendations. B. S. C. was supported by an Alexander Graham Bell Canada Graduate Scholarship-Doctoral. S. M. P. thanks the Canada Research Chairs program for its support and acknowledges grant support from NSERC and CIHR. No specific source of funding was used for this work. B. J. S. formerly served on the scientific advisory board for Tonal Corporation, a manufacturer of fitness equipment. No other authors reported conflicts relevant to this work. M. A. F. S., E. S. R., B. J. S., A. E. S. R., G. A. T., N. T. T., T. A. W., T. J. W., and S. M. P. designed and conceived the review; B. S. C. undertook the search of articles; B. S. C., A. C. D., C. V. L., and J. S. extracted data; B. S. C. synthesized data; B. S. C., M. A. F. S., E. S. R., B. J. S., A. E. S. R., G. T., N. T. T., T. A. W., T. J. W., and S. M. P. reviewed evidence recommendations; B. S. C. and S. M. P. wrote manuscript; B. S. C., A. C. D., M. A. F. S., C. V. L., E. S. R., B. J. S., A. E. S. R., J. S., G. A. T., N. T. T., T. A. W., T. J. W., and S. M. P. reviewed manuscript. The results of the study are presented clearly, honestly, and without fabrication, falsification, or inappropriate data manipulation. The results of the present study do not constitute endorsement by the American College of Sports Medicine.

REFERENCES

1. Phillips SM, Ma JK, Rawson ES. The coming of age of resistance exercise as a primary form of exercise for health. *ACSM'S Health Fit J.* 2023;27(6):19–25.
2. World Health Organization. *WHO Guidelines on Physical Activity and Sedentary Behaviour.* World Health Organization; 2020.
3. Delorme TL. Restoration of muscle power by heavy-resistance exercises. *J Bone Joint Surg.* 1945;27(4):645–67.
4. Shailendra P, Baldock KL, Li LSK, et al. Weight training and risk of all-cause, cardiovascular disease and cancer mortality among older adults. *Int J Epidemiol.* 2024;53(3):dyae074.
5. Paluch AE, Boyer WR, Franklin BA, et al; on behalf the American Heart Association Council on Lifestyle and Cardiometabolic Health; Council on Arteriosclerosis, Thrombosis and Vascular Biology; Council on Clinical Cardiology; Council on Cardiovascular and Stroke Nursing; Council on Epidemiology and Prevention; and Council on Peripheral Vascular Disease. Resistance exercise training in individuals with and without cardiovascular disease: 2023 update: a scientific statement from the American Heart Association. *Circulation.* 2024;149(3):e217–31.
6. Momma H, Kawakami R, Honda T, Sawada SS. Muscle-strengthening activities are associated with lower risk and mortality in major non-communicable diseases: a systematic review and meta-analysis of cohort studies. *Br J Sports Med.* 2022;56(13):755–63.
7. Carneiro L, Afonso J, Ramirez-Campillo R, Murawska-Ciałowicz E, Marques A, Clemente FM. The effects of exclusively resistance training-based supervised programs in people with depression: a systematic review and meta-analysis of randomized controlled trials. *Int J Environ Res Public Health.* 2020;17(18):6715.
8. Gordon BR, McDowell CP, Hallgren M, Meyer JD, Lyons M, Herring MP. Association of efficacy of resistance exercise training with depressive symptoms: meta-analysis and meta-regression analysis of randomized clinical trials. *JAMA Psychiatry.* 2018;75(6):566–76.
9. Kovacevic A, Mavros Y, Heisz JJ, Fiararone Singh MA. The effect of resistance exercise on sleep: a systematic review of randomized controlled trials. *Sleep Med Rev.* 2018;39:52–68.

10. Garber CE, Blissmer B, Deschenes MR, et al; American College of Sports Medicine. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: guidance for prescribing exercise. *Med Sci Sports Exerc.* 2011;43(7):1334–59.
11. American College of Sports Medicine. American College of Sports Medicine position stand. Progression models in resistance training for healthy adults. *Med Sci Sports Exerc.* 2009;41(3):687–708.
12. Kraemer WJ, Adams K, Cafarelli E, et al; American College of Sports Medicine. American College of Sports Medicine position stand. Progression models in resistance training for healthy adults. *Med Sci Sports Exerc.* 2002;34(2):364–80.
13. Belbasis L, Bellou V, Ioannidis JPA. Conducting umbrella reviews. *BMJ Med.* 2022;1(1):e000071.
14. Pollock M, Fernandes RM, Becker LA, Pieper D, Hartling L. Chapter V: overviews of reviews. In: Higgins J, Chandler J, Cumpston M, Li T, Page MJ, Welch VA, editors. *Cochrane Handbook for Systematic Reviews of Interventions Version 6.4 (Updated August 2023)*; Cochrane Publishing; 2023.
15. Hartling L, Chisholm A, Thomson D, Dryden DM. A descriptive analysis of overviews of reviews published between 2000 and 2011. *PLoS One.* 2012;7(11):e49667.
16. Gates M, Gates A, Pieper D, et al. Reporting guideline for overviews of reviews of healthcare interventions: development of the PRIOR statement. *BMJ.* 2022;378:e070849.
17. Currier BS, McLeod JC, Banfield L, et al. Resistance training prescription for muscle strength and hypertrophy in healthy adults: a systematic review and Bayesian network meta-analysis. *Br J Sports Med.* 2023;57(18):1211–20.
18. Gielen E, Beckwee D, Delaere A, De Breucker S, Vandewoude M, Bautmans I; Sarcopenia Guidelines Development Group of the Belgian Society of Gerontology and Geriatrics (BSGG). Nutritional interventions to improve muscle mass, muscle strength, and physical performance in older people: an umbrella review of systematic reviews and meta-analyses. *Nutr Rev.* 2021;79(2):121–47.
19. Shea BJ, Grimshaw JM, Wells GA, et al. Development of AMSTAR: a measurement tool to assess the methodological quality of systematic reviews. *BMC Med Res Methodol.* 2007;7:10.
20. Phillips SM, Lau KJ, D'Souza AC, Nunes EA. An umbrella review of systematic reviews of beta-hydroxy-beta-methyl butyrate supplementation in ageing and clinical practice. *J Cachexia Sarcopenia Muscle.* 2022;13(5):2265–75.
21. Austin TM, Richter RR, Sebelski CA. Introduction to the GRADE approach for guideline development: considerations for physical therapist practice. *Phys Ther.* 2014;94(11):1652–9.
22. Sadoyu S, Tanni KA, Punrum N, et al. Methodological approaches for assessing certainty of the evidence in umbrella reviews: a scoping review. *PLoS One.* 2022;17(6):e0269009.
23. Nunes EA, D'Souza AC, Steen JP, Phillips SM. Lack of evidence for Omega-3 fatty acid supplementation in enhancing lean mass, muscle strength, and physical function in healthy adults and clinical populations: an overview of reviews. *Clin Nutr ESPEN.* 2025;67:155–65.
24. McLeod JC, Currier BS, Lowisz CV, Phillips SM. The influence of resistance exercise training prescription variables on skeletal muscle mass, strength, and physical function in healthy adults: an umbrella review. *J Sport Health Sci.* 2024;13(1):47–60.
25. Lunny C, Pieper D, Thabet P, Kanji S. Managing overlap of primary study results across systematic reviews: practical considerations for authors of overviews of reviews. *BMC Med Res Methodol.* 2021;21(1):140.
26. Pieper D, Antoine SL, Mathes T, Neugebauer EA, Eikermann M. Systematic review finds overlapping reviews were not mentioned in every other overview. *J Clin Epidemiol.* 2014;67(4):368–75.
27. Kirvaldize M, Abbadi A, Dahlberg L, Sacco LB, Calderón-Larrañaga A, Morin L. Estimating pairwise overlap in umbrella reviews: considerations for using the corrected covered area (CCA) index methodology. *Res Synth Methods.* 2023;14(5):764–7.
28. Bougioukas KI, Diakonidis T, Mavromanoli AC, Haidich AB. ccaR: a package for assessing primary study overlap across systematic reviews in overviews. *Res Synth Methods.* 2023;14(3):443–54.
29. Amiri N, Fathei M, Mosaferi Ziaaldini M. Effects of resistance training on muscle strength, insulin-like growth factor-1, and insulin-like growth factor-binding protein-3 in healthy elderly subjects: a systematic review and meta-analysis of randomized controlled trials. *Hormones (Athens).* 2021;20(2):247–57.
30. Chen YC, Chen W-C, Liu C-W, et al. Is moderate resistance training adequate for older adults with sarcopenia? A systematic review and network meta-analysis of RCTs. *Eur Rev Aging Phys Act.* 2023;20(1):1–22.
31. Cordeiro LD, Linhares DG, dos Santos AOB, dos Santos LL, de Castro JBP, Vale RGD. Influence of resistance training on muscle architecture in older adults: a systematic review and meta-analysis of randomized controlled trials. *Arch Gerontol Geriatr.* 2023;112:105020.
32. Geng Q, Zhai H, Wang L, Wei H, Hou S. The efficacy of different interventions in the treatment of sarcopenia in middle-aged and elderly people: a network meta-analysis. *Medicine (Baltimore).* 2023;102(27):e34254.
33. González-Rocha A, Mendez-Sanchez L, Ortíz-Rodríguez MA, Denova-Gutiérrez E. Effect of exercise on muscle mass, fat mass, bone mass, muscular strength and physical performance in community dwelling older adults: systematic review and meta-analysis. *Aging Dis.* 2022;13(5):1421–35.
34. Kamiya M, Ihira H, Taniguchi Y, et al. Low-intensity resistance training to improve knee extension strength in community-dwelling older adults: systematic review and meta-analysis of randomized controlled studies. *Exp Gerontol.* 2023;172:112041.
35. Wang H, Huang WY, Zhao Y. Efficacy of exercise on muscle function and physical performance in older adults with sarcopenia: an updated systematic review and meta-analysis. *Int J Environ Res Public Health.* 2022;19(13):8212.
36. Zeng D, Ling X-Y, Fang Z-L, Lu Y-F. Optimal exercise to improve physical ability and performance in older adults with sarcopenia: a systematic review and network meta-analysis. *Geriatr Nurs.* 2023;52:199–207.
37. Zhao H, Cheng R, Song G, et al. The effect of resistance training on the rehabilitation of elderly patients with sarcopenia: a meta-analysis. *Int J Environ Res Public Health.* 2022;19(23):15491.
38. Chen N, He X, Feng Y, Ainsworth BE, Liu Y. Effects of resistance training in healthy older people with sarcopenia: a systematic review and meta-analysis of randomized controlled trials. *Eur Rev Aging Phys Act.* 2021;18(1):23.
39. Polito MD, Dias JR Jr, Papst RR. Resistance training to reduce resting blood pressure and increase muscle strength in users and non-users of anti-hypertensive medication: a meta-analysis. *Clin Exp Hypertens.* 2021;43(5):474–85.
40. Polito MD, Papst RR, Farinatti P. Moderators of strength gains and hypertrophy in resistance training: a systematic review and meta-analysis. *J Sports Sci* 2021;39(19):2189–98.
41. Andrews KL, Gallagher S, Herring MP. The effects of exercise interventions on health and fitness of firefighters: a meta-analysis. *Scand J Med Sci Sports* 2019;29(6):780–90.

42. Beckwee D, Delaere A, Aelbrecht S, et al. Exercise interventions for the prevention and treatment of sarcopenia. A systematic umbrella review. *J Nutr Health Aging*. 2019;23(6):494–502.
43. Grgic J, Garofolini A, Orazem J, Sabol F, Schoenfeld BJ, Pedisic Z. Effects of resistance training on muscle size and strength in very elderly adults: a systematic review and meta-analysis of randomized controlled trials. *Sports Med*. 2020;50(11):1983–99.
44. Hagstrom AD, Marshall PW, Halaki M, Hackett DA. The effect of resistance training in women on dynamic strength and muscular hypertrophy: a systematic review with meta-analysis. *Sports Med*. 2020;50(6):1075–93.
45. Beaudart C, Dawson A, Shaw SC, et al; IOF-ESCEO Sarcopenia Working Group. Nutrition and physical activity in the prevention and treatment of sarcopenia: systematic review. *Osteoporos Int*. 2017;28(6):1817–33.
46. Borde R, Hortobagyi T, Granacher U. Dose-response relationships of resistance training in healthy old adults: a systematic review and meta-analysis. *Sports Med*. 2015;45(12):1693–720.
47. De Mello RGB, Dalla Corte RR, Gioscia J, Moriguchi EH. Effects of physical exercise programs on sarcopenia management, dynapenia, and physical performance in the elderly: a systematic review of randomized clinical trials. *J Aging Res*. 2019;2019:1959486.
48. Miranda JM, Polito LFT, Rica RL, et al. Muscle strength training and prescribing in competitive tennis players: a systematic review. *Rev Bras Med Esporte*. 2020;26(1):87–92.
49. Guizelini PC, de Aguiar RA, Denadai BS, Caputo F, Greco CC. Effect of resistance training on muscle strength and rate of force development in healthy older adults: a systematic review and meta-analysis. *Exp Gerontol*. 2018;102:51–8.
50. Moreira OC, Oliveira CEP, Maroto-Izquierdo S, Cuevas MJ, De Paz JA. Effects of short-term strength training on body composition, muscle strength and functional capacity of elderly: a systematic review and meta-analysis. *Biosci J*. 2019;35(6):1941–57.
51. Arnold P, Bautmans I. The influence of strength training on muscle activation in elderly persons: a systematic review and meta-analysis. *Exp Gerontol*. 2014;58:58–68.
52. Asikainen TM, Kukkonen-Harjula K, Miilunpalo S. Exercise for health for early postmenopausal women: a systematic review of randomised controlled trials. *Sports Med*. 2004;34(11):753–78.
53. Beijersbergen CMI, Granacher U, Vandervoort AA, DeVita P, Hortobagyi T. The biomechanical mechanism of how strength and power training improves walking speed in old adults remains unknown. *Ageing Res Rev*. 2013;12(2):618–27.
54. Ferreira ML, Sherrington C, Smith K, et al. Physical activity improves strength, balance and endurance in adults aged 40–65 years: a systematic review. *J Physiother*. 2012;58(3):145–56.
55. Munn J, Herbert RD, Evia SC. Contralateral effects of unilateral resistance training: a meta-analysis. *J Appl Physiol (1985)*. 2004;96(5):1861–6.
56. Silva NL, Oliveira RB, Fleck SJ, Leon ACMP, Farinatti P. Influence of strength training variables on strength gains in adults over 55 years-old: a meta-analysis of dose-response relationships. *J Sci Med Sport*. 2014;17(3):337–44.
57. Ramos-Campo DJ, Andreu Caravaca L, Martinez-Rodriguez A, Rubio-Arias JA. Effects of resistance circuit-based training on body composition, strength and cardiorespiratory fitness: a systematic review and meta-analysis. *Biology (Basel)*. 2021;10(5):377.
58. Ramos-Campo DJ, Andreu-Caravaca L, Carrasco-Poyatos M, Benito PJ, Rubio-Arias JA. Effects of circuit resistance training on body composition, strength, and cardiorespiratory fitness in middle-aged and older women: a systematic review and meta-analysis. *J Aging Phys Act*. 2021;30(4):725–38.
59. Munoz-Martinez FA, Rubio-Arias JA, Ramos-Campo DJ, Alcaraz PE. Effectiveness of resistance circuit-based training for maximum oxygen uptake and upper-body one-repetition maximum improvements: a systematic review and meta-analysis. *Sports Med*. 2017;47(12):2553–68.
60. Colado JC, Mena R, Calatayud J, et al. Effects of strength training with variable elastic resistance across the lifespan: a systematic review. *Cult Cienc Deporte* 2020;15(44):147–64.
61. de Oliveira PA, Blasczyk JC, Souza Junior G, et al. Effects of elastic resistance exercise on muscle strength and functional performance in healthy adults: a systematic review and meta-analysis. *J Phys Act Health*. 2017;14(4):317–27.
62. Tsai YT, Su HH, Chou CH, et al. Impact of elastic band training on functional outcomes and muscle mass in the elderly with sarcopenia: a meta-analysis. *Int J Gerontol*. 2022;16(3):224–30.
63. Chaabene H, Prieske O, Herz M, et al. Home-based exercise programmes improve physical fitness of healthy older adults: a PRISMA-compliant systematic review and meta-analysis with relevance for COVID-19. *Ageing Res Rev*. 2021;67:101265.
64. Thiebaud RS, Funk MD, Abe T. Home-based resistance training for older adults: a systematic review. *Geriatr Gerontol Int*. 2014;14(4):750–7.
65. Baena-Marin M, Rojas-Jaramillo A, Gonzalez-Santamaria J, et al. Velocity-based resistance training on 1-rm, jump and sprint performance: a systematic review of clinical trials. *Sports (Basel)*. 2022;10(1):8.
66. Zhang X, Feng S, Peng R, Li H. The role of velocity-based training (VBT) in enhancing athletic performance in trained individuals: a meta-analysis of controlled trials. *Int J Environ Res Public Health*. 2022;19(15):9252.
67. Nunez Sanchez FJ, Saez de Villarreal E. Does flywheel paradigm training improve muscle volume and force? A meta-analysis. *J Strength Cond Res*. 2017;31(11):3177–86.
68. Medeiros DM, Marchiori C, Baroni BM. Effect of Nordic hamstring exercise training on knee flexors eccentric strength and fascicle length: a systematic review and meta-analysis. *J Sport Rehabil*. 2020;30(3):482–91.
69. Morris SJ, Oliver JL, Pedley JS, Haff GG, Lloyd RS. Comparison of weightlifting, traditional resistance training and plyometrics on strength, power and speed: a systematic review with meta-analysis. *Sports Med*. 2022;52(7):1533–54.
70. Behm DG, Muehlbauer T, Kibele A, Granacher U. Effects of strength training using unstable surfaces on strength, power and balance performance across the lifespan: a systematic review and meta-analysis. *Sports Med*. 2015;45(12):1645–69.
71. Androulakis-Korakakis P, Fisher JP, Steele J. The minimum effective training dose required to increase 1RM strength in resistance-trained men: a systematic review and meta-analysis. *Sports Med*. 2020;50(4):751–65.
72. Cuthbert M, Haff GG, Arent SM, et al. Effects of variations in resistance training frequency on strength development in well-trained populations and implications for in-season athlete training: a systematic review and meta-analysis. *Sports Med*. 2021;51(9):1967–82.
73. Kessinger TK, Melton B, Miyashita T, Ryan G. The effectiveness of frequency-based resistance training protocols on muscular performance and hypertrophy in trained males: a critically appraised topic. *J Sport Rehabil*. 2020;29(7):1024–31.
74. Kneffel Z, Murlasits Z, Reed J, Krieger J. A meta-regression of the effects of resistance training frequency on muscular strength and hypertrophy in adults over 60 years of age. *J Sports Sci*. 2021;39(3):351–8.
75. Ralston GW, Kilgore L, Wyatt FB, Buchan D, Baker JS. Weekly training frequency effects on strength gain: a meta-analysis. *Sports Med Open*. 2018;4(1):36.

76. Grgic J, Schoenfeld BJ, Davies TB, Lazinica B, Krieger JW, Pedisic Z. Effect of resistance training frequency on gains in muscular strength: a systematic review and meta-analysis. *Sports Med*. 2018;48(5):1207–20.
77. Bandeira-Guimaraes M, Blanco-Rambo E, Vieira AF, et al. Chronic effects of different intensities of power training on neuromuscular parameters in older people: a systematic review with meta-analysis. *Sports Med Open*. 2023;9(1):98.
78. Carvalho L, Junior RM, Barreira J, Schoenfeld BJ, Orazem J, Barroso R. Muscle hypertrophy and strength gains after resistance training with different volume-matched loads: a systematic review and meta-analysis. *Appl Physiol Nutr Metab*. 2022;47(4):357–68.
79. Csapo R, Alegre LM. Effects of resistance training with moderate vs heavy loads on muscle mass and strength in the elderly: a meta-analysis. *Scand J Med Sci Sports*. 2016;26(9):995–1006.
80. Lopez P, Radaelli R, Taaffe DR, et al. Resistance training load effects on muscle hypertrophy and strength gain: systematic review and network meta-analysis. *Med Sci Sports Exerc*. 2021;53(6):1206–16.
81. Refalo MC, Hamilton DL, Paval DR, Gallagher IJ, Feros SA, Fyfe JJ. Influence of resistance training load on measures of skeletal muscle hypertrophy and improvements in maximal strength and neuromuscular task performance: a systematic review and meta-analysis. *J Sports Sci*. 2021;39(15):1723–45.
82. Steib S, Schoene D, Pfeifer K. Dose-response relationship of resistance training in older adults: a meta-analysis. *Med Sci Sports Exerc*. 2010;42(5):902–14.
83. Schoenfeld BJ, Grgic J, Ogborn D, Krieger JW. Strength and hypertrophy adaptations between low- vs. high-load resistance training: a systematic review and meta-analysis. *J Strength Cond Res*. 2017;31(12):3508–23.
84. Schoenfeld BJ, Wilson JM, Lowery RP, Krieger JW. Muscular adaptations in low- versus high-load resistance training: a meta-analysis. *Eur J Sport Sci*. 2016;16(1):1–10.
85. Maroto-Izquierdo S, Garcia-Lopez D, Fernandez-Gonzalo R, Moreira OC, González-Gallego J, de Paz JA. Skeletal muscle functional and structural adaptations after eccentric overload flywheel resistance training: a systematic review and meta-analysis. *J Sci Med Sport*. 2017;20(10):943–51.
86. Oranchuk DJ, Storey AG, Nelson AR, Cronin JB. Isometric training and long-term adaptations: effects of muscle length, intensity, and intent: a systematic review. *Scand J Med Sci Sports*. 2019;29(4):484–503.
87. Pallares JG, Hernández-Belmonte A, Martínez-Cava A, Vetrovsky T, Steffl M, Courel-Ibáñez J. Effects of range of motion on resistance training adaptations: a systematic review and meta-analysis. *Scand J Med Sci Sports*. 2021;31(10):1866–81.
88. Bagenhammar S, Hansson EE. Repeated sets or single set of resistance training—a systematic review. *Adv Physiother*. 2007;9(4):154–60.
89. Ralston G, Kilgore L, Wyatt F, Baker J. The effect of weekly set volume on strength gain: a meta-analysis. *Sports Med*. 2017;47(12):2585–601.
90. Ralston GW, Kilgore L, Wyatt FB, et al. Re-examination of 1- vs. 3-sets of resistance exercise for pre-spaceflight muscle conditioning: a systematic review and meta-analysis. *Front Physiol*. 2019;10:864.
91. Rocha JNS, Pereira-Monteiro MR, Vasconcelos ABS, Pantoja-Cardoso A, Aragao-Santos JC, Da Silva-Grigoletto ME. Different resistance training volumes on strength, functional fitness, and body composition of older people: a systematic review with meta-analysis. *Arch Gerontol Geriatr*. 2023;119:105303.
92. Wolfe BL, LeMura LM, Cole PJ. Quantitative analysis of single- vs. multiple-set programs in resistance training. *J Strength Cond Res*. 2004;18(1):35–47.
93. Krieger JW. Single versus multiple sets of resistance exercise: a meta-regression. *J Strength Cond Res*. 2009;23(6):1890–901.
94. Eddens L, van Someren K, Howatson G. The role of intra-session exercise sequence in the interference effect: a systematic review with meta-analysis. *Sports Med*. 2018;48(1):177–88.
95. Murlasits Z, Kneffel Z, Thalib L. The physiological effects of concurrent strength and endurance training sequence: a systematic review and meta-analysis. *J Sports Sci*. 2018;36(11):1212–9.
96. Nunes JP, Grgic J, Cunha PM, et al. What influence does resistance exercise order have on muscular strength gains and muscle hypertrophy? A systematic review and meta-analysis. *Eur J Sport Sci*. 2021;21(2):149–57.
97. Davies T, Orr R, Halaki M, Hackett D. Effect of training leading to repetition failure on muscular strength: a systematic review and meta-analysis. *Sports Med*. 2016;46(4):487–502.
98. Grgic J, Schoenfeld BJ, Orazem J, Sabol F. Effects of resistance training performed to repetition failure or non-failure on muscular strength and hypertrophy: a systematic review and meta-analysis. *J Sport Health Sci*. 2022;11(2):202–11.
99. Vieira AF, Umpierre D, Teodoro JL, et al. Effects of resistance training performed to failure or not to failure on muscle strength, hypertrophy, and power output: a systematic review with meta-analysis. *J Strength Cond Res*. 2021;35(4):1165–75.
100. Haugen ME, Vårvik FT, Larsen S, Haugen AS, van den Tilhaar R, Bjørnsen T. Effect of free-weight vs. machine-based strength training on maximal strength, hypertrophy and jump performance—a systematic review and meta-analysis. *BMC Sports Sci Med Rehabil*. 2023;15(1):1–20.
101. Davies TB, Kuang K, Orr R, Halaki M, Hackett D. Effect of movement velocity during resistance training on dynamic muscular strength: a systematic review and meta-analysis. *Sports Med*. 2017;47(8):1603–17.
102. Grgic J, Lazinica B, Garofolini A, Schoenfeld BJ, Saner NJ, Mikulic P. The effects of time of day-specific resistance training on adaptations in skeletal muscle hypertrophy and muscle strength: a systematic review and meta-analysis. *Chronobiol Int*. 2019;36(4):449–60.
103. Grgic J, Schoenfeld BJ, Skrepnik M, Davies TB, Mikulic P. Effects of rest interval duration in resistance training on measures of muscular strength: a systematic review. *Sports Med*. 2018;48(1):137–51.
104. Roig M, O'Brien K, Kirk G, et al. The effects of eccentric versus concentric resistance training on muscle strength and mass in healthy adults: a systematic review with meta-analysis. *Br J Sports Med*. 2009;43(8):556–68.
105. Balachandran AT, Steele J, Angielczyk D, et al. Comparison of power training vs traditional strength training on physical function in older adults: a systematic review and meta-analysis. *JAMA Netw Open*. 2022;5(5):e2211623.
106. Pearson LT, Behm DG, Goodall S, Mason R, Stuart S, Barry G. Effects of maximal-versus submaximal-intent resistance training on functional capacity and strength in community-dwelling older adults: a systematic review and meta-analysis. *BMC Sports Sci Med Rehabil*. 2022;14(1):129.
107. Davies TB, Tran DL, Hogan CM, Haff GG, Latella C. Chronic effects of altering resistance training set configurations using cluster sets: a systematic review and meta-analysis. *Sports Med*. 2021;51(4):707–36.
108. Jukic I, Van Hooren B, Ramos AG, Helms ER, McGuigan MR, Tufano JJ. The effects of set structure manipulation on

- chronic adaptations to resistance training: a systematic review and meta-analysis. *Sports Med.* 2021;51(5):1061–86.
109. Marshall J, Bishop C, Turner A, Haff GG. Optimal training sequences to develop lower body force, velocity, power, and jump height: a systematic review with meta-analysis. *Sports Med.* 2021;51(6):1245–71.
 110. Centner C, Wiegel P, Gollhofer A, König D. Effects of blood flow restriction training on muscular strength and hypertrophy in older individuals: a systematic review and meta-analysis. *Sports Med.* 2018;49(1):95–108.
 111. Lixandrão ME, Ugrinowitsch C, Berton R, et al. Magnitude of muscle strength and mass adaptations between high-load resistance training versus low-load resistance training associated with blood-flow restriction: a systematic review and meta-analysis. *Sports Med.* 2018;48(2):361–78.
 112. Rodrigo-Mallorca D, Loaiza-Betancur AF, Monteagudo P, Blasco-Lafarga C, Chulvi-Medrano I. Resistance training with blood flow restriction compared to traditional resistance training on strength and muscle mass in non-active older adults: a systematic review and meta-analysis. *Int J Environ Res Public Health.* 2021;18(21):11441.
 113. Hickmott LM, Chilibeck PD, Shaw KA, Butcher SJ. The effect of load and volume autoregulation on muscular strength and hypertrophy: a systematic review and meta-analysis. *Sports Med Open.* 2022;8(1):9.
 114. Larsen S, Kristiansen E, van den Tillaar R. Effects of subjective and objective autoregulation methods for intensity and volume on enhancing maximal strength during resistance-training interventions: a systematic review. *PeerJ.* 2021;9:e10663.
 115. Kassiano W, Nunes JP, Costa B, Ribeiro AS, Schoenfeld BJ, Cyrino ES. Does varying resistance exercises promote superior muscle hypertrophy and strength gains? A systematic review. *J Strength Cond Res.* 2022;36(6):1753–62.
 116. Harries SK, Lubans DR, Callister R. Systematic review and meta-analysis of linear and undulating periodized resistance training programs on muscular strength. *J Strength Cond Res.* 2015;29(4):1113–25.
 117. Moesgaard L, Beck MM, Christiansen L, Aagaard P, Lundbye-Jensen J. Effects of periodization on strength and muscle hypertrophy in volume-equated resistance training programs: a systematic review and meta-analysis. *Sports Med.* 2022;52(7):1647–66.
 118. Rhea MR, Alderman BL. A meta-analysis of periodized versus nonperiodized strength and power training programs. *Res Q Exerc Sport.* 2004;75(4):413–22.
 119. Cadore EL, Izquierdo M. How to simultaneously optimize muscle strength, power, functional capacity, and cardiovascular gains in the elderly: an update. *Age (Dordr).* 2013;35(6):2329–44.
 120. Huijberts RO, Wüst RCI, van der Zwaard S. Concurrent strength and endurance training: a systematic review and meta-analysis on the impact of sex and training status. *Sports Med.* 2024;54(2):485–503.
 121. Petré H, Hemmingsson E, Rosdahl H, Psilander N. Development of maximal dynamic strength during concurrent resistance and endurance training in untrained, moderately trained, and trained individuals: a systematic review and meta-analysis. *Sports Med.* 2021;51(5):991–1010.
 122. Sabag A, Najafi A, Michael S, Esgin T, Halaki M, Hackett D. The compatibility of concurrent high intensity interval training and resistance training for muscular strength and hypertrophy: a systematic review and meta-analysis. *J Sports Sci.* 2018;36(21):2472–83.
 123. Trowell D, Vicenzino B, Saunders N, Fox A, Bonacci J. Effect of strength training on biomechanical and neuromuscular variables in distance runners: a systematic review and meta-analysis. *Sports Med.* 2020;50(1):133–50.
 124. Wilson JM, Marin PJ, Rhea MR, Wilson SMC, Loenneke JP, Anderson JC. Concurrent training: a meta-analysis examining interference of aerobic and resistance exercises. *J Strength Cond Res.* 2012;26(8):2293–307.
 125. Cirer-Sastre R, Beltrán-Garrido JV, Corbi F. Contralateral effects after unilateral strength training: a meta-analysis comparing training loads. *J Sports Sci Med.* 2017;16(2):180–6.
 126. Manca A, Dragone D, Dvir Z, Deriu F. Cross-education of muscular strength following unilateral resistance training: a meta-analysis. *Eur J Appl Physiol.* 2017;117(11):2335–54.
 127. Yagiz G, Akaras E, Kubis HP, Owen JA. The effects of resistance training on architecture and volume of the upper extremity muscles: a systematic review of randomised controlled trials and meta-analyses. *Appl Sci.* 2022;12(3):1593.
 128. Raya-Gonzalez J, Martinez Sanchez MA. Training methods and nutritional aspects to increase muscle mass: a systematic review. *Arch Med Deporte.* 2019;36(6):376–85.
 129. Schoenfeld BJ, Ogborn DI, Vigotsky AD, Franchi MV, Krieger JW. Hypertrophic effects of concentric vs. eccentric muscle actions: a systematic review and meta-analysis. *J Strength Cond Res.* 2017;31(9):2599–608.
 130. Baz-Valle E, Balsalobre-Fernández C, Alix-Fages C, Santos-Concejero J. A systematic review of the effects of different resistance training volumes on muscle hypertrophy. *J Hum Kinet.* 2022;81(1):199–210.
 131. Bernardez-Vazquez R, Raya-Gonzalez J, Castillo D, Beato M. Resistance training variables for optimization of muscle hypertrophy: an umbrella review. *Front Sports Act Living.* 2022;4:949021.
 132. Krieger JW. Single vs. multiple sets of resistance exercise for muscle hypertrophy: a meta-analysis. *J Strength Cond Res.* 2010;24(4):1150–9.
 133. Schoenfeld BJ, Ogborn D, Krieger JW. Dose-response relationship between weekly resistance training volume and increases in muscle mass: a systematic review and meta-analysis. *J Sports Sci.* 2017;35(11):1073–82.
 134. Schoenfeld BJ, Grgic J, Krieger J. How many times per week should a muscle be trained to maximize muscle hypertrophy? A systematic review and meta-analysis of studies examining the effects of resistance training frequency. *J Sports Sci.* 2019;37(11):1286–95.
 135. Schoenfeld BJ, Ogborn D, Krieger JW. Effects of resistance training frequency on measures of muscle hypertrophy: a systematic review and meta-analysis. *Sports Med.* 2016;46(11):1689–97.
 136. Grgic J. The effects of low-load vs. high-load resistance training on muscle fiber hypertrophy: a meta-analysis. *J Hum Kinet.* 2020;74(1):51–8.
 137. Refalo MC, Helms ER, Trexler ET, Hamilton DL, Fyfe JJ. Influence of resistance training proximity-to-failure on skeletal muscle hypertrophy: a systematic review with meta-analysis. *Sports Med.* 2023;53(3):649–65.
 138. Fuentes-García MA, Malchrowicz-Mosko E, Castañeda-Babarro A. Effects of variable resistance training versus conventional resistance training on muscle hypertrophy: a systematic review. *Sport Sci Health.* 2023;20(1):37–45.
 139. Hackett DA, Davies TB, Orr R, Kuang K, Halaki M. Effect of movement velocity during resistance training on muscle-specific hypertrophy: a systematic review. *Eur J Sport Sci.* 2018;18(4):473–82.
 140. Schoenfeld BJ, Ogborn DI, Krieger JW. Effect of repetition duration during resistance training on muscle hypertrophy: a systematic review and meta-analysis. *Sports Med.* 2015;45(4):577–85.

141. Grgic J, Lazinica B, Mikulic P, Schoenfeld BJ. Should resistance training programs aimed at muscular hypertrophy be periodized? A systematic review of periodized versus non-periodized approaches. *Sci Sports*. 2018;33(3):e97–e104.
142. Rosa A, Vazquez G, Grgic J, Balachandran AT, Orazem J, Schoenfeld BJ. Hypertrophic effects of single- versus multi-joint exercise of the limb muscles: a systematic review and meta-analysis. *Strength Cond J*. 2023;45(1):49–57.
143. Grgic J, Lazinica B, Mikulic P, Krieger JW, Schoenfeld BJ. The effects of short versus long inter-set rest intervals in resistance training on measures of muscle hypertrophy: a systematic review. *Eur J Sport Sci*. 2017;17(8):983–93.
144. Henselmans M, Schoenfeld B. The effect of inter-set rest intervals on resistance exercise-induced muscle hypertrophy. *Sports Med*. 2014;44(12):1635–43.
145. Schoenfeld BJ, Grgic J. Effects of range of motion on muscle development during resistance training interventions: a systematic review. *SAGE Open Med*. 2020;8:2050312120901559.
146. Sodal LK, Kristiansen E, Larsen S, van den Tillaar R. Effects of drop sets on skeletal muscle hypertrophy: a systematic review and meta-analysis. *Sports Med Open*. 2023;9(1):66.
147. Alcaraz-Ibanez M, Rodriguez-Perez M. Effects of resistance training on performance in previously trained endurance runners: a systematic review. *J Sports Sci*. 2018;36(6):613–29.
148. Straight CR, Lindheimer JB, Brady AO, Dishman RK, Evans EM. Effects of resistance training on lower-extremity muscle power in middle-aged and older adults: a systematic review and meta-analysis of randomized controlled trials. *Sports Med*. 2016;46(3):353–64.
149. Soriano MA, Jimenez-Reyes P, Rhea MR, Marin PJ. The optimal load for maximal power production during lower-body resistance exercises: a meta-analysis. *Sports Med*. 2015;45(8):1191–205.
150. Soriano MA, Suchomel TJ, Marin PJ. The optimal load for maximal power production during upper-body resistance exercises: a meta-analysis. *Sports Med*. 2017;47(4):757–68.
151. Berryman N, Mujika I, Arvaisis D, Roubeix M, Binet C, Bosquet L. Strength training for middle- and long-distance performance: a meta-analysis. *Int J Sports Physiol Perform*. 2018;13(1):57–63.
152. Almeida FJ, Melo MH, Nogueira R, Prazeres J, Costa C, Gambassi BB. Do all resistance exercise protocols improve the functional parameters of the elderly? A review study. *Asian J Sports Med*. 2020;11(4):1–7.
153. Hortobagyi T, Lesinski M, Gabler M, VanSwearingen JM, Malatesta D, Granacher U. Effects of three types of exercise interventions on healthy old adults' gait speed: a systematic review and meta-analysis. *Sports Med*. 2015;45(12):1627–43.
154. Howe TE, Rochester L, Neil F, Skelton DA, Ballinger C. Exercise for improving balance in older people. *Cochrane Database Syst Rev*. 2011;2011(11):CD004963.
155. Van Abbema R, De Greef M, Craje C, Krijnen W, Hobbelen H, Van Der Schans C. What type, or combination of exercise can improve preferred gait speed in older adults? A meta-analysis. *BMC Geriatr*. 2015;15:72.
156. Martins AD, Fernandes O, Pereira A, et al. The effects of high-speed resistance training on health outcomes in independent older adults: a systematic review and meta-analysis. *Int J Environ Res Public Health*. 2022;19(9):5390.
157. da Rosa Orssatto LB, de la Rocha Freitas C, Shield AJ, Silveira Pinto R, Trajano GS. Effects of resistance training concentric velocity on older adults' functional capacity: a systematic review and meta-analysis of randomised trials. *Exp Gerontol*. 2019;127:110731.
158. Thomas E, Battaglia G, Patti A, et al. Physical activity programs for balance and fall prevention in elderly: a systematic review. *Medicine (Baltimore)*. 2019;98(27):e16218.
159. Dipietro L, Campbell WW, Buchner DM, et al; 2018 Physical Activity Guidelines Advisory Committee. Physical activity, injurious falls, and physical function in aging: an umbrella review. *Med Sci Sports Exerc*. 2019;51(6):1303–13.
160. Tschopp M, Sattelmayer MK, Hilfiker R. Is power training or conventional resistance training better for function in elderly persons? A meta-analysis. *Age Ageing*. 2011;40(5):549–56.
161. Seitz LB, Reyes A, Tran TT, Saez de Villarreal E, Haff GG. Increases in lower-body strength transfer positively to sprint performance: a systematic review with meta-analysis. *Sports Med*. 2014;44(12):1693–702.
162. Raya-Gonzalez J, Prat-Luri A, Lopez-Valenciano A, Sabido R, Hernández-Davó JL. Effects of flywheel resistance training on sport actions. A systematic review and meta-analysis. *J Hum Kinet*. 2021;77(1):191–204.
163. Blazevich AJ, Wilson CJ, Alcaraz PE, Rubio-Arias JA. Effects of resistance training movement pattern and velocity on isometric muscular rate of force development: a systematic review with meta-analysis and meta-regression. *Sports Med*. 2020;50(5):943–63.
164. Morton RW, Colenso-Semple L, Phillips SM. Training for strength and hypertrophy: an evidence-based approach. *Cur Opin Physiol*. 2019;10:90–5.
165. Helms ER, Byrnes RK, Cooke DM, et al. RPE vs. percentage 1RM loading in periodized programs matched for sets and repetitions. *Front Physiol*. 2018;9:247.
166. Lorenz D, Morrison S. Current concepts in periodization of strength and conditioning for the sports physical therapist. *Int J Sports Phys Ther*. 2015;10(6):734–47.
167. Stone MH, Hornsby WG, Haff GG, et al. Periodization and block periodization in sports: emphasis on strength-power training—a provocative and challenging narrative. *J Strength Cond Res*. 2021;35(8):2351–71.
168. Cunanan AJ, DeWeese BH, Wagle JP, et al. The general adaptation syndrome: a foundation for the concept of periodization. *Sports Med*. 2018;48(4):787–97.
169. Hyde ET, Whitfield GP, Omura JD, Fulton JE, Carlson SA. Trends in meeting the physical activity guidelines: muscle-strengthening alone and combined with aerobic activity, United States, 1998-2018. *J Phys Act Health*. 2021;18(S1):S37–44.
170. Calatayud J, Lopez-Bueno R, Nunez-Cortes R, Yang L, Del Pozo Cruz B, Andersen LL. Trends in adherence to the muscle-strengthening activity guidelines in the US over a 20-year span. *Gen Hosp Psychiatry*. 2023;84:89–95.
171. Abildso CG, Daily SM, Renee Umstaddt Meyer MR, Perry CK, Eyler A. Prevalence of meeting aerobic, muscle-strengthening, and combined physical activity guidelines during leisure time among adults, by rural-urban classification and region—United States. *MMWR Morb Mortal Wkly Rep*. 2023;72(4):85–9.
172. Bennie JA, Lee DC, Khan A, et al. Muscle-strengthening exercise among 397,423 U.S. adults: prevalence, correlates, and associations with health conditions. *Am J Prev Med*. 2018;55(6):864–74.
173. Centers for Disease Control and Prevention. QuickStats: percentage of adults aged ≥18 years who met the federal guidelines for muscle-strengthening physical activity,† by age group and sex—National Health Interview Survey, United States, 2020. *MMWR Morb Mortal Wkly Rep*. 2022;71(18):642.
174. Gluchowski A, Bilsborough H, McDermott J, Hawley-Hague H, Todd C. “A lot of people just go for walks, and don't do anything else”: older adults in the UK are not aware of the strength component embedded in the chief medical officers'

- physical activity guidelines—a qualitative study. *Int J Environ Res Public Health*. 2022;19(16):10002.
175. Hyde ET, Brown DR, Webber BJ, et al. Meeting the aerobic and muscle-strengthening physical activity guidelines among older US adults, National Health Interview Survey 1998–2018. *J Appl Gerontol*. 2024;43(8):1003–14.
 176. Carpinelli RN. Challenging the American College of Sports Medicine 2009 position stand on resistance training. *Med Sport*. 2009;13(2):131–7.
 177. Fyfe JJ, Hamilton DL, Daly RM. Minimal-dose resistance training for improving muscle mass, strength, and function: a narrative review of current evidence and practical considerations. *Sports Med*. 2022;52(3):463–79.
 178. Cureton KJ, Collins MA, Hill DW, McElhannon FM Jr. Muscle hypertrophy in men and women. *Med Sci Sports Exerc*. 1988;20(4):338–44.
 179. Hickson RC, Hidaka K, Foster C. Skeletal muscle fiber type, resistance training, and strength-related performance. *Med Sci Sports Exerc*. 1994;26(5):593–8.
 180. Candow DG, Burke DG. Effect of short-term equal-volume resistance training with different workout frequency on muscle mass and strength in untrained men and women. *J Strength Cond Res*. 2007;21(1):204–7.
 181. Faigenbaum AD, Kamis D, Rial Rebullido T. A strong approach for overcoming the FoRE: fear of resistance exercise. *ACSM'S Health Fit J*. 2024;28(3):14–20.
 182. Niemeijer A, Lund H, Stafne SN, et al. Adverse events of exercise therapy in randomised controlled trials: a systematic review and meta-analysis. *Br J Sports Med*. 2020;54(18):1073–80.
 183. Hollings M, Mavros Y, Freeston J, Fiatarone Singh M. The effect of progressive resistance training on aerobic fitness and strength in adults with coronary heart disease: a systematic review and meta-analysis of randomised controlled trials. *Eur J Prev Cardiol*. 2017;24(12):1242–59.
 184. Swinton PA, Schoenfeld BJ, Murphy A. Dose-response modelling of resistance exercise across outcome domains in strength and conditioning: a meta-analysis. *Sports Med*. 2024;54(6):1579–94.
 185. Landers J. Maximum based on reps. *J Strength Cond Res*. 1984;6(6):60–1.
 186. Nuzzo JL, Pinto MD, Nosaka K, Steele J. Maximal number of repetitions at percentages of the one repetition maximum: a meta-regression and moderator analysis of sex, age, training status, and exercise. *Sports Med*. 2024;54(2):303–21.
 187. Pelland JC, Remmert JF, Robinson ZP, Hinson S, and Zourdos MC. The resistance training dose-response: meta-regressions exploring the effects of weekly volume and frequency on muscle hypertrophy and strength gains. *Sports Med*. 2025. doi: 10.1007/s40279-025-02344-w. Online ahead of print.
 188. Refalo MC, Helms ER, Robinson ZP, Hamilton DL, Fyfe JJ. Similar muscle hypertrophy following eight weeks of resistance training to momentary muscular failure or with repetitions-in-reserve in resistance-trained individuals. *J Sports Sci*. 2024;42(1):85–101.
 189. Helms ER, Cronin J, Storey A, Zourdos MC. Application of the repetitions in reserve-based rating of perceived exertion scale for resistance training. *Strength Cond J*. 2016;38(4):42–9.
 190. Robinson ZP, Pelland JC, Remmert JF, et al. Exploring the dose–response relationship between estimated resistance training proximity to failure, strength gain, and muscle hypertrophy: a series of meta-regressions. *Sports Med*. 2024;54(9):2209–31.
 191. Rhodes RE, Lubans DR, Karunamuni N, Kennedy S, Plotnikoff R. Factors associated with participation in resistance training: a systematic review. *Br J Sports Med*. 2017;51(20):1466–72.
 192. Room J, Hannink E, Dawes H, Barker K. What interventions are used to improve exercise adherence in older people and what behavioural techniques are they based on? A systematic review. *BMJ Open*. 2017;7(12):e019221.
 193. Gutierrez-Arias R, Pieper D, Lunny C, Torres-Castro R, Aguilera-Eguia R, Seron P. Strategies used to manage overlap of primary study data by exercise-related overviews: protocol for a systematic methodological review. *BMJ Open*. 2023;13(4):e069906.
 194. Martinez-Calderon J. Overviews of systematic reviews in sports and exercise medicine: what are they and why are they important? *Br J Sports Med*. 2023;57(16):1005–6.
 195. Nuzzo JL, Pinto MD, Kirk BJC, Nosaka K. Resistance exercise minimal dose strategies for increasing muscle strength in the general population: an overview. *Sports Med*. 2024;54(5):1139–62.
 196. Schoenfeld BJ, Grgic J, Van Every DW, Plotkin DL. Loading recommendations for muscle strength, hypertrophy, and local endurance: a re-examination of the repetition continuum. *Sports (Basel)*. 2021;9(2):32.
 197. Halperin I, Malleron T, Har-Nir I, et al. Accuracy in predicting repetitions to task failure in resistance exercise: a scoping review and exploratory meta-analysis. *Sports Med*. 2022;52(2):377–90.
 198. Lopez P, Izquierdo M, Radaelli R, et al. Effectiveness of multimodal training on functional capacity in frail older people: a meta-analysis of randomized controlled trials. *J Aging Phys Act*. 2018;26(3):407–18.
 199. Makaruk H, Starzak M, Plaszewski M, Winchester JB. Internal validity in resistance training research: a systematic review. *J Sports Sci Med*. 2022;21(2):308–31.
 200. Memon AR, Owen PJ, Anderson N, Verhagen E, Mundell NL, Belavy DL. Common issues of systematic reviews in the sports and exercise medicine field. *BMJ Open Sport Exerc Med*. 2024;10(1):e001784.
 201. Paras T, Sabzevari S, Solomon D, Smith C, McDonough C, Lin A. Trends in level of evidence of systematic reviews in sports medicine, 2010–2020: a systematic review and meta-analysis. *Orthop J Sports Med*. 2022;10(9):23259671221121330.
 202. Slade SC, Dionne CE, Underwood M, Buchbinder R. Consensus on exercise reporting template (CERT): explanation and elaboration statement. *Br J Sports Med*. 2016;50(23):1428–37.
 203. Ardern CL, Buttner F, Andrade R, et al. Implementing the 27 PRISMA 2020 Statement items for systematic reviews in the sport and exercise medicine, musculoskeletal rehabilitation and sports science fields: the PERSiST (implementing Prisma in Exercise, Rehabilitation, Sport medicine and Sport science) guidance. *Br J Sports Med*. 2022;56(4):175–95.