

REVIEW

OPEN

Physical exercise in liver diseases

Yunwei Zhang¹  | Chunyan Cao¹  | Chaofan Li^{2,3}  | Russell G. Witt¹  |
 Hai Huang⁴  | Allan Tsung¹  | Hongji Zhang¹ 

¹Department of Surgery, University of Virginia, Charlottesville, Virginia, USA

²Department of Medicine, Beirne B. Carter Center for Immunology Research, University of Virginia, Charlottesville, Virginia, USA

³Division of Infectious Disease and International Health, Department of Medicine, University of Virginia, Charlottesville, Virginia, USA

⁴Division of Hepatology, Center for Immunology and Inflammation, Departments of Molecular Medicine, Medicine, and Surgery at the School of Medicine, Feinstein Institutes for Medical Research, Manhasset, New York, USA

Correspondence

Allan Tsung, Department of Surgery, University of Virginia, Charlottesville, VA 22903, USA.
 Email: allantsung@virginia.edu

Hongji Zhang, Department of Surgery, University of Virginia, Charlottesville, VA 22903, USA.
 Email: jhn5wx@virginia.edu

Abstract

Liver diseases contribute to ~2 million deaths each year and account for 4% of all deaths globally. Despite various treatment options, the management of liver diseases remains challenging. Physical exercise is a promising nonpharmacological approach to maintain and restore homeostasis and effectively prevent and mitigate liver diseases. In this review, we delve into the mechanisms of physical exercise in preventing and treating liver diseases, highlighting its effects on improving insulin sensitivity, regulating lipid homeostasis, and modulating immune function. In addition, we evaluate the impact of physical exercise on various liver diseases, including liver ischemia/reperfusion injury, cardiogenic liver disease, metabolic dysfunction-associated steatotic liver disease, portal hypertension, cirrhosis, and liver cancer. In conclusion, the review underscores the effectiveness of physical exercise as a beneficial intervention in combating liver diseases.

INTRODUCTION

Liver diseases have become a major health concern and highly impact young individuals, ranking as the 12th leading cause of disability-adjusted life-years among those aged 25–49.^[1] Metabolic dysfunction-associated steatotic liver disease (MASLD), the most prevalent form of liver disease, impacts ~25% of the global adult population.^[2,3] The major causes of mortality related to liver diseases are cirrhosis and HCC.^[4,5] With a deep understanding of the pathophysiological mechanisms of liver diseases, the medical community's choice of therapeutic approaches has been evolving.^[6]

Due to its beneficial effects on metabolism regulation, physical exercise has been extensively validated

as an effective strategy in preventing and treating metabolic syndrome, including MASLD.^[7] In addition, recent studies underscored its potential value in managing other liver diseases, such as cirrhosis and HCC.^[8,9] Bed rest was once considered the cornerstone of the treatment of most liver diseases, such as viral hepatitis and cirrhosis.^[10,11] This approach stems from the research that less physical exercise reduces metabolic demands on the liver and lowers the risk of complications^[12] and that an upright posture further increases plasma renin levels.^[10] However, recent studies revealed the adverse consequences of prolonged inactivity, including muscle atrophy,^[13] deep vein thrombosis,^[14] and elevated risk of fibrosis.^[15] As a

Abbreviations: Akt, protein kinase B; AMPK, AMP-activated protein kinase; A β , amyloid- β peptide; FFAs, free fatty acids; HFD, high-fat diet; HIIT, high-intensity interval exercise; HMGCS2, hydroxy-3-methylglutaryl-CoA synthase 2; IHTG, intrahepatic triglyceride; I/R, ischemia/reperfusion; MASH, metabolic dysfunction-associated steatohepatitis; MASLD, metabolic dysfunction-associated steatotic liver disease; MET, metabolic equivalent of task; PH, portal hypertension; PPAR- α , peroxisome proliferator-activated receptor alpha; PPAR- γ , peroxisome proliferator-activated receptor gamma; RCTs, randomized controlled trials; TG, triglyceride.

Yunwei Zhang and Chunyan Cao have contributed equally to this work.

This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Copyright © 2024 The Author(s). Published by Wolters Kluwer Health, Inc.

result, the prescription of bed rest has been critically reevaluated. Physical exercise is an effective measure for managing cirrhosis-related complications, including HE and sarcopenia.^[16,17] Beyond that, regular physical exercise has also been linked to reduced risk of liver cancer.^[18] Furthermore, studies demonstrated that regular exercise not only diminishes the likelihood of MASLD but also elevates the overall quality of life for individuals afflicted with chronic liver diseases and liver transplantation.^[19,20] Physical exercise also ameliorates systemic metabolic disorders like obesity and type 2 diabetes, both of which can exacerbate liver injury.^[21,22]

In this review, we summarize the protective roles of physical exercise in liver diseases. We discuss physical exercise modalities and how physical exercise prevents and manages liver diseases. Furthermore, we delve into physical exercise's role in mitigating acute and chronic liver inflammation and its potential in managing liver cancer. This exploration underscores the vital role of physical exercise in liver diseases and advocates for its incorporation into liver disease prevention and management strategies.

PHYSICAL EXERCISE MODALITY, INTENSITY, AND DURATION

Physical exercise is mainly comprised of aerobic and resistance training, each with different benefits and precautions for liver diseases. Aerobic exercise effectively reduces body weight, glycosylated hemoglobin A1c, blood pressure, and serum cholesterol levels.^[23–25] However, it can induce fatigue and discomfort, potentially leading to poor long-term compliance, especially for individuals with lower cardiorespiratory fitness levels.^[26] Resistance exercise, which entails muscle contractions against external resistance to boost muscle strength, bone density, and endurance, has been shown to provide significant benefits in addressing dyslipidemia, hypertension, and insulin resistance,^[27,28] offering metabolic advantages with relatively lower energy expenditure; however, limited access to appropriate equipment and facilities can pose a barrier to engaging in effective resistance exercise.

No particular liver disease has been exclusively linked to a specific exercise modality for treatment or management.^[29] A comparative randomized controlled trial (RCT) investigated the effects of moderate-intensity aerobic exercise versus resistance exercise on patients with MASLD, finding both regimens equally effective in reducing intrahepatic triglycerides (IHTGs) and improving insulin resistance.^[30] In patients with cirrhosis, aerobic exercise enhances cardiopulmonary function and overall fitness, while resistance exercise is highly effective against sarcopenia.^[31] Combining these 2 forms of exercise can significantly improve hepatic complications and prognosis.^[8]

In mouse models focusing on liver diseases, physical exercise interventions primarily involve aerobic exercises such as treadmill running,^[32] voluntary wheel running,^[33] and swimming.^[34] Resistance exercise models include weightlifting,^[35] high-intensity interval exercise (HIIT),^[36] and ladder climbing.^[37] These exercise models offer valuable insights into the effects of physical exercise on liver health, metabolism, and the potential for treating liver diseases. Understanding the impacts of different exercise modalities on liver disease is crucial for developing effective therapeutic strategies and promoting overall liver well-being.

The intensity of physical exercise is assessed through absolute and relative intensity. Absolute intensity is measured in metabolic equivalent of task (MET) units, representing the energy expended during the exercise. Moderate-intensity activities range from 3 to 5.9 MET, while vigorous-intensity activities are 6 or higher.^[38] In contrast, relative intensity assesses effort to an individual's capacity. Intensity is a critical factor when considering physical exercise for patients with liver diseases.^[39] For instance, Keating et al^[40] showed that the minimum effective dose of exercise to improve hepatic steatosis was 135 minutes of moderate-intensity aerobic exercise per week and that increasing the intensity of exercise (including HIIT approaches) had no additional benefit for hepatic steatosis. It is essential to tailor physical exercise considering absolute and relative intensity to maximize benefits based on individual fitness levels.

Determining the appropriate exercise duration is equally important. Our group found that liver ischemia/reperfusion (I/R) injury decreased with exercise duration and was lowest after 4 weeks of preoperative exercise compared with sedentary mice. However, extended preoperative exercise periods beyond 4 weeks up to 16 weeks offered no further hepatic protection.^[32] Moreover, for those in poor physical condition, the recommendation is to initiate exercise with 20-minute sessions and progressively increase the duration by 5–10 minutes every 1–2 weeks.^[41]

In conclusion, careful selection of physical exercise modality, intensity, and duration is essential for effectively managing liver diseases. A personalized physical exercise plan tailored to the individual's health needs and capabilities is recommended to ensure improved adherence and optimal health outcomes.^[29,42]

THE MECHANISM OF PHYSICAL EXERCISE IN PROTECTING AGAINST LIVER DISEASES

Improve insulin sensitivity

Physical exercise effectively improves insulin sensitivity, which is critical to regulating glucose and lipid

TABLE 1 Molecular pathways involved in the improvement of hepatic insulin sensitivity by physical exercise

References	Exercise modeling	Exercise intervention	Molecular pathways
Marinho et al ^[45]	Aerobic exercise	Swimming, 60 min per day, 5 d per week for 8 wks	Improved the APPL1/TRB3/Akt signaling pathway
Tsuzuki et al ^[46]	Aerobic exercise	Voluntary wheel running for 20 wks	Suppressed the iNOS expression and S-nitrosylation of Akt
Zhang et al ^[47]	Aerobic exercise	Swimming, 60 min per day, 5 d per week for 8 wks	Reduced the NFE2/miR-423-5p and increased FAM3A/ATP/Akt pathway
Wang et al ^[48]	Aerobic exercise	Treadmill running, 60 min per day, 5 d per week for 10 wks	Improved the TLR-4-mediated ERK/AMPK signaling pathway
Zhang et al ^[49]	Aerobic exercise	Swimming, 60 min per day, 5 d per week for 8 wks	Suppressed hepatic TAK1/JNK/IRS1 signaling
Castaño et al ^[50]	Aerobic exercise	Treadmill running, 60 min per day, 3 d per week for 5 wks	Increased the miR-133a, miR-133b, and decreased FoxO1
Diniz et al ^[51]	Aerobic exercise	Treadmill running, 60% of maximum speed, 7 d per week for 8 wks	Improved the AMPK-PPAR- α /PPAR- γ signaling pathway
da Cruz Rodrigues, et al ^[52]	Resistance exercise	20 climbing series per day, 5 d per week for 3 wks	Increased the IRS-1/2 and AKT tyrosine phosphorylation
Zhang et al ^[53]	Aerobic exercise	Swimming, 60 min per day, 5 d per week for 8 wks	Decreased the ASK1 phosphorylation and improved JNK1/IRS1/Akt signaling pathway
Vieira et al ^[54]	Aerobic exercise	Treadmill running, 30–60 min per day, 7 d per week for 10 wks	Improved the IRS1/AKT signaling pathway

Abbreviations: Akt, protein kinase B; AMPK, AMP-activated protein kinase; APPL1, adaptor protein, phosphotyrosine interacting with PH domain and leucine zipper 1; ASK1, apoptosis signal-regulated kinase 1; ERK, extracellular signal-regulated kinase; FAM3A, family with sequence similarity 3 member A; FoxO1, forkhead box O1; IRS1, insulin receptor substrate 1; JNK1, c-Jun N-terminal kinase 1; PPAR- α , peroxisome proliferator-activated receptor alpha; PPAR- γ , peroxisome proliferator-activated receptor gamma; TAK1, TGF β -activated kinase 1; TLR-4, toll-like receptor 4.

metabolism.^[43] Insulin resistance is characterized by impaired responsiveness to insulin, leading to elevated insulin levels in the liver. This promotes MASLD through mechanisms that increase lipids' synthesis and accumulation. High levels of insulin resistance are a significant predictor of MASLD.^[44]

The positive effect of exercise on insulin sensitivity is attributed to its modulation of molecular pathways associated with insulin resistance (Table 1). For instance, Wang and colleagues found that treadmill running for 60 minutes per day, 5 days per week for 10 weeks, significantly improved the function of islet β -cells and effectively reduced insulin resistance in mice. These improvements were associated with the activation of the toll-like receptor 4 (TLR-4)-mediated extracellular signal-regulated kinase/AMP-activated protein kinase (AMPK) signaling pathway, which plays a crucial role in regulating insulin sensitivity and glucose metabolism.^[48] The research conducted by Zhang and colleagues demonstrated that 60 minutes per day, 5 days per week for 8 weeks of swimming training effectively improved hepatic insulin resistance induced by a high-fat diet (HFD) in male Sprague-Dawley rats through TGF- β -activated kinase 1-dependent signaling. The improvement in insulin sensitivity was attributed to multiple factors, including the enhancement of proteins ubiquitin-specific protease 4, dual-specificity phosphatases 14, and a reduction in the phosphorylation of critical signaling molecules such as TGF- β -activated

kinase 1, c-Jun N-terminal kinase 1, and insulin receptor substrate 1.^[49] Furthermore, this beneficial effect was associated with reduced apoptosis signal-regulated kinase 1 phosphorylation.^[53] Similarly, studies by Marinho et al and Zhang et al^[45,47] indicate that an identical regimen of swimming enhances insulin signaling through adaptor protein, phosphotyrosine interacting with PH domain and leucine zipper 1/protein kinase B (Akt) signaling pathway and the family with sequence similarity 3 member A/ATP/Akt pathway, respectively. AKT signaling plays a crucial role in insulin sensitivity, and disturbances in AKT signaling pathways may result in insulin resistance.^[46,52,54] Moreover, Diniz and colleagues have shown that treadmill running for 60 minutes per day, 5 days per week for 8 weeks, can attenuate the progression of HFD-induced hepatic steatosis and inflammation. This improvement is achieved through activating the AMPK-peroxisome proliferator-activated receptor alpha (PPAR- α) signaling pathway and PPAR-gamma (PPAR- γ) signaling pathways, leading to the amelioration of insulin resistance in obese mice.^[51]

HIIT has been shown to reduce insulin resistance and hepatic glucose production. Castaño et al^[50] revealed that HIIT can alter the microRNA profile of circulating exosomes in mice. This alteration leads to increased expression of miR-133a and miR-133b in plasma, which, in turn, downregulates the insulin-regulating transcription factor forkhead box O1 in the

liver, ultimately contributing to the attenuation of insulin resistance. These findings underscore exercise-induced molecular adaptations' significance in alleviating hepatic insulin resistance.

Regulate lipid homeostasis

The liver is vital in regulating lipid homeostasis and is the primary organ for fatty acid metabolism.^[55] Excess glucose ingested is converted into fatty acids by hepatocytes through a series of enzymatic reactions collectively known as *de novo* lipogenesis.^[56] These newly synthesized fatty acids can be stored in the liver as triglycerides or transported to adipose tissue for long-term storage. Failure to transport or use these fatty acids promptly can result in their accumulation in the liver, leading to the development of metabolic dysfunction-associated steatohepatitis (MASH).

Physical exercise involves the complex physiological process significantly affecting lipogenesis and lipolysis (Figure 1). Increasing energy expenditure is one of the primary mechanisms through which exercise influences this process.^[57] During aerobic exercise, lipase enzymes in the body are activated, leading to the hydrolysis of triglycerides into glycerol and free fatty acids (FFAs).^[58] Hormones, such as epinephrine and norepinephrine, further stimulate lipolysis.^[58] This process increases the release of FFAs into the bloodstream, serving as an energy source. Hepatic lipolysis is a complex metabolic process that relies on several critical regulatory enzymes to control the release of fatty acids from stored triglycerides. In addition, fatty acid oxidation and autophagy are involved in this intricate cellular mechanism.^[59] Physical exercise can decrease *de novo* lipogenesis and increase the fatty acid β -oxidation and autophagy. In addition, exercise can further regulate lipid metabolism by enhancing insulin sensitivity.^[59]

AMPK-dependent pathway plays a central role in exercise-mediated lipid metabolism. Zou et al^[60] demonstrated that swimming exercise for 4 hours per day, 5 days per week for 12 weeks decreased *de novo* lipogenesis and promoted liver lipophagy and fatty acid β -oxidation by activating AMPK/Sirtuin 1 signaling in MASLD Zebrafish. Qian et al^[61] noted that swimming training for 45 minutes per day, 5 days per week for 8 weeks induced hepatoprotection in HFD-fed mice, which was achieved by inhibiting the upregulation of hydroxy-3-methylglutaryl-CoA synthase 2 (HMGCS2), which serves as the key rate-limiting enzyme in ketogenesis. Another study showed that treadmill running for 60 minutes per day, 7 days per week for 12 weeks induced hepatic adenosine deaminases acting on RNA 2, which protect against lipogenesis during MASLD by decreasing the level of miR-34a.^[62] In addition, exercise regulates hepatic lipid metabolism by modulating gut microbiota.^[63]

Skeletal muscle is critical in absorbing and using FFAs for energy during exercise.^[64] Physical exercise enhances FFA utilization by skeletal muscle and promotes mitochondrial energy production, leading to more efficient utilization of FFAs. Exercise improves muscle cells' responsiveness to insulin, leading to more efficient absorption and utilization of glucose for energy production.^[65] This process, coupled with the indirect effects on FFA metabolism, contributes to the burning of stored fat. Consequently, physical exercise improves the body's overall energy metabolism efficiency.^[66]

Immunomodulation

Hepatic inflammation is a critical factor in the development and progression of liver diseases.^[67] Physical exercise is pivotal in regulating the hepatic inflammatory response by modulating key cytokines and signaling pathways,^[68] which emerges as a potent modulator of immune cell infiltration within the liver and the tumor immune microenvironment,^[69–71] effectively reducing hepatic inflammatory response (Figure 2).

In the acute hepatic injury model, our group has shown that 60 minutes per day, 5 days per week for 4 weeks of aerobic preoperative exercise regimen significantly attenuates liver injury and inflammation from ischemia and reperfusion in mice. We found that exercise specifically drives KCs toward an anti-inflammatory phenotype with trained immunity through metabolic reprogramming.^[32] In the context of acute liver injury induced by *N*-galactosamine and lipopolysaccharide, a median of 45 days of voluntary long-distance running has been found to alter the intrahepatic immunophenotype, which reduces the number of intrahepatic CD4+ T cells, B lymphocytes, and macrophages, thereby leading to a change of hepatic microenvironment that is less susceptible to acute liver injury in mice.^[72]

In MASH, treadmill training for 60 minutes per day, 5 days per week for 12 weeks has shown the capacity to attenuate hepatic inflammation, liver steatosis, and fibrosis by inhibiting the hepatic accumulation of bone marrow-derived macrophages and PD-1+ CD8+ T cells in mice.^[73] In addition, the study by Cai and colleagues revealed that swimming training for 60 minutes per day, 7 days per week for 12 weeks, upregulated the expression of key markers associated with insulin resistance, including PPAR- γ , carnitine palmitoyl transferase-1, and medium-chain acyl-CoA dehydrogenase, ultimately leading to improved insulin sensitivity in HFD-induced MASLD mice.^[74] A randomized clinical trial including 60 minutes per day, 3–5 times per week for the 12-week aerobic exercise regimen has been shown to elevate the levels of the apoptotic marker CD95 in mucosa-associated invariant T cells, both in the blood and within the liver in patients of MASLD. In addition,




FIGURE 1 Physical exercise on hepatic lipogenesis and lipolysis. Physical exercise suppresses hepatic lipogenesis and promotes liver triglyceride breakdown, enhancing lipolysis. Abbreviations: ACC, acetyl-CoA carboxylase; ATGL, adipose triglyceride lipase; FA, fatty acid; FAS, fatty acid synthase; FFA, free fatty acid; HSL, hormone-sensitive lipase; SCD1, stearoyl-CoA desaturase 1; SIRT1, silent information regulator T1; SREBP-1, sterol regulatory element-binding protein 1; TG, triglyceride.

this exercise regimen leads to decreased intrahepatic mucosa-associated invariant T cells and notable histological improvements.^[75] Exercise also hinders the activity of hepatic bone marrow-derived macrophages, thereby reducing the accumulation of inflammatory macrophages stemming from monocytes and bone marrow precursors in mice.^[71] Moreover, a specific subset of KC known as CD206+ ESAM+ KC has been identified to promote oxidative stress through the scavenger receptor CD36, a pivotal factor in MASH progression.^[76] Treadmill running for 60 minutes per

day, 5 days per week for 16 weeks, can reduce the expression of CD36 in KC,^[77] thereby decelerating the progression of MASH by altering the phenotype of these cells in mice.

Through targeted alterations in hepatic mRNA expression, exercise effectively suppresses the production of proinflammatory cytokines, including IL-6, IL-1 β , TNF- α , and monocyte chemoattractant protein-1,^[78] and elevates the expression of anti-inflammatory mediators, such as IL-10.^[79] Remarkably, this suppression is accompanied by a reduction in infiltrating macrophages,




FIGURE 2 Physical exercise on liver/tumor immune microenvironment in liver diseases. Physical exercise inhibits the infiltration of proinflammatory immune cells into the liver in benign liver diseases (left) while recruiting antitumor immune cells to both the liver and tumor microenvironment in malignant liver diseases (right).

distinctly mitigating immune cell-driven inflammation characteristic of MASH.^[71] In addition to immunomodulatory effects, push-ups and set squats for 20–30 minutes per day, 3 days per week for 12 weeks, reduce hepatocyte apoptosis, specifically involving cytokeratin 18 in patients with MASLD.^[80] In addition, treadmill running for 50 minutes per day, 5 days per week for 12 weeks, effectively inhibits the hepatic toll-like receptor 4-mediated NF- κ B pathway through apolipoprotein A5, further enhancing its protective effects on the liver.^[81] The level of physical exercise is inversely related to hepatic fibro inflammation, as measured by iron-corrected T1, underscoring its pivotal role in maintaining liver health and mitigating inflammation-associated liver diseases.^[82]

Natural killer cells are vital components of the innate immune system, primarily responsible for immunosurveillance and eliminating cells with low expression of major histocompatibility complex class I.^[83] IL-15 is a critical activator of NK cells and enhances their antitumor responses.^[84] A cross-sectional study of 133 medical students indicated that regular endurance training increases serum IL-15 expression.^[85] Also, treadmill running for 60 minutes per day, 5 days per week for 12 weeks, increases the expression of NK cell ligands retinoic acid early inducible gene 1 in the liver tissue of tumor-bearing mice, thereby enhancing the cytotoxic capabilities of NK cells.^[86] The efficacy of T cells in recognizing and eliminating cancer cells is paramount in preventing tumor growth and forms the basis of current immunotherapy.^[87] During exercise, higher levels of metabolites such as lactate released into the bloodstream from skeletal muscle enhance the effector profile of CD8+ T cells. In response to exercise, activated CD8+ T cells in mice adapt their central carbon metabolism. Transferring trained mouse CD8+ T cells into untrained tumor-bearing animals produces a more potent antitumor effect.^[88]

EXERCISE IN LIVER I/R INJURY

Liver I/R injury is a complex and multifaceted acute inflammation liver disease that occurs when the blood supply to the liver is temporarily interrupted and restored.^[89] This process may occur in various clinical situations, such as liver surgery, trauma, or liver transplantation.^[90] Ischemia is the initial phase that leads to hypoxia in liver tissue, resulting in cellular damage and metabolic disturbances. Subsequently, during the reperfusion phase, tissue damage is exacerbated due to the release of ROS and inflammatory mediators. This phenomenon leads to severe hepatic dysfunction and, in extreme cases, multiorgan failure.^[91] The pathophysiology of hepatic I/R injury involves a complex interplay of various cellular and molecular mechanisms.^[92] It is, therefore, an essential area of

research for developing therapeutic strategies to mitigate its harmful effects.

Exercise activates AMPK, an intracellular sensor that plays a role in metabolic reprogramming.^[93] The AMPK activator 5-aminoimidazole-4-carboxamide ribonucleotide enhances ischemic tolerance and fatty liver regeneration after hepatic I/R in diet-induced hepatic steatosis mice.^[94] Another key player in the liver I/R process is irisin, a hormone-like molecule released from muscle tissue during physical exercise.^[95] Irisin's ability to attenuate hepatic I/R injury by binding to the α V β 5 integrin receptor in hepatocytes has significant implications.^[96] Administering irisin at the onset of reperfusion has been shown to improve mitochondrial function, reduce oxidative stress, and alleviate endoplasmic reticulum stress in HFD mice, ultimately mitigating liver injury.^[97]

In liver transplantation, exercise plays a distinct role in enhancing the overall well-being of individuals undergoing this life-transforming procedure. Both pretransplant and posttransplant exercise training are recognized as safe and highly beneficial.^[98,99] A meta-analysis including 8 RCTs indicates that exercise training improves liver transplantation candidates' physical function and health-related quality of life, furthering their cardiorespiratory and muscle health.^[100] It optimizes their posttransplant recovery, ultimately contributing to an improved long-term quality of life following transplantation.

EXERCISE IN CARDIOGENIC LIVER DISEASE

Other conditions leading to liver injury, including cardiogenic liver disease triggered by myocardial infarction, may benefit from exercise. Treadmill running for 60 minutes per day, 5 days per week for 6 weeks has the potential to effectively mitigate liver injury by promoting the anti-inflammatory phenotype of hepatic macrophages, attenuating myocardial infarction-induced hepatic inflammation, enhancing the expression of fibronectin type III domain-containing protein 5 protein, and activating the PI3K/protein kinase B signaling pathway in the liver of myocardial infarction-afflicted mice.^[101]

EXERCISE IN MASLD

MASLD is indeed one of the most prevalent chronic liver disorders worldwide, affecting up to 25% of the global population.^[102] This condition encompasses a range of histological features, including hepatic steatosis, hepatocyte ballooning, hepatic lobular inflammation, and liver fibrosis. Approximately 4% of individuals with simple steatosis and over 20% of those with MASH

are expected to develop cirrhosis during their lifetime.^[103] MASLD is a complex multifactorial disease whose exact pathogenesis is not fully understood. Insulin resistance and liver steatosis may represent the first hit for the liver.^[104] However, it is important to note that steatosis, or fat accumulation in the liver, is typically an early event in MASLD and does not necessarily progress to MASH. Additional stresses are required to induce MASH onsets, such as oxidative stress, lipotoxicity, inflammation, subsequent stimulation of hepatocyte death, tissue regeneration, and fibrogenesis.^[104]

A large amount of evidence derived from clinical studies of MASLD supports the vital role of physical exercise in improving histological parameters and decelerating the progression of the disease. This includes its ability to enhance insulin resistance, reduce hepatic steatosis, and attenuate hepatic inflammation.^[30,105] For instance, a randomized clinical trial conducted by Charatcharoenwitthaya et al^[30] reported moderate-intensity aerobic exercise or resistance training for 60 minutes per day, 5 times per week for 12 weeks, coupled with dietary modifications equally reduced intrahepatic fat and substantially improved underlying insulin resistance in individuals diagnosed with MASLD. It is worth noting that the effectiveness of exercise in combating hepatic insulin resistance varies based on the duration of the intervention. Hari et al^[106] discovered that short-term (7 d) treadmill training consisting of 60 minutes per day at 80%–85% of maximal heart rate increased peripheral insulin sensitivity by 34% in patients with radiographically confirmed MASLD (>5% intrahepatic lipid content). Another meta-analysis that included 26 randomized clinical trials showed that short-term exercise interventions, lasting <12 weeks, have shown efficacy in reducing specific insulin resistance markers such as fasting blood glucose and the homeostatic model assessment of insulin resistance.^[107] Interestingly, an RCT indicated that combining moderate-intensity aerobic exercise (60 min per day, 5 times per week for 12 wks) with alternate-day fasting has been shown to reduce insulin resistance in adults with MASLD. However, this combined approach does not significantly enhance insulin resistance improvements compared to exercise alone.^[108] These clinical findings underscore the therapeutic potential of physical exercise as a potent tool in combatting insulin resistance in MASLD.

Several clinical studies demonstrate that physical exercise reduces IHTG levels, which is independent of weight loss.^[24,108–110] A systematic review has compared the therapeutic effects of aerobic and resistance exercises on hepatic steatosis in MASLD. They found that both aerobic and resistance exercise improved hepatic steatosis to a similar extent, with common parameters of duration, frequency, and training period (40–45 min/session, 3 times/week for 12 wks).^[7] However, resistance exercise may be more feasible for patients with MASLD with poor cardiorespiratory

fitness because it improves MASLD with lower intensity and less energy consumption.^[7] In addition, a meta-regression analysis from 17 studies (373 exercising participants) revealed a significant negative correlation between the duration of exercise and the reduction in IHTG ($\beta = -0.27$ [95% CI: -0.35 to -0.19], $p < 0.001$). This suggests that as the duration of exercise increases, the reduction in IHTG becomes more pronounced. Specifically, compared with shorter high-intensity exercise, continuous moderate-intensity protocols commonly exhibit a more significant decrease in IHTG in those patients with MASLD.^[111] The study revealed no statistically significant distinctions among the 3 levels of aerobic exercise regimens regarding reducing intrahepatic lipid content. The results demonstrated that even minimal engagement in exercise led to a noticeable reduction in intrahepatic lipid levels.^[111] These findings highlight the importance of developing individualized sustainable exercise interventions that may yield the most significant benefits for IHTG reduction in patients with MASLD.

Although the FDA has approved Rezdiffra (resmetirom) as the first drug treatment for patients with MASH with moderate to advanced liver scarring (fibrosis), lifestyle modification, including physical exercise, remains generally recommended as a first-line therapy for all patients with MASLD.^[112] In line with this recommendation, the American Gastroenterological Association recommends that all patients with MASLD should engage in regular physical activity with a target of 150–300 minutes of moderate-intensity (3–6 MET) or 75–150 minutes of vigorous-intensity (more than 6 MET) aerobic exercise per week. Resistance training can be complementary to aerobic exercise but not a replacement (Figure 3).^[112]

EXERCISE IN PORTAL HYPERTENSION

Portal hypertension (PH) is defined as abnormally high blood pressure within the portal vein system, primarily caused by cirrhosis and other portal vein obstructions, such as PVT.^[113] PH can lead to serious complications such as ascites, variceal bleeding, splenomegaly, and an elevated risk of spontaneous bacterial peritonitis or other infections, HE, hepatorenal syndrome, and liver failure.^[114] The gold standard for the diagnosis of PH is the HVP. An HVP value exceeding 5 mm Hg indicates the presence of PH, while a value above 10 mm Hg denotes clinically significant portal hypertension. This latter threshold is considered clinically significant because it marks the point at which complications of PH begin to develop.^[115] Treatment of PH focuses on reducing portal pressure, preventing complications, and managing symptoms. This approach mainly includes the medication (nonselective beta-blockers), the TIPS, and pericardial devascularization.^[116]



FIGURE 3 Aerobic exercise (upper) and resistance exercise (lower) are recommendations for normal-condition patients with MASLD, cirrhosis, and liver cancer. Abbreviation: MASLD, metabolic dysfunction–associated steatotic liver disease.

Research on the impact of physical exercise on PH is limited and has historically raised concerns about the potential for exacerbating complications such as HE or bleeding after exercise.^[117] Initial studies indicated that 8–10 minutes of cycling at 30% and 50% of peak workload increased HVPG.^[117] In an RCT, 40 patients with compensated cirrhosis of Child-Pugh A were recruited to evaluate the effects of a 12-week home-based exercise program. The regimen included moderate-intensity aerobic/isotonic continuous training exercises lasting 40 minutes per session, conducted at least 4 times weekly. This study reported no adverse events such as gastrointestinal bleeding, ascites, or musculoskeletal injuries, yet it did not find any beneficial impact on PH.^[118] Conversely, another pilot randomized clinical trial involving 29 patients with cirrhosis and PH (average HVPG > 10 mm Hg) who participated in a 14-week supervised exercise program showed significant benefits. The regimen, which consisted of physical exercises performed for 40 minutes, 3 times a week at an intensity of 12–14 on the Borg Rating of Perceived Exertion scale, led to an average reduction in HVPG of 2.5 mm Hg, with no episodes of variceal bleeding or HE observed.^[119] In addition, a prospective, multicenter, uncontrolled pilot study focused on the effects of an intensive lifestyle intervention on patients with compensated cirrhosis and PH (HVPG \geq 6 mm Hg). Participants were engaged in a 16-week intensive lifestyle modification program, which included a personalized hypocaloric normoproteic diet and 60 minutes of supervised physical activity (10 min of warm-up, 40 min of aerobic and strength exercising routine, and 10 min of cooling down) each week. Results showed a significant reduction in HVPG, decreasing from an average of 13.9 ± 5.6 mm Hg at baseline to 12.3 ± 5.2 mm Hg after intervention, with 42% of participants achieving an HVPG decrease of \geq 10%. However, the study's design does not confirm whether the observed reductions in PH were due to exercise, diet, or a combination of both.^[120]

Current research on PH is primarily focused on cirrhosis-induced cases, with other causes of PH not being well studied. In addition, there is a lack of comprehensive data on the appropriate intensity, duration, and type of exercise required for effective treatment and prevention of PH, and thus no consensus has been reached in this area.

EXERCISE IN CIRRHOSIS

Cirrhosis, primarily caused by MASLD, viral hepatitis, and alcohol consumption, accounted for 2.4% of global deaths in 2019.^[5,121] Cirrhosis manifests as extensive fibrosis, resulting in nodular changes in liver structure. Portal hypertension, in combination with cirrhosis, can lead to severe complications such as ascites, variceal bleeding, and HE.^[113] The liver's ability to synthesize and metabolize is impaired in cirrhosis, and the risk of HCC is elevated.^[3]

Physical exercise has been demonstrated to diminish hepatic inflammation and fibrosis by influencing different signaling pathways. The mechanisms are primarily associated with the regulation of several related signaling pathways, including the AMPK signaling pathway,^[122] the HMGCS2-regulated Wnt3a/ β -catenin pathway,^[61] monoacylglycerol O-acyltransferase 1 pathway,^[123] and MD2-toll-like receptor 4 pathway,^[124] etc. In addition, these mechanisms encompass the inhibition of NLR family pyrin domain-containing 3 inflammasome activation,^[125] modulation of myeloid cell monocytes infiltration in the liver,^[126] and restoration of gut microbiota.^[127]

Numerous clinical studies have highlighted the advantageous effects of exercise for individuals in the early stages of cirrhosis. Aamann et al^[31,128] found that a regimen of resistance training for 1 hour, 3 times weekly for 12 weeks, not only diminished the risk of initial hospitalization and mortality among patients with cirrhosis classified as Child-Pugh class A/B 3 years

post-trial entry but also resulted in increased muscle strength and size. Sirisunhirun et al^[118] implemented a 12-week program of aerobic moderate-intensity continuous training exercises, lasting 40 minutes per session, at least 4 times a week, in patients with compensated cirrhosis. This program significantly ameliorated the fatigue domain of the quality-of-life index without inducing adverse reactions. However, no benefits were observed in thigh muscle mass, liver stiffness, or spleen stiffness. Moreover, a meta-analysis of RCTs by Kawaguchi et al discovered that a median protocol of aerobic exercise for 30 minutes per session, 3 times a week for 12 weeks, in combination with a resistance exercise protocol of 60 minutes per session, 3 times a week for 10 weeks, significantly reduced the incidence of serious events in patients with liver cirrhosis and enhanced the 6-minute walk distance, an independent prognostic factor for these patients.^[8,129]

Given the heterogeneity of disease progression and variable clinical presentations in cirrhosis, standardized exercise guidelines, particularly for resistance exercise, remain elusive.^[130] Nonetheless, there are established recommendations for the prescription of aerobic exercise in patients with cirrhosis. Before initiating an exercise regimen in patients with cirrhosis, a comprehensive medical assessment is imperative. This assessment should encompass an evaluation of cardiopulmonary function, concurrent pathologies, and musculoskeletal constraints. Following this, it is crucial to delineate exercise objectives and modalities, adhering to the FITT (Frequency, Intensity, Time, and Type) principle, tailored to the patient's unique health condition. The clinical practice recommendations for patients with cirrhosis advocate for the engagement in physical exercise of mild to moderate intensity for 30–60 minutes per session, 5 days per week (cumulatively ≥ 150 min per week) through an activity such as walking or cycle ergometry. The intensity should be calibrated to 30%–40% of heart rate reserve or within a range of 10–14 on the Borg scale (Figure 3).^[41]

EXERCISE IN LIVER CANCER

Liver cancer represents a significant global health challenge and the third leading cause of cancer death worldwide, accounting for ~7.8% of total cancer deaths in 2022.^[131] The impending burden of liver cancer is expected to affect over 1 million individuals by 2025.^[132] Notably, physical exercise is emerging as a particularly promising strategy. Studies suggest that physical exercise might play a role in various aspects, including reducing the risk of developing liver cancer,^[133–135] controlling liver cancer progression^[111], enhancing efficacy, and reducing the side effects of anticancer treatments.^[136,137] This growing evidence suggests that physical exercise could be an effective component in combating liver cancer.

The accumulating epidemiological evidence supports the potential protective role of physical exercise against liver cancer, particularly HCC. In the United States, a study revealed a moderate inverse correlation ($r = -0.40$, $p = 0.004$) between state-level physical exercise and the incidence of HCC.^[9] This is further supported by comprehensive research from over 44 million participants across the United States and Europe, revealing that higher levels of leisure-time physical exercise reduce liver cancer risk by over 20%.^[138] Another cohort study quantifying the dose-response relationship between leisure-time physical exercise and liver cancer showed that engagement in 7.5–15 hours per week of leisure-time physical exercise was associated with an 18%–27% lower risk of liver cancer.^[133] However, the study also identified a nonlinear association between the risk of liver cancer and activity levels, implying a limit beyond which additional exercise does not further decrease liver cancer risk. This finding raises important questions about the optimal amount and intensity of physical exercise for liver cancer prevention. While these studies are groundbreaking, they are limited by their failure to adjust for other potentially important confounding factors, including viral hepatitis, MASLD, cirrhosis, and alcohol consumption. A prospective multinational cohort study further explored this association, finding physical exercise associated with a lower risk of liver cancers. This relationship persisted even after adjusting for liver cancer risk factors like alcohol consumption, smoking, waist circumference, or body mass index.^[139] Moreover, a meta-analysis provided further evidence, demonstrating a 23% reduction in liver cancer risk and a 19% decrease in mortality among individuals who engaged in moderate physical exercise, reinforcing the potential protective impact of exercise against liver cancer.^[140]

The beneficial effects of physical exercise on liver cancer can be attributed to various systemic and local biological mechanisms. Obesity, an independent risk factor for mortality in primary liver cancer, can be significantly mitigated by physical exercise, reducing the associated risk.^[117] In addition, in patients with HCC, a decline in skeletal muscle mass is significantly associated with severe adverse events from chemotherapy, low tolerability, increased tumor recurrence, and all-cause mortality.^[141] Significantly, exercise intervention has improved skeletal muscle strength in these patients.^[142] This underscores the effective role of exercise in enhancing the management of liver cancer, potentially improving treatment outcomes and survival rates for affected individuals. The benefits of exercise on liver cancer can be elucidated through 3 fundamental mechanisms. First, exercise regulates critical signaling pathways and modulatory proteins crucial in HCC progression. This includes the suppression of TGF- β ,^[143] downregulation of dynamin-related protein 1,^[144] activation of p53,^[145] and modulation of the AMPK/mTOR

signaling pathway.^[146] Second, exercise enhances anti-tumor immunity by increasing the CD8+ T-cell infiltration into the tumor microenvironment, creating a hostile environment for tumor growth and progression within the liver.^[147] Third, exercise induces metabolic reprogramming of tumor microenvironment,^[148] forming an exercise-induced metabolic shield unfavorable for tumor cell colonization. This metabolic reprogramming underscores the role of exercise in reshaping the metabolic landscape of the tumor microenvironment to impede cancer progression and metastasis. Beyond these specific mechanisms, exercise also influences other critical factors, such as angiogenesis and oxidative stress, which can affect tumor growth and survival.^[149] However, it is essential to acknowledge that current research on these mechanisms remains limited, necessitating further empirical investigation to substantiate these findings and fully elucidate their implications in the context of liver cancer management.

The American Society of Clinical Oncology and the American College of Sports Medicine actively endorse the integration of regular exercise into both cancer treatment regimens and posttreatment recovery.^[150,151] During the second roundtable conference held in 2018, an updated consensus on exercise programs for cancer survivors, including those with liver cancer, was presented. The most effective exercise prescription for addressing health-related outcomes caused by cancer diagnosis and treatment includes engaging in moderate-intensity aerobic training at least 3 times per week, for at least 30 minutes each session, for at least 8–12 weeks. In addition to aerobic training, incorporating resistance training at least twice a week, with at least 2 sets of 8–15 repetitions, where each repetition is performed with at least 60% of the maximum weight that can be lifted once, appears to offer similar benefits (Figure 3).^[152] This endorsement highlights the recognized importance of physical activity in enhancing the overall health and quality of life for individuals undergoing cancer therapy and those in the recovery phase after treatment. Further evidence is needed to formulate individualized and realistic exercise programs, including the type, intensity, and duration of exercise, ensuring their effectiveness and feasibility for patients with liver cancer.

PHYSICAL EXERCISE INDIRECTLY IMPROVES LIVER DISEASES THROUGH OTHER ORGANS

Physical exercise offers systemic benefits that improve health and disease resistance across various organs. In the brain, physical exercise enhances cognitive functions through multiple mechanisms: it increases the generation of new neurons in the hippocampus through brain-derived neurotrophic factor,^[153] suppresses

neuroinflammation by upregulating amyloid- β peptide (A β) transporter activity to clear A β ,^[154] and promotes neuroplastic changes by altering the synaptic structure and function.^[155] Cardiovascular health benefits significantly from physical exercise, which promotes myocardial mitochondrial biogenesis and autophagy through the peroxisome proliferator-activated receptor gamma coactivator 1-alpha.^[156] In addition, exercise attenuates myocardial inflammation and apoptosis by regulating the apelin peptide jejunum/signal transducer and activator of transcription 3 signaling pathway,^[157] and it improves heart function after myocardial infarction through the AMPK signaling pathway.^[158] For muscles, physical exercise increases the utilization of glucose and fatty acid,^[159] promotes the secretion of myokines that facilitate communication with other organs such as adipose tissue, liver, and pancreas,^[160] and reduces the risk of sarcopenia by maintaining muscle mass and strength.^[161] Moreover, physical exercise contributes to promoting adipose tissue utilization,^[162] boosts the anti-inflammatory gene expression in adipose tissue,^[163] and enhances paracrine and endocrine functions of adipose tissue by increasing the production of beneficial adipokines and extracellular vesicles.^[164] Exercise also benefits gut health by regulating the gut microbiota, increasing butyrate production,^[165] and restoring intestinal barrier integrity.^[166]

Physical exercise indirectly promotes liver health through its effects on other organs. For instance, physical exercise stimulates the brain to secrete IGF-1,^[167] which alleviates hepatic histologic lesions in MASH.^[168] In addition, exercise enhances cardio-pulmonary efficiency, which significantly improves overall well-being and quality of life in adult recipients of liver transplant.^[100] Physical exercise also triggers the release of myokines, such as Irisin, which can inhibit hepatic inflammation by competitively binding with MD2 to improve MASLD.^[169] Moreover, exercise helps reduce the accumulation of adipose tissue, addressing a key factor in the development of MASLD.^[109] It also regulates the gut microbiota, providing therapeutic benefits for liver diseases like MASLD and cirrhosis.^[170,171] Through these indirect mechanisms, physical exercise substantially supports liver health, highlighting its wide-ranging impact.

CONCLUSIONS AND FURTHER PERSPECTIVE

In summary, the existing literature demonstrates the impact of physical exercise on liver health, as it effectively improves critical factors such as insulin resistance, hepatic fatty acid metabolism, and the modulation of inflammatory cascades. These findings support the current guidelines recommending tailored exercise programs to enhance long-term adherence

and promote an active lifestyle among patients with liver diseases. However, there remains a pressing need for multicenter clinical trials to establish exercise norms for different disease stages, thereby further enriching our understanding of the role of exercise in liver disease management.

AUTHOR CONTRIBUTIONS

Yunwei Zhang and Chunyan Cao wrote the manuscript and prepared figures. Chaofan Li, Russell G. Witt, Hai Huang, and Allan Tsung helped with manuscript revision. Hongji Zhang provided supervision, manuscript review, and editing.

FUNDING INFORMATION

This work was supported by the National Institutes of Health R01-CA214865 and R01-GM95566 to Allan Tsung, National Institutes of Health R01-GM137203 to Hai Huang, and state funds within the UVA Comprehensive Cancer Center “IDEA-Cancer pilot award” and “Cancer Therapeutics (CRX) pilot award” to Hongji Zhang.

CONFLICTS OF INTEREST

The authors have no conflicts to report.

ORCID

Yunwei Zhang  <https://orcid.org/0000-0002-7609-8492>

Chunyan Cao  <https://orcid.org/0000-0002-7698-1381>

Chaofan Li  <https://orcid.org/0000-0002-5426-1498>

Russell G. Witt  <https://orcid.org/0000-0003-3747-8633>

Hai Huang  <https://orcid.org/0000-0001-7430-0119>

Allan Tsung  <https://orcid.org/0000-0002-3916-8965>

Hongji Zhang  <https://orcid.org/0000-0002-7704-4365>

REFERENCES

- Devarbhavi H, Asrani SK, Arab JP, Nartey YA, Pose E, Kamath PS. Global burden of liver disease: 2023 update. *J Hepatology*. 2023;79:516–37.
- Younossi ZM, Golabi P, Paik JM, Henry A, Van Dongen C, Henry L. The global epidemiology of nonalcoholic fatty liver disease (NAFLD) and nonalcoholic steatohepatitis (NASH): A systematic review. *Hepatology*. 2023;77:1335–47.
- Lomba R, Friedman SL, Shulman GI. Mechanisms and disease consequences of nonalcoholic fatty liver disease. *Cell*. 2021;184:2537–64.
- Bhala N, Mellinger J, Asrani SK, Shah VH. Tackling the burden of preventable liver disease in the USA. *Lancet Gastroenterol Hepatol*. 2024;9:9–10.
- Huang DQ, Terrault NA, Tacke F, Gluud LL, Arrese M, Bugianesi E, et al. Global epidemiology of cirrhosis—Aetiology, trends and predictions. *Nat Rev Gastroenterol Hepatol*. 2023;20:388–98.
- Gallo A, Dedionigi C, Civitelli C, Panzeri A, Corradi C, Squizzato A. Optimal management of cirrhotic ascites: A review for internal medicine physicians. *J Transl Int Med*. 2020;8:220–36.
- Hashida R, Kawaguchi T, Bekki M, Omoto M, Matsuse H, Nago T, et al. Aerobic vs. resistance exercise in non-alcoholic fatty liver disease: A systematic review. *J Hepatol*. 2017;66:142–52.
- Kawaguchi T, Kawaguchi A, Hashida R, Nakano D, Tsutsumi T, Kawaguchi M, et al. Resistance exercise in combination with aerobic exercise reduces the incidence of serious events in patients with liver cirrhosis: A meta-analysis of randomized controlled trials. *J Gastroenterol*. 2024;59:216–28.
- Lee YT, Wang JJ, Luu M, Tseng HR, Rich NE, Lu SC, et al. State-level HCC incidence and association with obesity and physical activity in the United States. *Hepatology*. 2021;74:1384–94.
- Bernardi M, Santini C, Trevisani F, Baraldini M, Ligabue A, Gasbarrini G. Renal function impairment induced by change in posture in patients with cirrhosis and ascites. *Gut*. 1985;26:629–35.
- Franken FH. Bed rest, diet and working capability in liver disease (author's transl). *Leber Magen Darm*. 1977;7:300–5.
- Messner M, Brissot P. Traditional management of liver disorders. *Drugs*. 1990;40(suppl 3):45–57.
- Koya S, Kawaguchi T, Hashida R, Goto E, Matsuse H, Saito H, et al. Effects of in-hospital exercise on liver function, physical ability, and muscle mass during treatment of hepatoma in patients with chronic liver disease. *Hepatol Res*. 2017;47:E22–e34.
- Lesmana CR, Inggriani S, Cahyadinata L, Lesmana LA. Deep vein thrombosis in patients with advanced liver cirrhosis: A rare condition? *Hepatol Int*. 2010;4:433–8.
- Chun HS, Lee M, Lee HA, Oh SY, Baek HJ, Moon JW, et al. Association of physical activity with risk of liver fibrosis, sarcopenia, and cardiovascular disease in nonalcoholic fatty liver disease. *Clin Gastroenterol Hepatol*. 2023;21:358–369. e312.
- Ghezal S, Clément M-A, Bosoi CR, Beauchamp R, Tremblay M, Rose CF, et al. Optimizing muscle mass: Therapeutic target to prevent experimental hepatic encephalopathy: 210. *Hepatology*. 2014;60:304A.
- Duarte-Rojo A, Ruiz-Margáin A, Montañó-Loza AJ, Macías-Rodríguez RU, Ferrando A, Kim WR. Exercise and physical activity for patients with end-stage liver disease: Improving functional status and sarcopenia while on the transplant waiting list. *Liver Transpl*. 2018;24:122–39.
- Diao X, Ling Y, Zeng Y, Wu Y, Guo C, Jin Y, et al. Physical activity and cancer risk: A dose-response analysis for the Global Burden of Disease Study 2019. *Cancer Commun (Lond)*. 2023;43:1229–43.
- Sung KC, Ryu S, Lee JY, Kim JY, Wild SH, Byrne CD. Effect of exercise on the development of new fatty liver and the resolution of existing fatty liver. *J Hepatol*. 2016;65:791–7.
- Choo YJ, Cho CW, Chang MC. Effects of supervised exercise on aerobic capacity and quality of life in patients with chronic liver disease and patients who underwent liver transplantation: A systematic review and meta-analysis. *Int J Rehabil Res*. 2022;45:1–11.
- Barb D, Repetto EM, Stokes ME, Shankar SS, Cusi K. Type 2 diabetes mellitus increases the risk of hepatic fibrosis in individuals with obesity and nonalcoholic fatty liver disease. *Obesity (Silver Spring)*. 2021;29:1950–60.
- Anstee QM, McPherson S, Day CP. How big a problem is non-alcoholic fatty liver disease? *Bmj*. 2011;343:d3897.
- Bianco A, Franco I, Curci R, Bonfiglio C, Campanella A, Mirizzi A, et al. Diet and exercise exert a differential effect on glucose metabolism markers according to the degree of NAFLD severity. *Nutrients*. 2023;15:2252.
- Zhang HJ, He J, Pan LL, Ma ZM, Han CK, Chen CS, et al. Effects of moderate and vigorous exercise on nonalcoholic fatty liver disease: A randomized clinical trial. *JAMA Intern Med*. 2016;176:1074–82.

25. Ghamarchehreh ME, Shamsoddini A, Alavian SM. Investigating the impact of eight weeks of aerobic and resistance training on blood lipid profile in elderly with non-alcoholic fatty liver disease: A randomized clinical trial. *Gastroenterol Hepatol Bed Bench*. 2019;12:190–6.
26. Smart TFF, Doleman B, Hatt J, Paul M, Toft S, Lund JN, et al. The role of resistance exercise training for improving cardiorespiratory fitness in healthy older adults: A systematic review and meta-analysis. *Age Ageing*. 2022;51:afac143.
27. Machado CLF, Botton CE, Brusco CM, Pfeifer LO, Cadore EL, Pinto RS. Acute and chronic effects of muscle power training on blood pressure in elderly patients with type 2 diabetes mellitus. *Clin Exp Hypertens*. 2020;42:153–9.
28. Moghamasi M, TaheriChadorneshin H, Papoli-Baravati SA, Teymuri A. Effects of upper-body resistance exercise training on serum nesfatin-1 level, insulin resistance, and body composition in obese paraplegic men. *Disabil Health J*. 2019;12:29–34.
29. van der Windt DJ, Sud V, Zhang H, Tsung A, Huang H. The effects of physical exercise on fatty liver disease. *Gene Expr*. 2018;18:89–101.
30. Charatcharoenwithaya P, Kuljiratitikal K, Aksornchanya O, Chaiyasoot K, Bandidnyamanon W, Charatcharoenwithaya N. Moderate-intensity aerobic vs resistance exercise and dietary modification in patients with nonalcoholic fatty liver disease: A randomized clinical trial. *Clin Transl Gastroenterol*. 2021;12:e00316.
31. Aamann L, Dam G, Borre M, Drljevic-Nielsen A, Overgaard K, Andersen H, et al. Resistance training increases muscle strength and muscle size in patients with liver cirrhosis. *Clin Gastroenterol Hepatol*. 2020;18:1179–187.e1176.
32. Zhang H, Chen T, Ren J, Xia Y, Onuma A, Wang Y, et al. Pre-operative exercise therapy triggers anti-inflammatory trained immunity of Kupffer cells through metabolic reprogramming. *Nat Metab*. 2021;3:843–58.
33. Kim YJ, Kim HJ, Lee SG, Kim DH, In Jang S, Go HS, et al. Aerobic exercise for eight weeks provides protective effects towards liver and cardiometabolic health and adipose tissue remodeling under metabolic stress for one week: A study in mice. *Metabolism*. 2022;130:155178.
34. Pi H, Liu M, Xi Y, Chen M, Tian L, Xie J, et al. Long-term exercise prevents hepatic steatosis: A novel role of FABP1 in regulation of autophagy-lysosomal machinery. *Faseb j*. 2019;33:11870–83.
35. Cui D, Drake JC, Wilson RJ, Shute RJ, Lewellen B, Zhang M, et al. A novel voluntary weightlifting model in mice promotes muscle adaptation and insulin sensitivity with simultaneous enhancement of autophagy and mTOR pathway. *Faseb j*. 2020;34:7330–44.
36. Izquierdo M, Merchant RA, Morley JE, Anker SD, Aprahamian I, Arai H, et al. International Exercise Recommendations in Older Adults (ICFSR): Expert consensus guidelines. *J Nutr Health Aging*. 2021;25:824–53.
37. Minuzzi LG, Kuga GK, Breda L, Gaspar RC, Muñoz VR, Pereira RM, et al. Short-term resistance training increases APPL1 content in the liver and the insulin sensitivity of mice fed a long-term high-fat diet. *Exp Clin Endocrinol Diabetes*. 2020;128:30–7.
38. Torres ER, Bendlin BB, Kassahun-Yimer W, Magnotta VA, Paradiso S. Transportation physical activity earlier in life and areas of the brain related to dementia later in life. *J Transp Health*. 2021;20:100992.
39. Słomko J, Zalewska M, Niemiro W, Kujawski S, Słupski M, Januszko-Giergielewicz B, et al. Evidence-based aerobic exercise training in metabolic-associated fatty liver disease: Systematic review with meta-analysis. *J Clin Med*. 2021;10:1659.
40. Keating SE, Sabag A, Hallsworth K, Hickman IJ, Macdonald GA, Stine JG, et al. Exercise in the management of metabolic-associated fatty liver disease (MAFLD) in adults: A position statement from Exercise and Sport Science Australia. *Sports Med*. 2023;53:2347–71.
41. Macías-Rodríguez RU, Ruiz-Margáin A, Román-Calleja BM, Moreno-Tavarez E, Weber-Sangri L, González-Arellano MF, et al. Exercise prescription in patients with cirrhosis: Recommendations for clinical practice. *Rev Gastroenterol Mex (Engl Ed)*. 2019;84:326–43.
42. Farrugia MA, Le Garf S, Chierici A, Piche T, Gual P, Iannelli A, et al. Therapeutic physical exercise programs in the context of NASH cirrhosis and liver transplantation: A systematic review. *Metabolites*. 2023;13:330.
43. Heiston EM, Eichner NZ, Gilbertson NM, Malin SK. Exercise improves adiposopathy, insulin sensitivity and metabolic syndrome severity independent of intensity. *Exp Physiol*. 2020;105:632–40.
44. Huang JF, Tsai PC, Yeh ML, Huang CF, Huang CI, Hsieh MH, et al. Risk stratification of non-alcoholic fatty liver disease across body mass index in a community basis. *J Formos Med Assoc*. 2020;119:89–96.
45. Marinho R, Ropelle ER, Cintra DE, De Souza CT, Da Silva AS, Bertoli FC, et al. Endurance exercise training increases APPL1 expression and improves insulin signaling in the hepatic tissue of diet-induced obese mice, independently of weight loss. *J Cell Physiol*. 2012;227:2917–26.
46. Tsuzuki T, Shinozaki S, Nakamoto H, Kaneki M, Goto S, Shimokado K, et al. Voluntary exercise can ameliorate insulin resistance by reducing iNOS-mediated S-nitrosylation of Akt in the liver in obese rats. *PLoS One*. 2015;10:e0132029.
47. Zhang Y, Wan J, Liu S, Hua T, Sun Q. Exercise induced improvements in insulin sensitivity are concurrent with reduced NFE2l3/miR-432-5p and increased FAM3A. *Life Sci*. 2018;207:23–9.
48. Wang M, Li S, Wang F, Zou J, Zhang Y. Aerobic exercise regulates blood lipid and insulin resistance via the toll-like receptor 4-mediated extracellular signal-regulated kinases/AMP-activated protein kinases signaling pathway. *Mol Med Rep*. 2018;17:8339–48.
49. Zhang Y, Wan J, Xu Z, Hua T, Sun Q. Exercise ameliorates insulin resistance via regulating TGFβ-activated kinase 1 (TAK1)-mediated insulin signaling in liver of high-fat diet-induced obese rats. *J Cell Physiol*. 2019;234:7467–74.
50. Castaño C, Mirasierra M, Vallejo M, Novials A, Párrizas M. Delivery of muscle-derived exosomal miRNAs induced by HIIT improves insulin sensitivity through down-regulation of hepatic FoxO1 in mice. *Proc Natl Acad Sci USA*. 2020;117:30335–43.
51. Diniz TA, de Lima Junior EA, Teixeira AA, Biondo LA, da Rocha LAF, Valadão IC, et al. Aerobic training improves NAFLD markers and insulin resistance through AMPK-PPAR-α signaling in obese mice. *Life Sci*. 2021;266:118868.
52. da Cruz Rodrigues KC, Martins Pereira R, Peruca GF, Torres Barbosa LW, Ramos Sant'Ana M, Rosetto Muñoz V, et al. Short-term strength exercise reduces hepatic insulin resistance in obese mice by reducing PTP1B content, regardless of changes in body weight. *Int J Mol Sci*. 2021;22:6402.
53. Zhang Y, Ye T, Zhou P, Li R, Liu Z, Xie J, et al. Exercise ameliorates insulin resistance and improves ASK1-mediated insulin signalling in obese rats. *J Cell Mol Med*. 2021;25:10930–8.
54. Vieira RFL, Muñoz VR, Junqueira RL, de Oliveira F, Gaspar RC, Nakandakari S, et al. Time-restricted feeding combined with aerobic exercise training can prevent weight gain and improve metabolic disorders in mice fed a high-fat diet. *J Physiol*. 2022;600:797–813.
55. Alves-Bezerra M, Cohen DE. Triglyceride metabolism in the liver. *Compr Physiol*. 2017;8:1–8.
56. Yao Z, Gong Y, Chen W, Shao S, Song Y, Guo H, et al. Upregulation of WDR6 drives hepatic de novo lipogenesis in insulin resistance in mice. *Nat Metab*. 2023;5:1706–25.

57. Damasceno de Lima R, Fudoli Lins Vieira R, Rosetto Muñoz V, Chaix A, Azevedo Macedo AP, Calheiros Antunes G, et al. Time-restricted feeding combined with resistance exercise training prevents obesity and improves lipid metabolism in the liver of mice fed a high-fat diet. *Am J Physiol Endocrinol Metab.* 2023;325:E513–28.
58. Kantor MA, Cullinane EM, Herbert PN, Thompson PD. Acute increase in lipoprotein lipase following prolonged exercise. *Metabolism.* 1984;33:454–7.
59. Donnelly KL, Smith CI, Schwarzenberg SJ, Jessurun J, Boldt MD, Parks EJ. Sources of fatty acids stored in liver and secreted via lipoproteins in patients with nonalcoholic fatty liver disease. *J Clin Invest.* 2005;115:1343–51.
60. Zou Y, Chen Z, Sun C, Yang D, Zhou Z, Peng X, et al. Exercise intervention mitigates pathological liver changes in NAFLD zebrafish by activating SIRT1/AMPK/NRF2 signaling. *Int J Mol Sci.* 2021;22:10940.
61. Qian X, Wang T, Gong J, Wang L, Chen X, Lin H, et al. Exercise in mice ameliorates high-fat diet-induced nonalcoholic fatty liver disease by lowering HMGCS2. *Aging (Albany NY).* 2021;13:8960–74.
62. Wang Z, Zhu Y, Xia L, Li J, Song M, Yang C. Exercise-induced ADAR2 protects against nonalcoholic fatty liver disease through miR-34a. *Nutrients.* 2022;15:121.
63. Huber Y, Pfirrmann D, Gebhardt I, Labenz C, Gehrke N, Straub BK, et al. Improvement of non-invasive markers of NAFLD from an individualised, web-based exercise program. *Aliment Pharmacol Ther.* 2019;50:930–9.
64. Schleh MW, Ahn C, Ryan BJ, Chugh OK, Luker AT, Luker KE, et al. Both moderate- and high-intensity exercise training increase intramyocellular lipid droplet abundance and modify myocellular distribution in adults with obesity. *Am J Physiol Endocrinol Metab.* 2023;325:E466–79.
65. Turcotte LP, Fisher JS. Skeletal muscle insulin resistance: Roles of fatty acid metabolism and exercise. *Phys Ther.* 2008;88:1279–96.
66. Goudriaan JR, Dahlmans VE, Teusink B, Ouwens DM, Febbraio M, Maassen JA, et al. CD36 deficiency increases insulin sensitivity in muscle, but induces insulin resistance in the liver in mice. *J Lipid Res.* 2003;44:2270–7.
67. Del Campo JA, Gallego P, Grande L. Role of inflammatory response in liver diseases: Therapeutic strategies. *World J Hepatol.* 2018;10:1–7.
68. Docherty S, Harley R, McAuley JJ, Crowe LAN, Pedret C, Kirwan PD, et al. The effect of exercise on cytokines: Implications for musculoskeletal health: a narrative review. *BMC Sports Sci Med Rehabil.* 2022;14:5.
69. Koelwyn GJ, Wennerberg E, Demaria S, Jones LW. Exercise in regulation of inflammation-immune axis function in cancer initiation and progression. *Oncology (Williston Park).* 2015;29:908–20; 922.
70. Callegari IOM, Rocha GZ, Oliveira AG. Physical exercise, health, and disease treatment: The role of macrophages. *Front Physiol.* 2023;14:1061353.
71. Fredrickson G, Barrow F, Dietsche K, Parthiban P, Khan S, Robert S, et al. Exercise of high intensity ameliorates hepatic inflammation and the progression of NASH. *Mol Metab.* 2021;53:101270.
72. Huber Y, Gehrke N, Biedenbach J, Helmig S, Simon P, Straub BK, et al. Voluntary distance running prevents TNF-mediated liver injury in mice through alterations of the intrahepatic immune milieu. *Cell Death Dis.* 2017;8:e2893.
73. Tsutsui Y, Mori T, Yoshio S, Sato M, Sakata T, Yoshida Y, et al. Exercise changes the intrahepatic immune cell profile and inhibits the progression of nonalcoholic steatohepatitis in a mouse model. *Hepatol Commun.* 2023;7:e0236.
74. Zheng F, Cai Y. Concurrent exercise improves insulin resistance and nonalcoholic fatty liver disease by upregulating PPAR- γ and genes involved in the beta-oxidation of fatty acids in ApoE-KO mice fed a high-fat diet. *Lipids Health Dis.* 2019;18:6.
75. Naimimohasses S, O’Gorman P, Wright C, Ni Fhloinn D, Holden D, Conlon N, et al. Differential effects of dietary versus exercise intervention on intrahepatic MAIT cells and histological features of NAFLD. *Nutrients.* 2022;14:2198.
76. Blériot C, Barreby E, Dunsmore G, Ballaire R, Chakarov S, Ficht X, et al. A subset of Kupffer cells regulates metabolism through the expression of CD36. *Immunity.* 2021;54:2101–116.e2106.
77. Kawanishi N, Mizokami T, Yada K, Suzuki K. Exercise training suppresses scavenger receptor CD36 expression in kupffer cells of nonalcoholic steatohepatitis model mice. *Physiol Rep.* 2018;6:e13902.
78. Gao C, Liu Y, Jiang C, Liu L, Li J, Li D, et al. Intensive running enhances NF- κ B activity in the mice liver and the intervention effects of quercetin. *Nutrients.* 2020;12:2770.
79. Cabral-Santos C, de Lima Junior EA, Fernandes I, Pinto RZ, Rosa-Neto JC, Bishop NC, et al. Interleukin-10 responses from acute exercise in healthy subjects: A systematic review. *J Cell Physiol.* 2019;234:9956–65.
80. Takahashi A, Abe K, Fujita M, Hayashi M, Okai K, Ohira H. Simple resistance exercise decreases cytokeratin 18 and fibroblast growth factor 21 levels in patients with nonalcoholic fatty liver disease: A retrospective clinical study. *Medicine (Baltimore).* 2020;99:e20399.
81. Yu Y, Yu L, Cheng N, Liu X, Fang C, Liu S, et al. Exercise alleviates the apolipoprotein A5-toll-like receptor 4 axis impairment in mice with high-fat diet-induced non-alcoholic steatohepatitis. *Front Physiol.* 2021;12:783341.
82. Shery AP, Willis SA, Yates T, Johnson W, Razieh C, Sargeant JA, et al. Physical activity is inversely associated with hepatic fibro-inflammation: A population-based cohort study using UK Biobank data. *JHEP Rep.* 2023;5:100622.
83. Dhatchinamoorthy K, Colbert JD, Rock KL. Cancer immune evasion through loss of MHC class I antigen presentation. *Front Immunol.* 2021;12:636568.
84. Sudholz H, Delconte RB, Huntington ND. Interleukin-15 cytokine checkpoints in natural killer cell anti-tumor immunity. *Curr Opin Immunol.* 2023;84:102364.
85. Hingorjo MR, Zehra S, Saleem S, Qureshi MA. Serum interleukin-15 and its relationship with adiposity indices before and after short-term endurance exercise. *Pak J Med Sci.* 2018;34:1125–31.
86. Wang Z, Cui Y, Zhang Y, Wang X, Li J, Li J, et al. Twelve-week treadmill endurance training in mice is associated with upregulation of interleukin-15 and natural killer cell activation and increases apoptosis rate in Hepa1-6 cell-derived mouse hepatomas. *Braz J Med Biol Res.* 2023;56:e12296.
87. Ahmed H, Mahmud AR, Faijanur-Rob-Siddiquee M, Shahriar A, Biswas P, Shimul MEK, et al. Role of T cells in cancer immunotherapy: Opportunities and challenges. *Cancer Pathogen Ther.* 2023;1:116–26.
88. Rundqvist H, Veliça P, Barbieri L, Gameiro PA, Bargiela D, Gojkovic M, et al. Cytotoxic T-cells mediate exercise-induced reductions in tumor growth. *Elife.* 2020;9:e59996.
89. Mendes-Braz M, Elias-Miró M, Jiménez-Castro MB, Casillas-Ramírez A, Ramalho FS, Peralta C. The current state of knowledge of hepatic ischemia-reperfusion injury based on its study in experimental models. *J Biomed Biotechnol.* 2012;2012:298657.
90. Zhang Y, Zhang X, Zhang H, Song P, Pan W, Xu P, et al. Mesenchymal stem cells derived extracellular vesicles alleviate traumatic hemorrhagic shock induced hepatic injury via IL-10/PTPN22-mediated M2 Kupffer cell polarization. *Front Immunol.* 2021;12:811164.
91. Rampes S, Ma D. Hepatic ischemia-reperfusion injury in liver transplant setting: Mechanisms and protective strategies. *J Biomed Res.* 2019;33:221–34.
92. Ozaki M. Cellular and molecular mechanisms of liver regeneration: Proliferation, growth, death and protection of hepatocytes. *Semin Cell Dev Biol.* 2020;100:62–73.

93. Batatinha H, Lira F, Kruger K, Rosa Neto J. Physical exercise and metabolic reprogramming. *Essential Aspects of Immuno-metabolism in Health and Disease*. Springer, Cham. 2022; 235–56.
94. Linecker M, Frick L, Kron P, Limani P, Kambakamba P, Tschuor C, et al. Exercise improves outcomes of surgery on fatty liver in mice: A novel effect mediated by the AMPK pathway. *Ann Surg*. 2020; 271:347–55.
95. Liu S, Cui F, Ning K, Wang Z, Fu P, Wang D, et al. Role of irisin in physiology and pathology. *Front Endocrinol (Lausanne)*. 2022;13:962968.
96. Bi J, Yang L, Wang T, Zhang J, Li T, Ren Y, et al. Irisin improves autophagy of aged hepatocytes via increasing telomerase activity in liver injury. *Oxid Med Cell Longev*. 2020;2020:6946037.
97. Zhang J, Ren Y, Bi J, Wang M, Zhang L, Wang T, et al. Involvement of kindlin-2 in irisin's protection against ischaemia reperfusion-induced liver injury in high-fat diet-fed mice. *J Cell Mol Med*. 2020;24:13081–92.
98. Williams FR, Vallance A, Faulkner T, Towey J, Durman S, Kyte D, et al. Home-based exercise in patients awaiting liver transplantation: A feasibility study. *Liver Transpl*. 2019;25: 995–1006.
99. Morkane CM, Kearney O, Bruce DA, Melikian CN, Martin DS. An outpatient hospital-based exercise training program for patients with cirrhotic liver disease awaiting transplantation: A feasibility trial. *Transplantation*. 2020;104:97–103.
100. De Smet S, O'Donoghue K, Lormans M, Monbaliu D, Pengel L. Does exercise training improve physical fitness and health in adult liver transplant recipients? A systematic review and meta-analysis. *Transplantation*. 2023;107:e11–26.
101. Wang T, Yu M, Li H, Qin S, Ren W, Ma Y, et al. FNDC5/irisin inhibits the inflammatory response and mediates the aerobic exercise-induced improvement of liver injury after myocardial infarction. *Int J Mol Sci*. 2023;24:4159.
102. Powell EE, Wong VW, Rinella M. Non-alcoholic fatty liver disease. *Lancet*. 2021;397:2212–24.
103. Sheka AC, Adeyi O, Thompson J, Hameed B, Crawford PA, Ikramuddin S. Nonalcoholic steatohepatitis: A review. *Jama*. 2020;323:1175–83.
104. Huby T, Gautier EL. Immune cell-mediated features of non-alcoholic steatohepatitis. *Nat Rev Immunol*. 2022;22:429–43.
105. Oh S, Tsujimoto T, Kim B, Uchida F, Suzuki H, Iizumi S, et al. Weight-loss-independent benefits of exercise on liver steatosis and stiffness in Japanese men with NAFLD. *JHEP Rep*. 2021;3: 100253.
106. Hari A, Fealy CE, Axelrod CL, Haus JM, Flask CA, McCullough AJ, et al. Exercise training rapidly increases hepatic insulin extraction in NAFLD. *Med Sci Sports Exerc*. 2020;52:1449–55.
107. Hejazi K, Hackett D. Effect of exercise on liver function and insulin resistance markers in patients with non-alcoholic fatty liver disease: A systematic review and meta-analysis of randomized controlled trials. *J Clin Med*. 2023;12:3011.
108. Ezpeleta M, Gabel K, Cienfuegos S, Kalam F, Lin S, Pavlou V, et al. Effect of alternate day fasting combined with aerobic exercise on non-alcoholic fatty liver disease: A randomized controlled trial. *Cell Metab*. 2023;35:56–70.e53.
109. Keating SE, Hackett DA, Parker HM, O'Connor HT, Gerofi JA, Sainsbury A, et al. Effect of aerobic exercise training dose on liver fat and visceral adiposity. *J Hepatol*. 2015;63:174–82.
110. Winn NC, Liu Y, Rector RS, Parks EJ, Ibdah JA, Kanaley JA. Energy-matched moderate and high intensity exercise training improves nonalcoholic fatty liver disease risk independent of changes in body mass or abdominal adiposity—A randomized trial. *Metabolism*. 2018;78:128–40.
111. Sargeant JA, Gray LJ, Bodicoat DH, Willis SA, Stensel DJ, Nimmo MA, et al. The effect of exercise training on intrahepatic triglyceride and hepatic insulin sensitivity: A systematic review and meta-analysis. *Obes Rev*. 2018;19:1446–59.
112. Younossi ZM, Corey KE, Lim JK. AGA clinical practice update on lifestyle modification using diet and exercise to achieve weight loss in the management of nonalcoholic fatty liver disease: Expert review. *Gastroenterology*. 2021;160:912–8.
113. Iwakiri Y. Pathophysiology of portal hypertension. *Clin Liver Dis*. 2014;18:281–91.
114. Villa E, Bianchini M, Blasi A, Denys A, Giannini EG, de Gottardi A, et al. EASL Clinical Practice Guidelines on prevention and management of bleeding and thrombosis in patients with cirrhosis. *J Hepatol*. 2022;76:1151–84.
115. Suk KT. Hepatic venous pressure gradient: Clinical use in chronic liver disease. *Clin Mol Hepatol*. 2014;20:6–14.
116. Rodrigues SG, Mendoza YP, Bosch J. Beta-blockers in cirrhosis: Evidence-based indications and limitations. *JHEP Rep*. 2020;2:100063.
117. García-Pagán JC, Santos C, Barberá JA, Luca A, Roca J, Rodríguez-Roisin R, et al. Physical exercise increases portal pressure in patients with cirrhosis and portal hypertension. *Gastroenterology*. 1996;111:1300–6.
118. Sirisunhirun P, Bandidniyamanon W, Jrerattakon Y, Muang-somboon K, Pramyothin P, Nimanong S, et al. Effect of a 12-week home-based exercise training program on aerobic capacity, muscle mass, liver and spleen stiffness, and quality of life in cirrhotic patients: A randomized controlled clinical trial. *BMC Gastroenterol*. 2022;22:66.
119. Macías-Rodríguez RU, Ibarra-Lomelí H, Ruiz-Margáin A, Ponce-de-León-Rosales S, Vargas-Vorácková F, García-Flores O, et al. Changes in hepatic venous pressure gradient induced by physical exercise in cirrhosis: Results of a pilot randomized open clinical trial. *Clin Transl Gastroenterol*. 2016;7:e180.
120. Berzigotti A, Albillos A, Villanueva C, Genescá J, Ardevol A, Augustin S, et al. Effects of an intensive lifestyle intervention program on portal hypertension in patients with cirrhosis and obesity: The SportDiet study. *Hepatology*. 2017;65:1293–305.
121. Cheemera S, Balakrishnan M. Global epidemiology of chronic liver disease. *Clin Liver Dis (Hoboken)*. 2021;17:365–70.
122. Wu J, Xue X, Fan G, Gu Y, Zhou F, Zheng Q, et al. Ferulic acid ameliorates hepatic inflammation and fibrotic liver injury by inhibiting PTP1B activity and subsequent promoting AMPK phosphorylation. *Front Pharmacol*. 2021;12:754976.
123. Yu JH, Song SJ, Kim A, Choi Y, Seok JW, Kim HJ, et al. Suppression of PPAR γ -mediated monoacylglycerol O-acyltransferase 1 expression ameliorates alcoholic hepatic steatosis. *Sci Rep*. 2016;6:29352.
124. Zhang Y, Wu B, Zhang H, Ge X, Ying S, Hu M, et al. Inhibition of MD2-dependent inflammation attenuates the progression of non-alcoholic fatty liver disease. *J Cell Mol Med*. 2018;22:936–47.
125. Yang W, Liu L, Wei Y, Fang C, Liu S, Zhou F, et al. Exercise suppresses NLRP3 inflammasome activation in mice with diet-induced NASH: A plausible role of adropin. *Lab Invest*. 2021; 101:369–80.
126. Puengel T, De Vos S, Hundertmark J, Kohlhepp M, Guldiken N, Pujuguet P, et al. The medium-chain fatty acid receptor GPR84 mediates myeloid cell infiltration promoting steatohepatitis and fibrosis. *J Clin Med*. 2020;9:1140.
127. Lin D, Sun Q, Liu Z, Pan J, Zhu J, Wang S, et al. Gut microbiota and bile acids partially mediate the improvement of fibroblast growth factor 21 on methionine-choline-deficient diet-induced non-alcoholic fatty liver disease mice. *Free Radic Biol Med*. 2023;195:199–218.
128. Aamann L, Dam G, Jepsen P, Borre M, Drljovic-Nielsen A, Overgaard K, et al. Reduced 3-year risk of hospital admission and mortality after 12-week resistance training of cirrhosis patients: A follow-up of a randomized clinical trial. *J Gastroenterol Hepatol*. 2023;38:1365–71.

129. Namisaki T, Sato S, Yoshiji H. Role of combined aerobic and resistance exercise in liver cirrhosis. *J Gastroenterol.* 2024;59:359–60.
130. Bellar A, Welch N, Dasarathy S. Exercise and physical activity in cirrhosis: Opportunities or perils. *J Appl Physiol.* 1985;2020:1547–67.
131. Bray F, Laversanne M, Sung H, Ferlay J, Siegel RL, Soerjomataram I, et al. Global cancer statistics 2022: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin.* 2024;74:229–63.
132. Llovet JM, Kelley RK, Villanueva A, Singal AG, Pikarsky E, Roayaie S, et al. Hepatocellular carcinoma. *Nat Rev Dis Primers.* 2021;7:6.
133. Matthews CE, Moore SC, Arem H, Cook MB, Trabert B, Håkansson N, et al. Amount and intensity of leisure-time physical activity and lower cancer risk. *J Clin Oncol.* 2020;38:686–97.
134. Wang J, Huang L, Gao Y, Wang Y, Chen S, Huang J, et al. Physically active individuals have a 23% lower risk of any colorectal neoplasia and a 27% lower risk of advanced colorectal neoplasia than their non-active counterparts: Systematic review and meta-analysis of observational studies. *Br J Sports Med.* 2020;54:582–91.
135. Wang Q, Zhou W. Roles and molecular mechanisms of physical exercise in cancer prevention and treatment. *J Sport Health Sci.* 2021;10:201–10.
136. Kang DW, Fairey AS, Boulé NG, Field CJ, Wharton SA, Courneya KS. Effects of exercise on cardiorespiratory fitness and biochemical progression in men with localized prostate cancer under active surveillance: The ERASE Randomized Clinical Trial. *JAMA Oncol.* 2021;7:1487–95.
137. Pedisic Z, Shrestha N, Kovalchik S, Stamatakis E, Liangruenrom N, Grgic J, et al. Is running associated with a lower risk of all-cause, cardiovascular and cancer mortality, and is the more the better? A systematic review and meta-analysis. *Br J Sports Med.* 2020;54:898–905.
138. Moore SC, Lee IM, Weiderpass E, Campbell PT, Sampson JN, Kitahara CM, et al. Association of leisure-time physical activity with risk of 26 types of cancer in 1.44 million adults. *JAMA Intern Med.* 2016;176:816–25.
139. Baumeister SE, Schlesinger S, Aleksandrova K, Jochem C, Jenab M, Gunter MJ, et al. Association between physical activity and risk of hepatobiliary cancers: A multinational cohort study. *J Hepatol.* 2019;70:885–92.
140. Lee J. Associations between physical activity and liver cancer risks and mortality: A systematic review and meta-analysis. *Int J Environ Res Public Health.* 2020;17:8943.
141. Chang KV, Chen JD, Wu WT, Huang KC, Hsu CT, Han DS. Association between loss of skeletal muscle mass and mortality and tumor recurrence in hepatocellular carcinoma: A systematic review and meta-analysis. *Liver Cancer.* 2018;7:90–103.
142. Sweegers MG, Altenburg TM, Brug J, May AM, van Vulpen JK, Aaronson NK, et al. Effects and moderators of exercise on muscle strength, muscle function and aerobic fitness in patients with cancer: A meta-analysis of individual patient data. *Br J Sports Med.* 2019;53:812.
143. Zhang QB, Zhang BH, Zhang KZ, Meng XT, Jia QA, Zhang QB, et al. Moderate swimming suppressed the growth and metastasis of the transplanted liver cancer in mice model: With reference to nervous system. *Oncogene.* 2016;35:4122–31.
144. Zhao T, Guo BJ, Xiao CL, Chen JJ, Lü C, Fang FF, et al. Aerobic exercise suppresses hepatocellular carcinoma by downregulating dynamin-related protein 1 through PI3K/AKT pathway. *J Integr Med.* 2021;19:418–27.
145. Arfianti A, Pok S, Barn V, Haigh WG, Yeh MM, Ioannou GN, et al. Exercise retards hepatocarcinogenesis in obese mice independently of weight control. *J Hepatol.* 2020;73:140–8.
146. Piguet AC, Saran U, Simillion C, Keller I, Terracciano L, Reeves HL, et al. Regular exercise decreases liver tumors development in hepatocyte-specific PTEN-deficient mice independently of steatosis. *J Hepatol.* 2015;62:1296–303.
147. Liu XF, Zhu XD, Feng LH, Li XL, Xu B, Li KS, et al. Physical activity improves outcomes of combined lenvatinib plus anti-PD-1 therapy in unresectable hepatocellular carcinoma: A retrospective study and mouse model. *Exp Hematol Oncol.* 2022;11:20.
148. Sheinboim D, Parikh S, Manich P, Markus I, Dahan S, Parikh R, et al. An exercise-induced metabolic shield in distant organs blocks cancer progression and metastatic dissemination. *Cancer Res.* 2022;82:4164–78.
149. Rogers CJ, Colbert LH, Greiner JW, Perkins SN, Hursting SD. Physical activity and cancer prevention: Pathways and targets for intervention. *Sports Med.* 2008;38:271–96.
150. Ligibel JA, Bohlke K, May AM, Clinton SK, Demark-Wahnefried W, Gilchrist SC, et al. Exercise, diet, and weight management during cancer treatment: ASCO guideline. *J Clin Oncol.* 2022;40:2491–507.
151. Schmitz KH, Campbell AM, Stuiver MM, Pinto BM, Schwartz AL, Morris GS, et al. Exercise is medicine in oncology: Engaging clinicians to help patients move through cancer. *CA Cancer J Clin.* 2019;69:468–84.
152. Campbell KL, Winters-Stone KM, Wiskemann J, May AM, Schwartz AL, Courneya KS, et al. Exercise guidelines for cancer survivors: Consensus statement from International Multidisciplinary Roundtable. *Med Sci Sports Exerc.* 2019;51:2375–90.
153. Liu PZ, Nusslock R. Exercise-mediated neurogenesis in the hippocampus via BDNF. *Front Neurosci.* 2018;12:52.
154. He XF, Liu DX, Zhang Q, Liang FY, Dai GY, Zeng JS, et al. Voluntary exercise promotes glymphatic clearance of amyloid beta and reduces the activation of astrocytes and microglia in aged mice. *Front Mol Neurosci.* 2017;10:144.
155. Lin TW, Tsai SF, Kuo YM. Physical exercise enhances neuroplasticity and delays Alzheimer's disease. *Brain Plast.* 2018;4:95–110.
156. Nijholt KT, Sánchez-Aguilera PI, Voorrips SN, de Boer RA, Westenbrink BD. Exercise: A molecular tool to boost muscle growth and mitochondrial performance in heart failure? *Eur J Heart Fail.* 2022;24:287–98.
157. Wang L, Tu W, Li X, Li C, Lu J, Dai P, et al. Exercise improves cardiac function and attenuates myocardial inflammation and apoptosis by regulating APJ/STAT3 in mice with stroke. *Life Sci.* 2023;332:122041.
158. Zhang X, Zhao Y, Guo D, Luo M, Zhang Q, Zhang L, et al. Exercise improves heart function after myocardial infarction: The merits of AMPK. *Cardiovasc Drugs Ther.* 2024. Online ahead of print.
159. Hargreaves M, Spriet LL. Skeletal muscle energy metabolism during exercise. *Nat Metab.* 2020;2:817–28.
160. Laurens C, Bergouignan A, Moro C. Exercise-released myokines in the control of energy metabolism. *Front Physiol.* 2020;11:91.
161. Bao W, Sun Y, Zhang T, Zou L, Wu X, Wang D, et al. Exercise programs for muscle mass, muscle strength and physical performance in older adults with sarcopenia: A systematic review and meta-analysis. *Aging Dis.* 2020;11:863–73.
162. Mika A, Macaluso F, Barone R, Di Felice V, Sledzinski T. Effect of exercise on fatty acid metabolism and adipokine secretion in adipose tissue. *Front Physiol.* 2019;10:26.
163. Fu P, Zhu R, Jia J, Hu Y, Wu C, Cieszczyk P, et al. Aerobic exercise promotes the functions of brown adipose tissue in obese mice via a mechanism involving COX2 in the VEGF signaling pathway. *Nutr Metab.* 2021;18:56.
164. Vechetti IJ Jr, Peck BD, Wen Y, Walton RG, Valentino TR, Alimov AP, et al. Mechanical overload-induced muscle-derived extracellular vesicles promote adipose tissue lipolysis. *Faseb j.* 2021;35:e21644.

165. Monda V, Villano I, Messina A, Valenzano A, Esposito T, Moscatelli F, et al. Exercise modifies the gut microbiota with positive health effects. *Oxid Med Cell Longev*. 2017;2017:3831972.
166. Carbajo-Pescador S, Porras D, García-Mediavilla MV, Martínez-Flórez S, Juárez-Fernández M, Cuevas MJ, et al. Beneficial effects of exercise on gut microbiota functionality and barrier integrity, and gut-liver crosstalk in an in vivo model of early obesity and non-alcoholic fatty liver disease. *Dis Model Mech*. 2019;12:dmm039206.
167. Khalid K, Szewczyk A, Kiszalkiewicz J, Migdalska-Sęk M, Domańska-Senderowska D, Brzeziański M, et al. Type of training has a significant influence on the GH/IGF-1 axis but not on regulating miRNAs. *Biol Sport*. 2020;37:217–28.
168. Dichtel LE, Corey KE, Misdraji J, Bredella MA, Schorr M, Osganian SA, et al. The association between IGF-1 levels and the histologic severity of nonalcoholic fatty liver disease. *Clin Transl Gastroenterol*. 2017;8:e217.
169. Zhu W, Sahar NE, Javaid HMA, Pak ES, Liang G, Wang Y, et al. Exercise-induced irisin decreases inflammation and improves NAFLD by competitive binding with MD2. *Cells*. 2021;10:3306.
170. Cheng R, Wang L, Le S, Yang Y, Zhao C, Zhang X, et al. A randomized controlled trial for response of microbiome network to exercise and diet intervention in patients with nonalcoholic fatty liver disease. *Nat Commun*. 2022;13:2555.
171. Solé C, Guilly S, Da Silva K, Llopis M, Le-Chatelier E, Huelin P, et al. Alterations in gut microbiome in cirrhosis as assessed by quantitative metagenomics: Relationship with acute-on-chronic liver failure and prognosis. *Gastroenterology*. 2021;160:206–218.e213.

How to cite this article: Zhang Y, Cao C, Li C, Witt RG, Huang H, Tsung A, et al. Physical exercise in liver diseases. *Hepatology*. 2026;83:915–930. <https://doi.org/10.1097/HEP.0000000000000941>